

•) Exercícios 17/09•) Linearização de  $f(\theta) = \cos \theta$ - Em torno de  $\theta_0 = 0$ 

$$\cos \theta = \cos \theta_0 + \left. \frac{\partial (\cos \theta)}{\partial \theta} \right|_{\theta_0} \cdot (\theta - \theta_0) = 1 - \sin \theta_0 \cdot (\theta - \theta_0) = \underline{1}$$

- Em torno de  $\theta_0 = \pi/4$ 

$$\cos \theta = \cos \theta_0 + \left. \frac{\partial (\cos \theta)}{\partial \theta} \right|_{\theta_0} \cdot (\theta - \theta_0) = \frac{\sqrt{2}}{2} - \sin(\pi/4) \cdot (\theta - \pi/4)$$

$$\Rightarrow \cos \theta = \frac{\sqrt{2}}{2} - \frac{\sqrt{2}}{2} \cdot (\theta - \pi/4) = \underline{\underline{-\frac{\sqrt{2}}{2} \cdot \theta + \frac{\sqrt{2}(4+\pi)}{8}}}$$

•) Linearização por expansão em série de Taylor

$$F = m \cdot \vec{r} + m \cdot \vec{r} \cdot \vec{r} = m \cdot \vec{r} \cdot \vec{r} ; \quad \vec{r} = \vec{r} = \vec{r} = \vec{F} = 0;$$

$$m \cdot \vec{r} = f(F, r, \dot{r}, \ddot{r}, \ddot{r}) = F - m \cdot \vec{r} \cdot \vec{r} + m \cdot \vec{r} \cdot \dot{r}$$

Série de Taylor:

$$f(F, r, \dot{r}, \ddot{r}, \ddot{r}) = f(\bar{F}, \bar{r}, \bar{\dot{r}}, \bar{\ddot{r}}, \bar{\ddot{r}}) + \frac{\partial f}{\partial F} \cdot (F - \bar{F}) + \frac{\partial f}{\partial r} \cdot (r - \bar{r}) + \frac{\partial f}{\partial \dot{r}} \cdot (\dot{r} - \bar{\dot{r}}) + \frac{\partial f}{\partial \ddot{r}} \cdot (\ddot{r} - \bar{\ddot{r}})$$

Desenvolvendo cada termo:

$$f(\bar{F}, \bar{\theta}, \bar{n}, \bar{i}, \bar{\alpha}) = 0$$

$$\frac{\partial f}{\partial F} \cdot (F - \bar{F}) = F$$

$$\frac{\partial f}{\partial \theta} (\theta - \bar{\theta}) = -m \cdot \bar{n}^0 \cdot (\theta - \bar{\theta}) = 0$$

$$\frac{\partial f}{\partial n} (n - \bar{n}) = -m \cdot \bar{\theta} \cdot (n - 0) = -m \cdot \bar{\theta} \cdot n$$

$$\frac{\partial f}{\partial i} (i - \bar{i}) = +m \cdot \bar{\alpha} \cdot (i - 0) = +m \cdot \bar{\alpha} \cdot i$$

$$\frac{\partial f}{\partial \alpha} (\alpha - \bar{\alpha}) = m \cdot \bar{i} \cdot (\alpha - \bar{\alpha}) = 0$$

Nessa forma:

$$\underline{f(F, \theta, n, i, \alpha) = m \cdot \dot{\theta} = F - m \bar{\theta} \cdot n + m \bar{\alpha} \cdot i}$$