

Microarqueología – ARQ5095

Profa. Dra. Ximena S. Villagrán

Microarqueobotánica

Dra. Jennifer Watling (MAE/USP) - docente convidada

Bibliografía básica

Piperno DR. Phytoliths: A comprehensive guide for archaeologists and paleoecologists. Oxford: Altimira Press; 2006.

Pearsall DM. Paleoethnobotany: A handbook of procedures. 3rd ed. New York: Academic Press; 2015.

Torrence R, Barton H, editors. Ancient Starch Research. Walnut Creek, California: Left Coast Press Inc.; 2006.

Bibliografía citada na aula

Babot M del P. Starch grain damage as an indicator of food processing. In: Hart DM, Wallis LA, editors. Phytolith and Starch Research in the Australian-Pacific-Asian regions: the state of the art. Canberra, Australia: Pandanus Books, Research School of Pacific and Asian Studies, The Australian National University; 2003. p. 69–81.

Bremond L, Alexandre A, Wooller MJ, Hély C, Williamson D, Schäfer P a., et al. Phytolith indices as proxies of grass subfamilies on East African tropical mountains. *Glob Planet Change* 2008; 61(3–4):209–24.

Carson JF, Watling J, Mayle FE, Whitney BS, Iriarte J, Prumers H, et al. Pre-Columbian land use in the ring-ditch region of the Bolivian Amazon. *The Holocene*. 2015; 25(8):1285–300.

Crowther A, Haslam M, Oakden N, Walde D, Mercader J. Documenting contamination in ancient starch laboratories. *J Archaeol Sci*, 2014; 49(1):90–104

Harvey EL, Fuller DQ. Investigating crop processing using phytolith analysis: the example of rice and millets. 2005; 32(5):739–52.

Henry AG, Hudson HF, Piperno DR. Changes in starch grain morphologies from cooking. *J Archaeol Sci*; 2009; 36(3):915–22.

Henry AG, Brooks AS, Piperno DR. Microfossils in calculus demonstrate consumption of plants and cooked foods in Neanderthal diets (Shanidar III, Iraq; Spy I and II, Belgium). *Proc Natl Acad Sci*. 2011; 108(2):486–91.

Hodson MJ. The development of phytoliths in plants and its influence on their chemistry and isotopic composition. Implications for palaeoecology and archaeology. *J Archaeol Sci*; 2016; 68:62–9.

Holst I, Moreno JE, Piperno DR. Identification of teosinte, maize, and *Tripsacum* in Mesoamerica by using pollen, starch grains, and phytoliths. *Proc Natl Acad Sci*. 2007; 104(45):17608–13.

Iriarte J, Glaser B, Watling J, Wainwright A, Birk JJ, Renard D, et al. Late Holocene Neotropical agricultural landscapes: phytolith and stable carbon isotope analysis of raised fields from French Guianan coastal savannahs. *J Archaeol Sci*; 2010; 37(12):2984–94.

Iriarte J, Dickau R. ¿Las culturas del maíz? Arqueobotânica de las sociedades hidráulicas de las tierras bajas sudamericanas. *Amazôntica*. 2012; 4:30–58.

Madella M, Lancelotti C. Taphonomy and phytoliths: a user manual. *Quat Int and INQUA*; 2012; 275:76–83.

Morcote-Ríos G, Bernal R, Raz L. Phytoliths as a tool for archaeobotanical, palaeobotanical and palaeoecological studies in Amazonian palms. *Bot J Linn Soc*. 2016; 182(2):348–60

Pagán-Jiménez JR, Guachamín-Tello AM, Romero-Bastidas ME, Constantine-Castro AR. Late ninth millennium B.P. use of *Zea mays* L. at Cubilán area, highland Ecuador, revealed by ancient starches. *Quat Int*. 2016; 404:137–55.

Perry L. Starch analyses reveal multiple functions of quartz “manioc” grater flakes from the Orinoco Basin, Venezuela. *Interciencia*. 2002; 27(11):635–9.

Perry L, Dickau R, Zarrillo S, Holst I, Pearsall DM, Piperno DR, et al. Starch fossils and the domestication and dispersal of chili peppers (*Capsicum* spp. L.) in the Americas. *Science* (80-. 2007; 315(5814):986–8.

Pearsall DM, Piperno DR, Dinan EH, Umlauf M, Zhao Z, Benfer RA. Distinguishing rice (*Oryza sativa* Poaceae) from wild *Oryza* species through phytolith analysis: results of preliminary research. *Econ Bot*. 1995; 49(2):183–96.

Piperno DR. Identifying crop plants with phytoliths (and starch grains) in Central and South America: A review and an update of the evidence. *Quat Int*. 2009; 193(1–2):146–59.

Piperno DR. The origins of plant cultivation and domestication in the New World tropics: patterns, process, and new developments. *Curr Anthropol*. 2011; 52(S4):S453–70.

Piperno DR, Ranere a J, Holst I, Hansell P. Starch grains reveal early root crop horticulture in the Panamanian tropical forest. *Nature*. 2000 ;407(6806):894–7.

Piperno DR, Stothert KE. Phytolith evidence for early Holocene Cucurbita domestication in southwest Ecuador. *Science* (80-) 2003; 299(5609):1054–7.

Pagán-Jiménez JR, Guachamín-Tello AM, Romero-Bastidas ME, Constantine-Castro AR. Late ninth millennium B.P. use of *Zea mays* L. at Cubilán area, highland Ecuador, revealed by ancient starches. *Quat Int.* 2016; 404:137–55.

Twiss PC. Predicted world distribution of C3 and C4 grass phytoliths. In: Rapp Jr. G, Mulholland SC, editors. *Phytolith Systematics: Emerging Issues*. New York: Plenum Press; 1992. p. 113–28.

Watling J, Iriarte J, Whitney BS, Consuelo E, Mayle FE, Castro W, et al. Differentiation of neotropical ecosystems by modern soil phytoliths assemblages and its implications for palaeoenvironmental and archaeological reconstructions II: southwestern Amazonian forests. *Rev Palaeobot Palynol.* 2016; 226:30–43.

Watling J, Iriarte J, Mayle FE, Schaan D, Pessenda LCR, Loader NJ, et al. Impact of pre-Columbian “geoglyph” builders on Amazonian forests. *Proc Natl Acad Sci.* 2017; 114(8):1868–73.

Wesolowski V, De Souza SMFM, Reinhard KJ, Ceccantini G. Evaluating microfossil content of dental calculus from Brazilian sambaquis. *J Archaeol Sci.* 2010; 37:1326–38.

Tutorial p/ análises Arqueobotânicas:

Charles, M., Crowther, A., Ertug, F., Herbig, C., Jones, G., Kutterer, J., Longford, C., Madella, M., Maier, U., Out, W., Pessin, H., Zurro, D., (2009) Archaeobotanical Online Tutorial <http://archaeobotany.dept.shef.ac.uk/wiki/index.php/Main_Page>

ICSN 2011, The International Code for Starch Nomenclature
<<http://fossilfarm.org/ICSN/Code.html>>

Madella M, Alexandre A, Ball T. International code for phytolith nomenclature 1.0. *Ann Bot.* 2005; 96(2):253–60.