



Published in final edited form as:

Man Ther. 2009 October ; 14(5): 531–538. doi:10.1016/j.math.2008.09.001.

The Mechanisms of Manual Therapy in the Treatment of Musculoskeletal Pain: A Comprehensive Model

Joel E Bialosky, PT, MS¹, Mark D Bishop, PT, PhD¹, Don D Price, PhD², Michael E Robinson, PhD³, and Steven Z George, PT, PhD¹

¹University of Florida Department of Physical Therapy, Gainesville, Florida

²University of Florida Department of Dentistry, Gainesville, Florida

³University of Florida Department of Clinical and Health Psychology, Gainesville, Florida

Abstract

Prior studies suggest manual therapy (MT) as effective in the treatment of musculoskeletal pain; however, the mechanisms through which MT exerts its effects are not established. In this paper we present a comprehensive model to direct future studies in MT. This model provides visualization of potential individual mechanisms of MT that the current literature suggests as pertinent and provides a framework for the consideration of the potential interaction between these individual mechanisms. Specifically, this model suggests that a mechanical force from MT initiates a cascade of neurophysiological responses from the peripheral and central nervous system which are then responsible for the clinical outcomes. This model provides clear direction so that future studies may provide appropriate methodology to account for multiple potential pertinent mechanisms.

Keywords

Manual Therapy; Rehabilitation; Pain

Introduction

Available evidence suggests manual therapy (MT) as effective in the treatment of musculoskeletal disorders including low back pain (Childs et al., 2004; Licciardone et al., 2003), carpal tunnel syndrome (Akalin et al., 2002; Rozmaryn et al., 1998), knee osteoarthritis (Deyle et al., 2000), and hip osteoarthritis (MacDonald et al., 2006). Moreover, recent studies have provided even stronger evidence when participants are classified into sub-groups (Childs et al., 2004; Cleland et al., 2006). Despite the literature supporting its effectiveness, the mechanisms of MT are not established leading to a National Institutes of Health (NIH) call to specifically address this shortcoming (Khalsa et al., 2006).

A better understanding of the mechanisms of MT is necessary for several reasons. First, recent evidence suggests successful outcomes in MT are dependent on identifying individuals likely

© 2009 Elsevier Ltd. All rights reserved.

Address Correspondence to: Joel E Bialosky University of Florida, Department of Physical Therapy PO Box 100154 Gainesville, Florida 32610-0154 bialosky@phhp.ufl.edu Phone: 352-870-9116 Fax: 352-273-6109.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

to respond rather than identification of a specific lesion. Subsequently, clinical prediction rules based on clusters of signs and symptoms have been proposed to identify responders to MT (Flynn et al., 2002; Cleland et al., 2007). While helpful in directing clinical practice, an explanation is lacking as to why such patterns of signs and symptoms predicts successful clinical outcomes. Subsequently, the biological plausibility of current clinical prediction rules may not be established leading to concern for chance associations rather than causation. Highlighting this concern, only one clinical prediction rule (Flynn et al., 2002) has, to our knowledge, been validated with a follow up study (Childs et al., 2004). An understanding of the mechanisms behind MT could assist in the identification of individuals likely to respond to MT by allowing a priori hypotheses as to pertinent predictive factors for future clinical prediction rules and a better understanding of the factors which are determined as predictive.

A second benefit of the identification of MT mechanisms is the potential for increased acceptance of these techniques by health care providers. Despite the literature supporting the effectiveness of MT in specific musculoskeletal conditions, health care practitioners at times provide or refer for MT at a lower than expected rate (Bishop & Wing, 2003; Jette & Delitto, 1997; Li & Bombardier, 2001). The lack of an identifiable mechanism of action for MT may limit the acceptability of these techniques as they may be viewed as less scientific. Knowledge of mechanisms may promote more appropriate use of MT by healthcare providers.

The intention of this manuscript is to present a comprehensive model to guide future studies of MT mechanisms. For our purposes, MT includes a variety of techniques used in clinical practice for the treatment of musculoskeletal pain which target the skeletal system, soft tissue, and nervous system (Table 1).

Need for a Comprehensive Model

MT likely works through biomechanical and/or neurophysiological mechanisms. A limitation of the current literature is the failure to acknowledge the potential for a combined effect of these mechanisms. For example, prior studies have noted individual biomechanical (Coppiters & Butler, 2007; Gal et al., 1997) and neurophysiological effects (DeVocht et al., 2005; Dishman & Bulbulian, 2000; Suter et al., 1999; Vicenzino et al., 1998) associated with MT; however the potential interaction of these effects is frequently overlooked. Combined effects may be important to consider as the biomechanical parameters of a given MT may produce unique or dose dependent neurophysiological responses. For example, associated hypoalgesic response (McLean et al., 2002) and EMG response (Colloca et al., 2006) have an observed dependence on the force and force/time profile of a given MT. Additionally, prior studies often focus on a single neurophysiological mechanism without consideration for competing explanations. For example neuromuscular changes such as decreased resting EMG activity (DeVocht et al., 2005) and decreased muscle inhibition (Suter et al., 1999; Suter & McMorland, 2002) have been associated with MT and theorized to occur due to stimulation of the mechanoreceptors or proprioceptors producing a spinal cord mediated effect (Suter et al., 2000; Suter & McMorland, 2002). While helpful in establishing the groundwork for the mechanistic study of MT, conclusions based on studies designed in this fashion may fail to consider other potentially pertinent mechanisms. Psychological factors have an observed association with muscular response in individuals with low back pain (Thomas et al., 2008) and MT has an observed effect on these psychological factors (Williams et al., 2007). Subsequently, outcomes reported in the prior studies (DeVocht et al., 2005; Suter et al., 1999; Suter & McMorland, 2002) could be explained by a descending supraspinal mediating effect due to changes in psychological factors such as fear. A consideration of the interaction between biomechanical and multiple potential neurophysiological effects necessitates a comprehensive model to synthesize the current literature and direct future research.

Proposed Model

We propose the following model which provides a compilation of the existing mechanistic literature of MT as a framework for interpreting current and conducting future mechanistic research (Figure 1). Briefly, this model suggests a mechanical stimulus initiates a number of potential neurophysiological effects which produce the clinical outcomes associated with MT in the treatment of musculoskeletal pain.

Mechanical Stimulus

Biomechanical effects are associated with MT as motion has been quantified with joint biased MT (Colloca et al., 2006;Gal et al., 1997) and nerve biased MT (Coppieters & Alshami, 2007;Coppieters & Butler, 2007); however, the direct implication on clinical outcomes is questionable. First, only transient biomechanical effects are supported by studies which quantify motion (Colloca et al., 2006;Gal et al., 1997;Coppieters & Butler, 2007;Coppieters & Alshami, 2007) but not a lasting positional change (Tullberg et al., 1998;Hsieh et al., 2002). Second, biomechanical assessment is not reliable. Palpation for position and movement faults has demonstrated poor reliability (Seffinger et al., 2004;Trojanovich et al., 1998) suggesting an inability to accurately determine a specific area requiring MT. Third, MT techniques lack precision as nerve biased techniques are not specific to a single nerve (Kleinrensink et al., 2000) and joint biased technique forces are dissipated over a large area (Herzog et al., 2001;Ross et al., 2004). Additionally, different kinetic parameters are observed between clinicians in the performance of the same technique (Hessell et al., 1990;Ngan et al., 2005) and the choice of technique does not seem to matter as much as identifying an individual likely to respond (Cleland et al., 2006;Kent et al., 2005). Finally, studies have reported improvements in signs and symptoms away from the site of application such as treating cervical pain with MT directed to the thoracic spine (Cleland et al., 2005;Cleland et al., 2007) and lateral epicondylitis with MT directed to the cervical spine (Vicenzino et al., 1996). Collectively, the literature suggests a biomechanical effect of MT; however, lasting structural changes have not been identified, clinicians are unable to reliably identify areas requiring MT, the forces associated with MT are not specific to a given location and vary between clinicians, choice of technique does not seem to affect outcomes, and sign and symptom responses occur in areas separate from the region of application. The effectiveness of MT despite the inconsistencies associate with a purported biomechanical mechanism suggests that additional mechanisms may be pertinent. Subsequently, we suggest, that as illustrated by the model, a mechanical force is necessary to initiate a chain of neurophysiological responses which produce the outcomes associated with MT.

Neurophysiological Mechanism

The proposed model accounts for the complex interactions of both the peripheral and central nervous system which comprise the pain experience. Current mechanistic studies of MT in humans are frequently unable to directly observe the central or peripheral nervous system. Subsequently, in the absence of direct observation, conclusions are drawn from associated neurophysiological responses which indirectly implicate specific mechanisms. Studies have measured associated responses of hypoalgesia and sympathetic activity following MT to suggest a mechanism of action mediated by the periaqueductal gray (Wright, 1995) and lessening of temporal summation following MT to suggest a mechanism mediated by the dorsal horn of the spinal cord (George et al., 2006) The model makes use of directly measurable associated responses to imply specific neurophysiological mechanisms when direct observations are not possible. The model categorizes neurophysiological mechanisms as those likely originating from a peripheral mechanism, spinal cord mechanisms, and/or supraspinal mechanisms.

Peripheral mechanism

Musculoskeletal injuries induce an inflammatory response in the periphery which initiates the healing process and influences pain processing. Inflammatory mediators and peripheral nociceptors interact in response to injury and MT may directly affect this process. For example, (Teodorczyk-Injeyan et al., 2006) observed a significant reduction of blood and serum level cytokines in individuals receiving joint biased MT which was not observed in those receiving sham MT or in a control group. Additionally, changes of blood levels of β -endorphin, anandamide, N-palmitoylethanolamide, serotonin, (Degenhardt et al., 2007) and endogenous cannabinoids (McPartland et al., 2005) have been observed following MT. Finally, soft tissue biased MT has been shown to alter acute inflammation in response to exercise (Smith et al., 1994) and substance P levels in individuals with fibromyalgia (Field et al., 2002). Collectively, these studies suggest a potential mechanism of action of MT on musculoskeletal pain mediated by the peripheral nervous system for which mechanistic studies may wish to account.

Spinal mechanisms

MT may exert an effect on the spinal cord. For example, MT has been suggested to act as a counter irritant to modulate pain (Boal & Gillette, 2004) and joint biased MT is speculated to “bombard the central nervous system with sensory input from the muscle proprioceptors (Pickar & Wheeler, 2001).” Subsequently, a spinal cord mediated mechanism of MT must be considered and is accounted for in the model. Direct evidence for such an effect comes from a study (Malisza et al., 2003b) in which joint biased MT was applied to the lower extremity of rats following capsaicin injection. A spinal cord response was quantified by functional MRI during light touch to the hind paw. A trend was noted towards decreased activation of the dorsal horn of the spinal cord following the MT. The model uses associated neuromuscular responses following MT to provide indirect evidence for a spinal cord mediated mechanism. For example, MT is associated with hypoalgesia (George et al., 2006; Mohammadian et al., 2004; Vicenzino et al., 2001), afferent discharge (Colloca et al., 2000; Colloca et al., 2003), motoneuron pool activity (Bulbulian et al., 2002; Dishman & Burke, 2003), and changes in muscle activity (Herzog et al., 1999; Symons et al., 2000) all of which may indirectly implicate a spinal cord mediated effect.

Supraspinal mechanisms

Finally, the pain literature suggests the influence of specific supraspinal structures in response to pain. Structures such as the anterior cingulate cortex (ACC), amygdala, periaqueductal gray (PAG), and rostral ventromedial medulla (RVM) are considered instrumental in the pain experience. (Peyron et al., 2000; Vogt et al., 1996; Derbyshire et al., 1997; Iadarola et al., 1998; Hsieh et al., 1995; Oshiro et al., 2007; Moulton et al., 2005; Staud et al., 2007; Bee & Dickenson, 2007; Guo et al., 2006). Subsequently, the model considers potential supraspinal mechanisms of MT. Direct support for a supraspinal mechanism of action of MT comes from (Malisza et al., 2003a) who applied joint biased MT to the lower extremity of rats following capsaicin injection. Functional MRI of the supraspinal region quantified the response of the hind paw to light touch following the injection. A trend was noted towards decreased activation of the supraspinal regions responsible for central pain processing. The model accounts for direct measures of supraspinal activity along with associated responses such as autonomic responses (Moulton & Watson, 2006; Sterling et al., 2001; Vicenzino et al., 1998) (Delaney et al., 2002; Zhang et al., 2006), and opioid responses (Vernon et al., 1986) (Kaada & Torsteinbo, 1989) to indirectly imply a supraspinal mechanism. Additionally, variables such as placebo, expectation, and psychosocial factors may be pertinent in the mechanisms of MT (Ernst, 2000; Kaptchuk, 2002). For example expectation for the effectiveness of MT is associated with functional outcomes (Kalauokalani et al., 2001) and a recent systematic review of the literature has noted that joint biased MT is associated with improved psychological outcomes (Williams

et al., 2007). For this paper we categorize such factors as neurophysiological effects related to supraspinal descending inhibition due to associated changes in the opioid system (Sauro & Greenberg, 2005), dopamine production (Fuente-Fernandez et al., 2006), and central nervous system (Petrovic et al., 2002; Wager et al., 2004; Matre et al., 2006) which have been observed in studies unrelated to MT.

Implementation of Comprehensive Model

This comprehensive model delineates potential mechanisms associated with pain relief from MT allowing researchers to identify domains of interest their studies are designed to evaluate and potential mechanisms not adequately considered. The model is intended to highlight differing possibilities when conclusions are drawn which may be further explored in subsequent studies. For example, studies have reported hypoalgesia following MT (George et al., 2006; Mohammadian et al., 2004). (George et al., 2006) suggested a spinal cord mediated mechanism due to associated hypoalgesia of temporal summation. The model indicates that while monitoring a spinal cord mediating effect (temporal summation), the potential for a peripheral or supraspinal mediating effects was not considered (Figure 2). A recent study attempted to replicate these prior findings while accounting for potential supraspinal influence (Bialosky et al., 2008). Specifically, a spinal cord mediated effect was measured through an associated response of temporal summation. Additionally, a potential supraspinal mechanism (expectation) was manipulated by randomly assigning participants to receive an instructional set stating MT was expected to either increase, decrease, or have no effect on their pain perception. The model pathway of this study is visualized in Figure 3.

In addition to guiding research, the model also allows clinicians to visualize the potential multiple mechanisms likely involved in the clinical effects of MT. The clinical use of MT is frequently dependent upon a purported biomechanical mechanism in evaluation and treatment. For instance, a clinical examination may focus on locating a mal- aligned joint or a hypomobile joint or soft tissue. A MT technique may then be used as treatment to impart a specific movement to the observed dysfunction. Clinical outcomes are then attributed to alleviation of the biomechanical fault. Such practice is common and has lead to many continuing education dollars and valuable clinic time spent in search of biomechanical dysfunction of questionable validity (Seffinger et al., 2004) and treatments of questionable specificity (Ross et al., 2004). The model provides visualization of what the current literature suggests as mechanisms pertinent to MT and while acknowledging a biomechanical effect allows clinicians to consider other potential mechanisms in the MT evaluation and treatment of individuals with musculoskeletal pain.

Limitations of Proposed Model

The model is intended to be applicable to all forms of MT. While the biomechanical application of joint biased, soft tissue biased and nerve biased MT are different, the related neurophysiological responses are similar and adequately encompassed within the model given the current state of knowledge. The proposed model provides a platform to empirically test hypotheses related to different biomechanical and neurophysiological effects specific to types of MT, an area that is currently lacking in the literature. The proposed comprehensive model is intended to explain the mechanisms of MT on musculoskeletal pain. MT has a postulated role in the treatment of disorders of other body systems such as asthma (Balon & Mior, 2004) and high blood pressure (Plaughter & Bachman, 1993); however, those effects are beyond the scope of the current model. Finally, this model is strictly intended to guide research questions regarding the mechanisms of MT. A body of literature already exists suggesting the effectiveness of MT. The proposed model is intended to compliment and provide underlying explanations to the existing body of literature suggesting the effectiveness of MT.

Future Directions

A limitation in the current literature is the failure to account for the non-specific mechanisms associated with MT in the treatment of musculoskeletal pain. A number of neurophysiological responses associated with MT are also associated with non-specific effects such as placebo (Figure 4). Current study designs have not adequately accounted for non-specific effects, and subsequently, their role in the clinical outcomes associated with MT is unknown. Future mechanistic studies in MT should consider determining the influence of non-specific effects. The model presents a guide to design future mechanistic studies so that all relevant possibilities are included.

The model is based primarily on associated responses as the current body of mechanistic literature is lacking in studies which directly observe regions of interest. As technology improves, the means to directly observe specific regions is becoming possible. More recent studies in the acupuncture literature have reported direct observation of the spinal cord (Wang et al., 2006; Chen et al., 2007) and supraspinal centers (Fang et al., 2008; Dougherty et al., 2008) in response to treatment. Similar studies are possible in MT and will allow direct observation of the nervous system response to MT with a subsequent improved understanding of where the techniques exert their effect.

Interdisciplinary collaboration has been recommended in the study of the mechanisms of MT (Khalsa et al., 2006). The comprehensive model provides a framework for such efforts to study both specific sections of the model and their interaction. For example, a team of researchers could work together including a manual therapist to provide treatment, a biomechanist to monitor the biomechanical parameters of the studied MT, an endocrinologist to monitor peripheral inflammatory mediators, a neurophysiologist to monitor potential spinal cord and supraspinal mechanisms, and a psychologist to monitor the influence of non-specific effects such as expectation, fear, and catastrophizing.

Conclusion

The mechanisms behind the clinical effectiveness of MT are not established. Limitations of prior mechanistic studies are the study of individual mechanisms without regard for others and a failure to adequately account for non-specific effects. We have proposed a comprehensive model to consolidate the current research and guide future research into the mechanisms of MT.

Acknowledgments

The project was supported by Grant Number R-21 AT002796-01 from the National Institutes of Health – National Center for Complimentary and Alternative Medicine (SZG, MDB, MER, DDP).

This manuscript was written while JEB received support from the National Institutes of Health T-32 Neural Plasticity Research Training Fellowship (T32HD043730)

Reference List

- Akalin E, El O, Peker O, Senocak O, Tamci S, Gulbahar S, Cakmur R, Oncel S. Treatment of carpal tunnel syndrome with nerve and tendon gliding exercises. *Am.J.Phys.Med.Rehabil* 2002;81:108–113. [PubMed: 11807347]
- Amanzio M, Pollo A, Maggi G, Benedetti F. Response variability to analgesics: a role for non-specific activation of endogenous opioids. *Pain* 2001;90:205–215. [PubMed: 11207392]
- Balon JW, Mior SA. Chiropractic care in asthma and allergy. *Ann.Allergy Asthma Immunol* 2004;93:S55–S60. [PubMed: 15330012]

- Bee LA, Dickenson AH. Rostral ventromedial medulla control of spinal sensory processing in normal and pathophysiological states. *Neuroscience* 2007;147:786–793. [PubMed: 17570596]
- Benedetti F, Amanzio M, Vighetti S, Asteggiano G. The biochemical and neuroendocrine bases of the hyperalgesic nocebo effect. *J.Neurosci* 2006;26:12014–12022. [PubMed: 17108175]
- Benedetti F, Pollo A, Lopiano L, Lanotte M, Vighetti S, Rainero I. Conscious expectation and unconscious conditioning in analgesic, motor, and hormonal placebo/nocebo responses. *J.Neurosci* 2003;23:4315–4323. [PubMed: 12764120]
- Bialosky JE, Bishop MD, Robinson ME, Barabas JA, George SZ. The influence of expectation on spinal manipulation induced hypoalgesia: an experimental study in normal subjects. *BMC.Musculoskelet.Disord* 2008;9:19. [PubMed: 18267029]
- Bingel U, Lorenz J, Schoell E, Weiller C, Buchel C. Mechanisms of placebo analgesia: rACC recruitment of a subcortical antinociceptive network. *Pain* 2006;120:8–15. [PubMed: 16364549]
- Bishop PB, Wing PC. Compliance with clinical practice guidelines in family physicians managing worker's compensation board patients with acute lower back pain. *Spine J* 2003;3:442–450. [PubMed: 14609688]
- Boal RW, Gillette RG. Central neuronal plasticity, low back pain and spinal manipulative therapy. *J.Manipulative Physiol Ther* 2004;27:314–326. [PubMed: 15195039]
- Bulbulian R, Burke J, Dishman JD. Spinal reflex excitability changes after lumbar spine passive flexion mobilization. *J.Manipulative Physiol Ther* 2002;25:526–532. [PubMed: 12381975]
- Chen YX, Kong KM, Wang WD, Xie CH, Wu RH. Functional MR imaging of the spinal cord in cervical spinal cord injury patients by acupuncture at LI 4 (Hegu) and LI 11 (Quchi). *Conf.Proc.IEEE Eng Med.Biol.Soc* 2007;2007:3388–3391. [PubMed: 18002724]
- Childs JD, Fritz JM, Flynn TW, Irrgang JJ, Johnson KK, Majkowski GR, Delitto A. A clinical prediction rule to identify patients with low back pain most likely to benefit from spinal manipulation: a validation study. *Ann.Intern.Med* 2004;141:920–928. [PubMed: 15611489]
- Cleland JA, Childs JD, Fritz JM, Whitman JM, Eberhart SL. Development of a clinical prediction rule for guiding treatment of a subgroup of patients with neck pain: use of thoracic spine manipulation, exercise, and patient education. *Phys.Ther* 2007;87:9–23. [PubMed: 17142640]
- Cleland JA, Childs JD, McRae M, Palmer JA, Stowell T. Immediate effects of thoracic manipulation in patients with neck pain: a randomized clinical trial. *Man.Ther* 2005;10:127–135. [PubMed: 15922233]
- Cleland JA, Fritz JM, Whitman JM, Childs JD, Palmer JA. The use of a lumbar spine manipulation technique by physical therapists in patients who satisfy a clinical prediction rule: a case series. *J.Orthop.Sports Phys.Ther* 2006;36:209–214. [PubMed: 16676870]
- Colloca CJ, Keller TS, Gunzburg R. Neuromechanical characterization of in vivo lumbar spinal manipulation. Part II. Neurophysiological response. *J.Manipulative Physiol Ther* 2003;26:579–591. [PubMed: 14673407]
- Colloca CJ, Keller TS, Gunzburg R, Vandeputte K, Fuhr AW. Neurophysiologic response to intraoperative lumbosacral spinal manipulation. *J.Manipulative Physiol Ther* 2000;23:447–457. [PubMed: 11004648]
- Colloca CJ, Keller TS, Harrison DE, Moore RJ, Gunzburg R, Harrison DD. Spinal manipulation force and duration affect vertebral movement and neuromuscular responses. *Clin.Biomech.(Bristol., Avon.)* 2006;21:254–262.
- Coppieters MW, Alshami AM. Longitudinal excursion and strain in the median nerve during novel nerve gliding exercises for carpal tunnel syndrome. *J.Orthop.Res* 2007;25:972–980. [PubMed: 17415752]
- Coppieters MW, Butler DS. Do 'sliders' slide and 'tensioners' tension? An analysis of neurodynamic techniques and considerations regarding their application. *Man.Ther.* 2007
- Degenhardt BF, Darmani NA, Johnson JC, Towns LC, Rhodes DC, Trinh C, McClanahan B, DiMarzo V. Role of osteopathic manipulative treatment in altering pain biomarkers: a pilot study. *J.Am.Osteopath.Assoc* 2007;107:387–400. [PubMed: 17908831]
- Delaney JP, Leong KS, Watkins A, Brodie D. The short-term effects of myofascial trigger point massage therapy on cardiac autonomic tone in healthy subjects. *J.Adv.Nurs* 2002;37:364–371. [PubMed: 11872106]

- Derbyshire SW, Jones AK, Gyulai F, Clark S, Townsend D, Firestone LL. Pain processing during three levels of noxious stimulation produces differential patterns of central activity. *Pain* 1997;73:431–445. [PubMed: 9469535]
- DeVocht JW, Pickar JG, Wilder DG. Spinal manipulation alters electromyographic activity of paraspinal muscles: a descriptive study. *J.Manipulative Physiol Ther* 2005;28:465–471. [PubMed: 16182019]
- Deyle GD, Henderson NE, Matekel RL, Ryder MG, Garber MB, Allison SC. Effectiveness of manual physical therapy and exercise in osteoarthritis of the knee. A randomized, controlled trial. *Ann.Intern.Med* 2000;132:173–181. [PubMed: 10651597]
- Dishman JD, Bulbulian R. Spinal reflex attenuation associated with spinal manipulation. *Spine* 2000;25:2519–2524. [PubMed: 11013505]
- Dishman JD, Burke J. Spinal reflex excitability changes after cervical and lumbar spinal manipulation: a comparative study. *Spine J* 2003;3:204–212. [PubMed: 14589201]
- Dougherty DD, Kong J, Webb M, Bonab AA, Fischman AJ, Gollub RL. A combined [11C]diprenorphine PET study and fMRI study of acupuncture analgesia. *Behav.Brain Res.* 2008
- Ernst E. Does spinal manipulation have specific treatment effects? *Fam.Pract* 2000;17:554–556. [PubMed: 11120730]
- Fang J, Jin Z, Wang Y, Li K, Kong J, Nixon EE, Zeng Y, Ren Y, Tong H, Wang Y, Wang P, Hui KK. The salient characteristics of the central effects of acupuncture needling: Limbic-paralimbic-neocortical network modulation. *Hum.Brain Mapp.* 2008
- Field T, Diego M, Cullen C, Hernandez-Reif M, Sunshine W, Douglas S. Fibromyalgia pain and substance P decrease and sleep improves after massage therapy. *J.Clin.Rheumatol* 2002;8:72–76. [PubMed: 17041326]
- Flynn T, Fritz J, Whitman J, Wainner R, Magel J, Rendeiro D, Butler B, Garber M, Allison S. A clinical prediction rule for classifying patients with low back pain who demonstrate short-term improvement with spinal manipulation. *Spine* 2002;27:2835–2843. [PubMed: 12486357]
- Fuente-Fernandez R, Lidstone S, Stoessl AJ. Placebo effect and dopamine release. *J.Neural Transm* 2006; (Suppl):415–418.
- Gal J, Herzog W, Kawchuk G, Conway PJ, Zhang YT. Movements of vertebrae during manipulative thrusts to unembalmed human cadavers. *J.Manipulative Physiol Ther* 1997;20:30–40. [PubMed: 9004120]
- George SZ, Bishop MD, Bialosky JE, Zeppieri G Jr, Robinson ME. Immediate effects of spinal manipulation on thermal pain sensitivity: an experimental study. *BMC.Musculoskelet.Disord* 2006;7:68. [PubMed: 16911795]
- Goebel MU, Trebst AE, Steiner J, Xie YF, Exton MS, Frede S, Canbay AE, Michel MC, Heemann U, Schedlowski M. Behavioral conditioning of immunosuppression is possible in humans. *FASEB J* 2002;16:1869–1873. [PubMed: 12468450]
- Goffaux P, Redmond WJ, Rainville P, Marchand S. Descending analgesia - When the spine echoes what the brain expects. *Pain.* 2007
- Guo W, Robbins MT, Wei F, Zou S, Dubner R, Ren K. Spinal brain-derived neurotrophic factor signaling: a novel mechanism for descending pain facilitation. *J.Neurosci* 2006;26:126–137. [PubMed: 16399679]
- Herzog W, Kats M, Symons B. The effective forces transmitted by high-speed, low-amplitude thoracic manipulation. *Spine* 2001;26:2105–2110. [PubMed: 11698887]
- Herzog W, Scheele D, Conway PJ. Electromyographic responses of back and limb muscles associated with spinal manipulative therapy. *Spine* 1999;24:146–152. [PubMed: 9926385]
- Hessell BW, Herzog W, Conway PJ, McEwen MC. Experimental measurement of the force exerted during spinal manipulation using the Thompson technique. *J.Manipulative Physiol Ther* 1990;13:448–453. [PubMed: 2146356]
- Hsieh CY, Vicenzino B, Yang CH, Hu MH, Yang C. Mulligan's mobilization with movement for the thumb: a single case report using magnetic resonance imaging to evaluate the positional fault hypothesis. *Man.Ther* 2002;7:44–49. [PubMed: 11884156]
- Hsieh JC, Belfrage M, Stone-Elander S, Hansson P, Ingvar M. Central representation of chronic ongoing neuropathic pain studied by positron emission tomography. *Pain* 1995;63:225–236. [PubMed: 8628589]

- Iadarola MJ, Berman KF, Zeffiro TA, Byas-Smith MG, Gracely RH, Max MB, Bennett GJ. Neural activation during acute capsaicin-evoked pain and allodynia assessed with PET. *Brain* 1998;121(Pt 5):931–947. [PubMed: 9619195]
- Jette AM, Delitto A. Physical therapy treatment choices for musculoskeletal impairments. *Phys. Ther* 1997;77:145–154. [PubMed: 9037215]
- Johansen O, Brox J, Flaten MA. Placebo and Nocebo responses, cortisol, and circulating beta-endorphin. *Psychosom. Med* 2003;65:786–790. [PubMed: 14508021]
- Kaada B, Torsteinbo O. Increase of plasma beta-endorphins in connective tissue massage. *Gen. Pharmacol* 1989;20:487–489. [PubMed: 2526775]
- Kalauokalani D, Cherkin DC, Sherman KJ, Koepsell TD, Deyo RA. Lessons from a trial of acupuncture and massage for low back pain: patient expectations and treatment effects. *Spine* 2001;26:1418–1424. [PubMed: 11458142]
- Kaptchuk TJ. The placebo effect in alternative medicine: can the performance of a healing ritual have clinical significance? *Ann. Intern. Med* 2002;136:817–825. [PubMed: 12044130]
- Kent P, Marks D, Pearson W, Keating J. Does clinician treatment choice improve the outcomes of manual therapy for nonspecific low back pain? A metaanalysis. *J. Manipulative Physiol Ther* 2005;28:312–322.
- Khalsa PS, Eberhart A, Cotler A, Nahin R. The 2005 conference on the biology of manual therapies. *J. Manipulative Physiol Ther* 2006;29:341–346. [PubMed: 16762659]
- Kleinrensink GJ, Stoeckart R, Mulder PG, Hoek G, Broek T, Vleeming A, Snijders CJ. Upper limb tension tests as tools in the diagnosis of nerve and plexus lesions. Anatomical and biomechanical aspects. *Clin. Biomech. (Bristol., Avon.)* 2000;15:9–14.
- Kong J, Gollub RL, Rosman IS, Webb JM, Vangel MG, Kirsch I, Kaptchuk TJ. Brain activity associated with expectancy-enhanced placebo analgesia as measured by functional magnetic resonance imaging. *J. Neurosci* 2006;26:381–388. [PubMed: 16407533]
- Lanotte M, Lopiano L, Torre E, Bergamasco B, Colloca L, Benedetti F. Expectation enhances autonomic responses to stimulation of the human subthalamic limbic region. *Brain Behav. Immun* 2005;19:500–509. [PubMed: 16055306]
- Li LC, Bombardier C. Physical therapy management of low back pain: an exploratory survey of therapist approaches. *Phys. Ther* 2001;81:1018–1028. [PubMed: 11276184]
- Licciardone JC, Stoll ST, Fulda KG, Russo DP, Siu J, Winn W, Swift J Jr. Osteopathic manipulative treatment for chronic low back pain: a randomized controlled trial. *Spine* 2003;28:1355–1362. [PubMed: 12838090]
- MacDonald CW, Whitman JM, Cleland JA, Smith M, Hoeksma HL. Clinical outcomes following manual physical therapy and exercise for hip osteoarthritis: A case series. *J. Orthop. Sports Phys. Ther* 2006;36:588–599. [PubMed: 16915980]
- Malisza KL, Gregorash L, Turner A, Foniok T, Stroman PW, Allman AA, Summers R, Wright A. Functional MRI involving painful stimulation of the ankle and the effect of physiotherapy joint mobilization. *Magn Reson. Imaging* 2003a;21:489–496. [PubMed: 12878258]
- Malisza KL, Stroman PW, Turner A, Gregorash L, Foniok T, Wright A. Functional MRI of the rat lumbar spinal cord involving painful stimulation and the effect of peripheral joint mobilization. *J. Magn Reson. Imaging* 2003b;18:152–159. [PubMed: 12884326]
- Matre D, Casey KL, Knardahl S. Placebo-induced changes in spinal cord pain processing. *J. Neurosci* 2006;26:559–563. [PubMed: 16407554]
- McLean S, Naish R, Reed L, Urry S, Vicenzino B. A pilot study of the manual force levels required to produce manipulation induced hypoalgesia. *Clin. Biomech. (Bristol., Avon.)* 2002;17:304–308.
- McPartland JM, Giuffrida A, King J, Skinner E, Scotter J, Musty RE. Cannabimimetic effects of osteopathic manipulative treatment. *J. Am. Osteopath. Assoc* 2005;105:283–291. [PubMed: 16118355]
- Mohammadian P, Gonsalves A, Tsai C, Hummel T, Carpenter T. Areas of capsaicin-induced secondary hyperalgesia and allodynia are reduced by a single chiropractic adjustment: a preliminary study. *J. Manipulative Physiol Ther* 2004;27:381–387. [PubMed: 15319760]

- Moulson A, Watson T. A preliminary investigation into the relationship between cervical snags and sympathetic nervous system activity in the upper limbs of an asymptomatic population. *Man. Ther* 2006;11:214–224. [PubMed: 16820316]
- Moulton EA, Keaser ML, Gullapalli RP, Greenspan JD. Regional intensive and temporal patterns of functional MRI activation distinguishing noxious and innocuous contact heat. *J. Neurophysiol* 2005;93:2183–2193. [PubMed: 15601733]
- Ngan JM, Chow DH, Holmes AD. The kinematics and intra- and inter-therapist consistencies of lower cervical rotational manipulation. *Med. Eng Phys* 2005;27:395–401. [PubMed: 15863348]
- Oshiro Y, Quevedo AS, McHaffie JG, Kraft RA, Coghill RC. Brain mechanisms supporting spatial discrimination of pain. *J. Neurosci* 2007;27:3388–3394. [PubMed: 17392455]
- Petrovic P, Kalso E, Petersson KM, Ingvar M. Placebo and opioid analgesia-- imaging a shared neuronal network. *Science* 2002;295:1737–1740. [PubMed: 11834781]
- Peyron R, Laurent B, Garcia-Larrea L. Functional imaging of brain responses to pain. A review and meta-analysis (2000). *Neurophysiol. Clin* 2000;30:263–288. [PubMed: 11126640]
- Pickar JG, Wheeler JD. Response of muscle proprioceptors to spinal manipulative-like loads in the anesthetized cat. *J. Manipulative Physiol Ther* 2001;24:2–11. [PubMed: 11174689]
- Plaugher G, Bachman TR. Chiropractic management of a hypertensive patient. *J. Manipulative Physiol Ther* 1993;16:544–549. [PubMed: 8263434]
- Pollo A, Vighetti S, Rainero I, Benedetti F. Placebo analgesia and the heart. *Pain* 2003;102:125–133. [PubMed: 12620603]
- Price DD, Craggs J, Verne GN, Perlstein WM, Robinson ME. Placebo analgesia is accompanied by large reductions in pain-related brain activity in irritable bowel syndrome patients. *Pain* 2007;127:63–72. [PubMed: 16963184]
- Price DD, Milling LS, Kirsch I, Duff A, Montgomery GH, Nicholls SS. An analysis of factors that contribute to the magnitude of placebo analgesia in an experimental paradigm. *Pain* 1999;83:147–156. [PubMed: 10534585]
- Price DD, Staud R, Robinson ME, Mauderli AP, Cannon R, Vierck CJ. Enhanced temporal summation of second pain and its central modulation in fibromyalgia patients. *Pain* 2002;99:49–59. [PubMed: 12237183]
- Ross JK, Bereznic DE, McGill SM. Determining cavitation location during lumbar and thoracic spinal manipulation: is spinal manipulation accurate and specific? *Spine* 2004;29:1452–1457. [PubMed: 15223938]
- Rozmaryn LM, Dovel S, Rothman ER, Gorman K, Olvey KM, Bartko JJ. Nerve and tendon gliding exercises and the conservative management of carpal tunnel syndrome. *J. Hand Ther* 1998;11:171–179. [PubMed: 9730093]
- Sauro MD, Greenberg RP. Endogenous opiates and the placebo effect: a meta-analytic review. *J. Psychosom. Res* 2005;58:115–120. [PubMed: 15820838]
- Seffinger MA, Najm WI, Mishra SI, Adams A, Dickerson VM, Murphy LS, Reinsch S. Reliability of spinal palpation for diagnosis of back and neck pain: a systematic review of the literature. *Spine* 2004;29:E413–E425. [PubMed: 15454722]
- Smith LL, Keating MN, Holbert D, Spratt DJ, McCammon MR, Smith SS, Israel RG. The effects of athletic massage on delayed onset muscle soreness, creatine kinase, and neutrophil count: a preliminary report. *J. Orthop. Sports Phys. Ther* 1994;19:93–99. [PubMed: 8148868]
- Staud R, Craggs JG, Robinson ME, Perlstein WM, Price DD. Brain activity related to temporal summation of C-fiber evoked pain. *Pain* 2007;129:130–142. [PubMed: 17156923]
- Sterling M, Jull G, Wright A. Cervical mobilisation: concurrent effects on pain, sympathetic nervous system activity and motor activity. *Man. Ther* 2001;6:72–81. [PubMed: 11414776]
- Suter E, McMorland G. Decrease in elbow flexor inhibition after cervical spine manipulation in patients with chronic neck pain. *Clin. Biomech. (Bristol., Avon.)* 2002;17:541–544.
- Suter E, McMorland G, Herzog W, Bray R. Decrease in quadriceps inhibition after sacroiliac joint manipulation in patients with anterior knee pain. *J. Manipulative Physiol Ther* 1999;22:149–153. [PubMed: 10220713]

- Suter E, McMorland G, Herzog W, Bray R. Conservative lower back treatment reduces inhibition in knee-extensor muscles: a randomized controlled trial. *J.Manipulative Physiol Ther* 2000;23:76–80. [PubMed: 10714531]
- Symons BP, Herzog W, Leonard T, Nguyen H. Reflex responses associated with activator treatment. *J.Manipulative Physiol Ther* 2000;23:155–159. [PubMed: 10771499]
- Teodorczyk-Injeyan JA, Injeyan HS, Ruegg R. Spinal manipulative therapy reduces inflammatory cytokines but not substance P production in normal subjects. *J.Manipulative Physiol Ther* 2006;29:14–21. [PubMed: 16396725]
- Thomas JS, France CR, Sha D, Wiele NV. The influence of pain-related fear on peak muscle activity and force generation during maximal isometric trunk exertions. *Spine* 2008;33:342–348.
- Troyanovich SJ, Harrison DD, Harrison DE. Motion palpation: it's time to accept the evidence. *J.Manipulative Physiol Ther* 1998;21:568–571. [PubMed: 9798186]
- Tullberg T, Blomberg S, Branth B, Johnsson R. Manipulation does not alter the position of the sacroiliac joint. A roentgen stereophotogrammetric analysis. *Spine* 1998;23:1124–1128. [PubMed: 9615363]
- Vase L, Riley JL III, Price DD. A comparison of placebo effects in clinical analgesic trials versus studies of placebo analgesia. *Pain* 2002;99:443–452. [PubMed: 12406519]
- Vernon HT, Dhami MS, Howley TP, Annett R. Spinal manipulation and beta-endorphin: a controlled study of the effect of a spinal manipulation on plasma beta-endorphin levels in normal males. *J.Manipulative Physiol Ther* 1986;9:115–123. [PubMed: 2942618]
- Vicenzino B, Collins D, Benson H, Wright A. An investigation of the interrelationship between manipulative therapy-induced hypoalgesia and sympathoexcitation. *J.Manipulative Physiol Ther* 1998;21:448–453. [PubMed: 9777544]
- Vicenzino B, Collins D, Wright A. The initial effects of a cervical spine manipulative physiotherapy treatment on the pain and dysfunction of lateral epicondylalgia. *Pain* 1996;68:69–74. [PubMed: 9252000]
- Vicenzino B, Paungmali A, Buratowski S, Wright A. Specific manipulative therapy treatment for chronic lateral epicondylalgia produces uniquely characteristic hypoalgesia. *Man.Ther* 2001;6:205–212. [PubMed: 11673930]
- Vogt BA, Derbyshire S, Jones AK. Pain processing in four regions of human cingulate cortex localized with co-registered PET and MR imaging. *Eur.J.Neurosci* 1996;8:1461–1473. [PubMed: 8758953]
- Wager TD, Rilling JK, Smith EE, Sokolik A, Casey KL, Davidson RJ, Kosslyn SM, Rose RM, Cohen JD. Placebo-induced changes in FMRI in the anticipation and experience of pain. *Science* 2004;303:1162–1167. [PubMed: 14976306]
- Wang WD, Kong KM, Xiao YY, Wang XJ, Liang B, Qi WL, Wu RH. Functional MR imaging of the cervical spinal cord by use of electrical stimulation at LI4 (Hegu). *Conf.Proc.IEEE Eng Med.Biol.Soc* 2006;1:1029–1031. [PubMed: 17946437]
- Williams NH, Hendry M, Lewis R, Russell I, Westmoreland A, Wilkinson C. Psychological response in spinal manipulation (PRISM): a systematic review of psychological outcomes in randomised controlled trials. *Complement Ther.Med* 2007;15:271–283. [PubMed: 18054729]
- Wright A. Hypoalgesia post-manipulative therapy: a review of a potential neurophysiological mechanism. *Man.Ther* 1995;1:11–16. [PubMed: 11327789]
- Zhang J, Dean D, Nosco D, Strathopoulos D, Floros M. Effect of chiropractic care on heart rate variability and pain in a multisite clinical study. *J.Manipulative Physiol Ther* 2006;29:267–274. [PubMed: 16690380]
- Zubieta JK, Bueller JA, Jackson LR, Scott DJ, Xu Y, Koeppe RA, Nichols TE, Stohler CS. Placebo effects mediated by endogenous opioid activity on mu-opioid receptors. *J.Neurosci* 2005;25:7754–7762. [PubMed: 16120776]




Figure 1. Comprehensive model of the mechanisms of manual therapy

Figure Key: The model suggests a transient, mechanical stimulus to the tissue produces a chain of neurophysiological effects. Solid arrows denote a direct mediating effect. Broken arrows denote an associative relationship which may include: -----> = an association between a construct and its measure Bold boxes indicate the measurement of a construct ACC = anterior cingular cortex; PAG = periaqueductal gray; RVM = rostral ventromedial medulla




Figure 2. Pathway for a spinal cord mediated effect of manual therapy from George et al (2006)
 Figure Key: Proposed model pathway of study by George et al (2006) suggesting a spinal cord mediating effect of manual therapy. Bold arrows indicate suggested mechanism. Note mediating effect is suggested to be through the spinal cord due to measurement of the associated relationship of temporal summation. Also note, the design of this study neglects to consider potential supraspinal mediated effects.




Figure 3. Pathway considering both a spinal cord and supraspinal mediated effect from Bialosky et al (2008)

Figure Key: Proposed model pathway of study by Bialosky et al (2008) which considers both a spinal cord and supraspinal mediating effect of MT. Bold arrows indicate suggested mechanism. Note mediating effect is suggested to be through both the spinal cord due to measurement of the associated relationship of temporal summation and through a supraspinal mechanism due to measurement of the associated relationship of expectation.




Figure 4. Comprehensive model for the mechanisms of manual therapy illustrating similar neurophysiological activity in response to non-specific effects such as placebo and expectation
 A limitation of the current mechanistic literature in MT is the failure to adequately account for non-specific effects such as placebo and expectation. Italicized references are examples of studies from the placebo and expectation literature which have reported similar neurophysiological effects as have been associated with MT. These similarities emphasize the potential for non-specific effects to play a significant role in the mechanisms behind MT and the need to specifically address these factors in future studies.

Table 1

Categorization of manual therapy techniques

Manual Therapy Technique	Definition	Desired Outcomes
Joint Biased <ul style="list-style-type: none"> • Manipulation • Mobilization 	<ul style="list-style-type: none"> • Passive movement of a joint beyond the normal range of motion. • Passive movement of a joint within its normal range of motion. 	<ul style="list-style-type: none"> • Improved range of motion • Decrease muscle spasm • Decreased pain
Soft Tissue Biased <ul style="list-style-type: none"> • Swedish Massage • Deep Tissue Massage • Trigger Point Massage • Shiatsu Massage 	<ul style="list-style-type: none"> • Stroking and kneading of the skin and underlying soft tissue • Deep stroking and pressure across the muscles and soft tissue • Deep pressure to areas of local tenderness • Varying, rhythmic pressure from the fingers 	<ul style="list-style-type: none"> • Improve circulation • Decrease muscle spasm • Relaxation • Re-align soft tissue • Break adhesions • Increase range of motion • Release muscle spasm • Remove cellular exudates • Improve Circulation • Decrease muscle spasm • Relaxation
Nerve Biased <ul style="list-style-type: none"> • Neural Dynamics 	<ul style="list-style-type: none"> • Passive, combined movement of the spine and extremities, within their normal range of motion, in ways to elongate or tension specific nerves. 	<ul style="list-style-type: none"> • Improve range of motion • Decrease pain

Classification of manual therapy techniques referenced in manuscript along with specific examples of each. Proposed model is general and accounts for all techniques regardless of their theorized anatomical emphasis. Adapted from NCCAM website.(<http://nccam.nih.gov/>, 2007)