Neuroimunoendocrinologia

Mediadores Lipídicos

Pedro Augusto Carlos Magno Fernandes
Departamento de Fisiologia, sala 317.
e-mail: pacmf@usp.br

Mediadores Lipídicos

Ativação da PLA2

Ativação da PLA2

Respiratory Research

Transformação do ácido araquidônico

Diacylglycerol or phospholipid

Ciclooxigenases

COX1: constitutiva na maioria dos tecidos (microglia)

COX2: induzida (constitutiva em neurônios; coração)

Neuroimmunomodulação

Leukotriene pathway

Mediadores Lipídicos - Ações

Mediadores Lipídicos - Ações

Nature Reviews | Immunology

T-cell trafficking in asthma: lipid mediators grease the way

Andrew D. Luster & Andrew M. Tager Nature Reviews Immunology 4, 711-724 (September 2004)

Mediadores Lipídicos - Ações

Neuroimunoendocrinologia

Resposta imune inata

Cyclooxygenase

Ciclooxigenase

Fosfolipídios da membrana celular

Glicocorticóides

+

Anexinas
(Lipocortinas)

-

Fosfolipase A2

COX1

Ácido araquidônico

COX2

-

AINEs

PGG_{2α}

L
P
O
X

Prostaglandinas

- Vasodilatação
- Sensibilização à dor
- Febre
- Contração uterina
- Amadurecimento do colo uterino
- Diminuição da secreção de HCl
- Homeostasia renal
- Drenagem do humor aquoso

Prostaciclina

- Vasodilatação
- Sensibilização à dor
- Citoproteção gástrica
- Homeostasia renal
- Inibição da agregação plaquetária
- Homeostasia renal

Tromboxanos

- Broncoconstrição
- Agregação plaquetária

5-HPTE

Leucotrienos

- Quimiotaxia
- Ativação dos fagócitos
- Broncoconstrição

AINEs

Anti-inflamatórios não esteroidais

ácido acetil-salicílico (Aspirina)

Schematic depiction of the structural differences between the substrate-binding channels of COX-1 and COX-2 that allowed the design of selective inhibitors

Receptores de PGE2

Receptores de PGE2

Brief Reviews

THE JOURNAL OF
IMMUNOLOGY

Regulation of Immune Responses by Prostaglandin E₂

Pawel Kalinski

Mediadores Lipídicos - imunomodulação

Azul: relevante para patógenos intracelulares e cancer

Verde: relevante para patógenos extracelulares

Mediadores Lipídicos - imunomodulação

Azul: relevante para patógenos intracelulares e cancer

Verde: relevante para patógenos extracelulares

Mediadores Lipídicos - imunomodulação

Azul: relevante para patógenos intracelulares e cancer

Verde: relevante para patógenos extracelulares

ELSEVIER

Contents lists available at ScienceDirect

Journal of Neuroimmunology

journal homepage: www.elsevier.com/locate/jneuroim

Inhibition of prostaglandin E₂ EP3 receptors improves stroke injury via anti-inflammatory and anti-apoptotic mechanisms

Yuri Ikeda-Matsuo^{a,*}, Hayato Tanji^a, Shuh Narumiya^b, Yasuharu Sasaki^a

^a Laboratory of Pharmacology, School of Pharmaceutical Sciences, Kitasato University, Tokyo, Japan

^b Department of Pharmacology, Kyoto University Faculty of Medicine, Kyoto, Japan

E-Prostanoid (EP)2/EP4 Receptor-Dependent Maturation of Human Monocyte-Derived Dendritic Cells and Induction of Helper T2 Polarization

Shinichiro Kubo, Hideo Kohka Takahashi, Masao Takei, Hiromi Iwagaki, Tadashi Yoshino, Noriaki Tanaka, Shuji Mori, and Masahiro Nishibori

Departments of Pharmacology (S.K., H.K.T., S.M., M.N.), Tumour Biology (S.K., H.K.T., H.I., N.T.), and Pathology (T.Y.), Okayama University Graduate School of Medicine and Dentistry, Okayama, Japan; and Forshungazentrum Borstel (M.T.), Borstel, Germany

Received November 11, 2003; accepted February 9, 2004

ELSEVIER

Contents lists available at SciVerse ScienceDirect

Molecular Immunology

journal homepage: www.elsevier.com/locate/molimm

PGE₂ differentially regulates monocyte-derived dendritic cell cytokine responses depending on receptor usage (EP₂/EP₄)[☆]

Neil J. Poloso^{a,*}, Paula Urquhart^b, Anna Nicolaou^b, Jenny Wang^a, David F. Woodward^a

^a 2525 DuPont Dr, RD3-2B, Department Biological Sciences, Allergan Inc., Irvine, CA 92612, USA

^b Bradford School of Pharmacy, School of Life Sciences, University of Bradford, Bradford BD7 1PD, UK

RESEARCH

Open Access

Differential expression of E-type prostanoid receptors 2 and 4 in microglia stimulated with lipopolysaccharide

Ester Bonfill-Teixidor^{1,2}, Amaia Otxoa-de-Amezaga^{1,2}, Miriam Font-Nieves¹, M. Glòria Sans-Fons² and Anna M. Planas^{1,2,3*}

Balanço Th1/Th2

Normal Balance

Autoimmune Disease

Cancer

Immunomodulation

Balanço Th1/Th2

Normal Balance

Autoimmune Disease

Cancer

Immunomodulation

Mediadores Lipídicos - imunomodulação

Mediadores Lipídicos - Neuroimunomodulação

Mediadores Lipídicos - Ações

Figure 1 | The proposed P2X4 pathway to pain. ATP, released following adjuvant stimulation of Toll-like receptors, opens the P2X4 receptor, in turn causing calcium influx. A cascade of events ensues, including calcium-dependent translocations of cPLA₂ and 5-LO to the nucleus and endoplasmic reticulum. cPLA₂ liberates AA, the substrate for COX1, COX2 and 5-LO. These enzymes catalyze the formation of PGH₂ and LTA₄, respectively, which are further metabolized to terminal prostaglandins and leukotrienes. PGE₂, PGI₂ and LTB₄ evoke pain responses via their respective receptors. Abbreviations: AA, arachidonic acid; COX, cyclooxygenase; cPLA₂, cytosolic phospholipase A₂; 5-LO, 5-lipoxygenase; LTA₄, leukotriene A₄; LTB₄, leukotriene B₄; PGE₂, prostaglandin E₂; PGI₂, prostacyclin; PGH₂, prostaglandin H₂.

Mediadores Lipídicos - Ações

Papel das PGs- DOR

PGE₂ inibe MnPO levando a desinibição de vias descendentes inibitórias da MPO (medial preoptic area) para o sistema antinociceptivo.

PAG : periaqueductal gray matter

RMR: rostral medullary raphe

Papel das PGs na neuroimunomodulação

Reflexo vagal

a

Via aferente

Reflexo vagal

Fig. 1. A: mean core body temperature for sham surgery (Sham) and vagotomized (Vag) subjects injected at *time 0* with saline (Sal) or 0.1 $\mu\text{g}/\text{kg}$ recombinant human interleukin-1 β (rh-IL-1 β). B: the difference between core body temperature after injection with rh-IL-1 β and after Sal for Sham and Vag subjects.

Efeito relativo do vago no controle do aumento da temperatura corporal induzida por administração de IL- β

Via aferente

Reflexo vagal

Efeito relativo do vago no controle do aumento da temperatura corporal induzida por administração de IL- β

Fig. 2. A: mean core body temperature for Sham and Vag subjects injected at *time 0* with Sal or 0.5 $\mu\text{g/kg}$ rh-IL-1 β . B: the difference between core body temperature after injection with rh-IL-1 β and after Sal for Sham and Vag subjects.

Via aferente

Reflexo vagal

Efeito relativo do vago no controle do aumento da temperatura corporal induzida por administração de IL- β

Fig. 3. A: mean core body temperature for Sham and Vag subjects injected at *time 0* with Sal or 1.0 μ g/kg rh-IL-1 β . B: the difference between core body temperature after injection with rh-IL-1 β and after Sal for Sham and Vag subjects.

Via aferente

Intravenous Lipopolysaccharide Induces Cyclooxygenase 2-Like Immunoreactivity in Rat Brain Perivascular Microglia and Meningeal Macrophages

JOEL K. ELMQUIST,^{1*} CHRISTOPHER D. BREDER,² JONATHAN E. SHERIN,¹
THOMAS E. SCAMMELL,¹ WILLIAM F. HICKEY,³ DAVID DEWITT,⁴
AND CLIFFORD B. SAPER¹

¹Department of Neurology and Program in Neuroscience, Harvard Medical School,
Beth Israel Hospital, Boston, Massachusetts 02115

²Department of Anesthesiology and Critical Care Medicine, Johns Hopkins University,
Baltimore, Maryland 21287

³Department of Pathology, Dartmouth Medical School, Lebanon, New Hampshire 03756

⁴Department of Biochemistry, Michigan State University, East Lansing, Michigan 48824

NTS

córtex

Papel das PGE2 na neuroimunomodulação

Dual modulation of synaptic transmission in the nucleus tractus solitarius by prostaglandin E₂ synthesized downstream of IL-1 β

Vincent Marty, Mickaël El Hachmane and Thierry Amédée

Psychoneuroimmunology, Nutrition and Genetics (PsyNuGen), UMR 5226 CNRS-UB2/UMR 1286 INRA-UB2, Université Bordeaux 2, Institut François Magendie, 146 Rue Léo-Saignat, F-33077 Bordeaux, France

Keywords: brain stem slices, inflammation, nucleus tractus solitarius, PGE₂, synaptic transmission

Abstract

The activation of the innate immune system induces the production of blood-borne proinflammatory cytokines like interleukin-1 β (IL-1 β), which in turn triggers brain-mediated adaptive responses referred to as sickness behaviour. These responses involve the modulation of neural networks in key regions of the brain. The nucleus tractus solitarius (NTS) of the brainstem is a key nucleus for immune-to-brain signalling. It is the main site of termination of vagal afferents and is adjacent to the area postrema, a circumventricular organ allowing blood-borne action of circulating IL-1 β . Although it is well described that IL-1 β activates cerebral endothelial and glial cells, it is still unknown if and how IL-1 β or downstream-synthesized molecules impact NTS synaptic function. In this study we report that IL-1 β did not modulate NTS synaptic transmission *per se*, whereas prostaglandin E₂ (PGE₂), which is produced downstream of IL-1 β , produced opposite effects on spontaneous and evoked release. On the one hand, PGE₂ facilitated glutamatergic transmission between local NTS neurons by enhancing the frequency of spontaneous excitatory postsynaptic currents through a presynaptic receptor different from the classical EP1–4 subtypes. On the other hand, PGE₂ also depressed evoked excitatory input from vagal afferent terminals through presynaptic EP3 receptors coupled to G-proteins linked to adenylyl cyclase and protein kinase A activity. Our data show that IL-1 β -induced PGE₂ can modulate evoked and spontaneous release in the NTS differentially through different mechanisms. These data unravel some molecular mechanisms by which innate immune stimuli could signal to, and be integrated within, the brainstem to produce central adaptive responses.

Papel das PGE2 na neuroimunomodulação

Papel das PGE2 na neuroimunomodulação

Key points

- Prostaglandin E₂ (PGE₂) regulates multiple responses to illness and psychological stress via distinct PGE receptor subtypes (EP1, EP2, EP3 and EP4)
- During illness-induced fever, PGE₂ activates EP3 in neurons in the preoptic hypothalamus, which promotes two independent effector mechanisms: heat production in brown adipose tissue and vasoconstriction of the skin
- In illness-induced activation of the hypothalamic–pituitary–adrenal axis, EP1 and EP3 mediate distinct neuronal pathways that converge on neurons in the paraventricular hypothalamus that contain corticotropin-releasing factor
- Under psychological stress, the PGE₂–EP1 signaling pathway regulates impulsive behaviors via the midbrain dopaminergic system
- PGE₂ regulates illness-induced impairment of memory formation, but under physiological conditions (without illness), it supports hippocampus-dependent memory formation and synaptic plasticity via EP2
- PGE₂ is synthesized by enzymes induced inside and outside the brain in response to illness and by constitutive enzymes in neurons and/or microglia in response to psychological stress

Papel das PGs- FEBRE

Agonista EP4: efeito oposto

PGE₂ é produzida por células endoteliais e perivasculares ao longo de vênulas nas extremidades do cérebro, em especial, na área pré-óptica do hipotálamo (controle da termogênese).

MnPO: median preoptic nucleus

DMH: dorsomedial nucleus of the hypothalamus

RMR: rostral medullary raphe

Neural circuitry engaged by prostaglandins during the sickness Syndrome. Clifford B Saper, Andrej A Romanovsky & Thomas E Scammell Nature Neuroscience 15, 1088–1095 (2012).

Papel das PGs- DOR

Estresse

RVLM, rostral ventrolateral medulla
 POA, preoptic area
 BNST, bed nucleus of the stria terminalis
 CeA, central nucleus of the amygdala
 PVH, paraventricular

NSAIDs: Bloqueiam a ativação do eixo HPA

Efeitos dos glicocorticóides - Sistema Nervoso

IL-1 β induz produção de prostanglandinas no sistema nervosos central

(reflexo imunológico)

PGE₂ induz a febre e alterações comportamentais ativando o núcleo preóptico medial

Glicocorticóides inibem PLA₂ e COX

Table 1. Reported changes of COX-1 and COX-2 in neurodegenerative diseases

Disorder	COX-1	COX-2
AD	Small trial with indomethacin (preferential COX-1 inhibitor) beneficial in protecting from cognitive decline. Use of aspirin (preferential COX-1 inhibitor) associated with reduced risk of AD. Increased COX-1-expressing microglia surrounding amyloid plaques.	Upregulation of COX-2 in early AD, and downregulation in advanced AD stages. Randomized trials with COX-2-selective inhibitors not effective.
PD	COX-1 expression unchanged by MPTP, but >65% of ventral midbrain PGE ₂ originated from COX-1. Aspirin (preferential COX-1 inhibitor) and mixed COX inhibitors neuroprotective in parkinsonian models.	COX-2 expressed within dopaminergic neurons of substantia nigra in the PD postmortem brain. COX-2 induced in the MPTP model of PD during destruction of nigrostriatal pathway. COX-2-selective inhibitors neuroprotective in parkinsonian models.
TBI	Accumulation of COX-1 expressing microglia at peri-lesional areas and in the developing core in patients with TBI.	Conflicting reports on COX-2-selective inhibitors in experimental TBI: nimesulide reversed motor and cognitive abnormalities; celecoxib worsened motor but not cognitive performance; and rofecoxib did not limit neuronal cell death. No difference in lesion volume 3 weeks after TBI and no changes in Morris water maze performance in COX-2-null versus wild-type mice. Not determined.
Prion diseases	Accumulation of COX-1-expressing microglia in areas of severe tissue damage and around vacuoles in CJD patients.	
HIV dementia	Twofold increase in COX-1 mRNA expression in the demented versus non-demented patients.	COX-2 mRNA expression is unchanged.
Stroke	COX-1 deletion increases vulnerability to focal cerebral ischemia. COX-1 inhibition reduced neuronal injury and oxidative stress in hippocampus during transient global ischemia.	COX-2 expression upregulated in neurons in early ischemia. COX-2 inhibition or genetic deletion reduces infarct volume and neurological deficits.
ALS	COX-1, and not COX-2, is the primary source of PGE ₂ in spinal cords of SOD1 ^{G93A} mice. However, deletion of COX-1 does not improve preservation of motor neurons and survival of transgenic mutant mice.	Levels of COX-2 mRNA and PGE ₂ increased in mutant SOD1 mice. Selective COX-2 inhibitors (celecoxib and rofecoxib) improve motor performance, extend survival and reduce CSF levels of PGE ₂ . COX-2 expression increased in motor neurons, interneurons and glial cells in seven human sporadic ALS patients. Double-blinded, placebo-controlled clinical trial with celecoxib (800 mg/day) administered for 12 months in ALS subjects did not show any beneficial effect.
MS	Not determined.	No correlation between PGE ₂ and clinical scores of MS severity or biomarkers of axonal or astroglial injury.

Inflamação e Doença de Alzheimer