

Tabela de Transformadas de Laplace (autores: W. E. Boyce, R. C. DiPrima& D. B. Meade)

$f(t) = \mathcal{L}^{-1}\{F(s)\}$	$F(s) = \mathcal{L}\{f(t)\}$	Notes
1. 1	$\frac{1}{s}, \quad s > 0$	Sec. 6.1; Ex. 4
2. e^{at}	$\frac{1}{s-a}, \quad s > a$	Sec. 6.1; Ex. 5
3. t^n , n a positive integer	$\frac{n!}{s^{n+1}}, \quad s > 0$	Sec. 6.1; Prob. 24
4. t^p , $p > -1$	$\frac{\Gamma(p+1)}{s^{p+1}}, \quad s > 0$	Sec. 6.1; Prob. 24
5. $\sin(at)$	$\frac{a}{s^2 + a^2}, \quad s > 0$	Sec. 6.1; Ex. 7
6. $\cos(at)$	$\frac{s}{s^2 + a^2}, \quad s > 0$	Sec. 6.1; Prob. 5
7. $\sinh(at)$	$\frac{a}{s^2 - a^2}, \quad s > a $	Sec. 6.1; Prob. 7
8. $\cosh(at)$	$\frac{s}{s^2 - a^2}, \quad s > a $	Sec. 6.1; Prob. 6
9. $e^{at} \sin(bt)$	$\frac{b}{(s-a)^2 + b^2}, \quad s > a$	Sec. 6.1; Prob. 10
10. $e^{at} \cos(bt)$	$\frac{s-a}{(s-a)^2 + b^2}, \quad s > a$	Sec. 6.1; Prob. 11
11. $t^n e^{at}$, n a positive integer	$\frac{n!}{(s-a)^{n+1}}, \quad s > a$	Sec. 6.1; Prob. 14
12. $u_c(t) = \begin{cases} 0 & t < c \\ 1 & t \geq c \end{cases}$	$\frac{e^{-cs}}{s}, \quad s > 0$	Sec. 6.3
13. $u_c(t) f(t-c)$	$e^{-cs} F(s)$	Sec. 6.3
14. $e^{ct} f(t)$	$F(s-c)$	Sec. 6.3
15. $f(ct)$	$\frac{1}{c} F\left(\frac{s}{c}\right), \quad c > 0$	Sec. 6.3; Prob. 17
16. $(f * g)(t) = \int_0^t f(t-\tau)g(\tau) d\tau$	$F(s)G(s)$	Sec. 6.6
17. $\delta(t-c)$	e^{-cs}	Sec. 6.5
18. $f^{(n)}(t)$	$s^n F(s) - s^{n-1} f(0) - \dots - f^{(n-1)}(0)$	Sec. 6.2; Cor. 6.2.2
19. $(-t)^n f(t)$	$F^{(n)}(s)$	Sec. 6.2; Prob. 21