






A biodiversidade no tempo profundo **(ou como a Paleontologia nos ajuda a** **entender o mundo que nos cerca)**

Catalogue of Life



Checklist 2021 > 2 milhões spp






“we estimate that there exist globally about 0.8–1.6 million prokaryotic OTUs”

Diversidade atual (2 Ma) vs pretérita (500 Ma)

250 x 2 milhões = 500 milhões



c. 500 mil / 250 = 2.000 spp

Hexapoda:
900 mil vs 7 mil



Brachiopoda:
300 vs 12 mil



Por que o eixo temporal é importante no estudo da biodiversidade?

- 1 – Entender que a **discrepância** atual não é limite morfológico absoluto
- 2 – Comparar a diversidade biológica atual com aquela de épocas passadas (e estas entre si), ajudando a entender aspectos do processo **evolutivo**
- 3 – Investigar os eventos de **extinção** como parâmetros para estudos das relações da biota com o meio físico
- 4 – Desmistificar conceitos fixistas que permeiam o discurso **ambientalista**




entender o mundo que nos cerca

1 – Entender que a **discrepância** atual não é limite morfológico absoluto



Membracidae





Curculionidae



Paleontologia:
“morfo-espécies”



Tamanho como parâmetro para diversidade












Amphicoelias fragillimus

Afinidad: Ninguna actual,
Saurópodos
58m de largo


















- 1 – Entender que a **discrepância** atual não é limite morfológico absoluto
- 2 – Comparar a diversidade biológica atual com aquela de épocas passadas (e estas entre si), ajudando a entender aspectos do processo **evolutivo**

“Explosão” do
Cambriano











2 – Comparar a diversidade biológica atual com aquela de épocas passadas (e estas entre si), **ajudando a entender aspectos do processo evolutivo**





G. G. Simpson







Heinrich Bronn (1800-1862): diagramas de fuso (spindle)



“Early burst”



Tempo and Mode in Evolution

George Gaylord Simpson

A Columbia Classic in Evolution

With a new introduction by
George Gaylord Simpson

Irradiação Adaptativa

- 1 – Novas estruturas
- 2 – Mudança ambiental
- 3 – Extinções





-

Tempo and Mode in Evolution



George Gaylord Simpson

A Columbia Classic in Evolution



With a new introduction by
George Gaylord Simpson

Irradiação Adaptativa

- 1 – Novas estruturas
- 2 – Mudança ambiental
- 3 – Extinções




Teria a biodiversidade variado no passado?





Groenlândia

Permiano sup.





Triássico inf.




China






1860: registro fóssil da Ilhas
Britânicas com correção para
maior amostragem do Paleozóico




Evolução da fauna
marinha fanerozóica


John Sepkoski




Evolução da fauna marinha fanerozóica




Evolução da fauna marinha fanerozóica





Evolução da fauna marinha fanerozóica



Enquanto isso nos continentes ...




Enquanto isso nos continentes ...







Curvas de diversidade por categorias taxonômicas




David Raup



Diversidade biológica pode ter se mantido praticamente estável desde meados do Paleozóico




O problema da amostragem





Mais rochas
=
Mais fósseis
=
Mais táxons


O problema da amostragem




John Alroy: correção para viés de amostragem




Extrapolação de padrões locais



Diversidade total é fator de
presença de mais rochas
Mas unidades mais novas
não tem maior diversidade




3 – Investigar os eventos de **extinção** como parâmetros para estudos das relações da biota com o meio físico



Extinções (estudos de caso)

“Positivos”

2 bilhões de anos




360 milhões de anos



“Negativos”



240 milhões de anos





65 milhões de anos












420 Ma






360 Ma




Sequestro de carbono


Carbon Sequestration
















REVIEW

doi:10.1038/nature09678

Has the Earth's sixth mass extinction already arrived?


Anthony D. Barnosky^{1,2,3}, Nicholas Matzke¹, Susumu Tomiya^{1,2,3}, Guinevere O. U. Wogan^{1,3}, Brian Swartz^{1,2}, Tiago B. Quental^{1,2†}, Charles Marshall^{1,2}, Jenny L. McGuire^{1,2,3,4†}, Emily L. Lindsey^{1,2}, Kaitlin C. Maguire^{1,2}, Ben Mersey^{1,4} & Elizabeth A. Ferrer^{1,2}

Palaeontologists characterize mass extinctions as times when the Earth loses more than three-quarters of its species in a geologically short interval, as has happened only five times in the past 540 million years or so. Biologists now suggest that a sixth mass extinction may be under way, given the known species losses over the past few centuries and millennia. Here we review how differences between fossil and modern data and the addition of recently available palaeontological information influence our understanding of the current extinction crisis. Our results confirm that current extinction rates are higher than would be expected from the fossil record, highlighting the need for effective conservation measures.

ScienceAdvances Home News Journals Topics Careers

Webinar

Cell and gene therapies for cancer:
Future promises and challenges



January 23, 2019

12 noon ET, 9 a.m. PT,
5 p.m. GMT, 6 p.m. CET

[Register](#)

Science
AAAS

Sponsored by
Bio-Techne

Institution: US
Log in | My

SHARE



0



0



0

RESEARCH ARTICLE | ENVIRONMENTAL SCIENCES

Accelerated modern human-induced species losses: Entering the sixth mass extinction

Gerardo Ceballos^{1,4}, Paul R. Ehrlich², Anthony D. Barnosky³, Andrés García⁴, Robert M. Pringle⁵ and Todd M. Palmer⁶

† See all authors and affiliations

Science Advances 19 Jun 2015:
Vol. 1, no. 5, e1400253
DOI: 10.1126/sciadv.1400253

[Article](#)

[Figures & Data](#)

[Info & Metrics](#)

[eLetters](#)

[PDF](#)

Abstract

The oft-repeated claim that Earth's biota is entering a sixth "mass extinction" depends on


4 – Desmistificar conceitos fixistas que permeiam o discurso **ambientalista**



Perguntas:

- 1: Sabemos se o número de espécies está mesmo diminuindo?
- 2: É possível comparar essa possível diminuição com o registro fóssil?

Perda e “surgimento” de espécies






Crédito: Instituto Tamandua


Preguiça-de-coleira
do-Nordeste




Crédito: Suelen Sanches

Preguiça-de-coleira
do-Sudeste








Rinoceronte branco
“do norte”



Sapo das
cachoeiras de Kihansi



Global meta-analysis reveals no net change in local-scale plant biodiversity over time

Mark Vellend^{a,1}, Lander Baeten^{b,c}, Isla H. Myers-Smith^d, Sarah C. Elmendorf^e, Robin Beauséjour^a, Carissa D. Brown^a, Pieter De Frenne^b, Kris Verheyen^b, and Sonja Wipf^f

^aDépartement de Biologie, Université de Sherbrooke, Sherbrooke, QC, Canada J1K 2R1; ^bDepartment of Forest and Water Management, Forest and Nature Lab, Ghent University, BE-9090 Melle-Gontrode, Belgium; ^cDepartment of Biology, Terrestrial Ecology Unit, Ghent University, BE-9000 Gent, Belgium; ^dSchool of GeoSciences, University of Edinburgh, Edinburgh EH9 3JW, United Kingdom; ^eNational Ecology Observatory Network, Boulder, CO 80301; and ^fWSL Institute for Snow and Avalanche Research SLF, CH-7260 Davos, Switzerland

Edited by Peter M. Kareiva, The Nature Conservancy, Seattle, WA, and approved October 1, 2013 (received for review July 9, 2013)

Global biodiversity is in decline. This is of concern for aesthetic and ethical reasons, but possibly also for practical reasons, as suggested by experimental studies, mostly with plants, showing that biodiversity reductions in small study plots can lead to compromised ecosystem function. However, inferring that ecosystem functions will decline due to biodiversity loss in the real world rests on the untested assumption that such loss is actually occurring at these small scales in nature. Using a global database of 168 published studies and >16,000 nonexperimental, local-scale vegetation plots, we show that mean temporal change in species diversity over periods of 5–261 y is not different from zero, with increases at least as likely as declines over time. Sites influenced primarily by plant species' invasions showed a tendency for declines in species richness, whereas sites undergoing postdisturbance succession showed increases in richness over time. Other distinctions among studies had little influence on temporal richness trends. Although maximizing diversity is likely important for maintaining ecosystem function in intensely managed systems such as restored grasslands or tree plantations, the clear lack of any general tendency for plant biodiversity to decline at small scales in nature directly contradicts the key assumption linking experimental results to ecosystem function as a motivation for biodiversity conservation in nature. How often real world changes in the diversity and composition of plant communities at the local scale cause ecosystem function to deteriorate, or actually to improve, remains unknown and is in critical need of further study.

spatial scale | permanent plots | ecosystem services

A huge number of experiments has investigated the effects of species diversity (typically the number of species) on ecosystem function in small study plots (<400 m²), with a general consensus emerging that processes such as primary productivity and nutrient uptake increase as a function of the number of species in a community (1–6). These experiments thus appear to provide a powerful motivation for biodiversity conservation, given that ecosystem functions underpin many ecosystem services from which people benefit, such as forage production and carbon sequestration (1). However, the link between diversity-function experiments and the widespread argument that ecosystem function should motivate biodiversity conservation (7–11) hinges on the untested assumption that global biodiversity declines apply to the small scale (2). Experimental studies typically focus on small spatial scales not only for practical reasons, but also because organisms, plants in particular, typically interact over short distances (12), and so it is at the small scale that biodiversity is most likely to have an important impact on the functioning of ecosystems (13–15).

Habitat loss, invasive species, and overexploitation, among other factors, have accelerated global species' extinction well beyond the background rate (16–18), and it is tempting to assume that a global decline in biodiversity is necessarily accompanied by declines at smaller spatial scales. However, this is not

a logical inevitability because, unlike other key variables involved in global environmental change, biodiversity at large scales (often termed gamma diversity) is not an additive function of biodiversity at smaller scales (alpha diversity). If global temperature or atmospheric CO₂ concentrations, for example, are increasing at the global scale, the net change over time within local areas must, on average, be positive. However, because local species losses may be accompanied by immigration of species from elsewhere, decreases in biodiversity at the global scale do not necessarily result in any biodiversity change at smaller scales (16, 19, 20). Here we present a global synthesis testing for directional changes in local-scale biodiversity of terrestrial plants, which have been the focus of most well-replicated biodiversity-ecosystem function (BDEF) experiments. We focus on the most commonly studied component of biodiversity—species diversity—estimated by metrics that reflect the number of species (richness) and/or the equitability of their abundances (indices of diversity or evenness).

Results and Discussion

We conducted a systematic global meta-analysis of plant species diversity change over time in >16,000 plots (median plot area = 44 m²) from all major vegetation types (Fig. 1), including areas under profound and direct human influence (e.g., urban areas or pastures) and more pristine areas. Effect sizes were calculated as the log ratio of species diversity [e.g., species richness (*SR*)] in the year of the final survey (*Y2*) and the initial survey (*Y1*) divided by

Significance

A major advance of the last 20 y at the interface of biological, environmental, and conservation sciences has been the demonstration that plant biodiversity positively influences ecosystem function. Linking these results to applied conservation efforts hinges on the assumption that biodiversity is actually declining at the local scale at which diversity-function relationships are strongest. Our compilation and analysis of a global database of >16,000 repeat survey vegetation plots from habitats across the globe directly contradict this assumption. We find no general tendency for local-scale plant species diversity to decline over the last century, calling into question the widespread use of ecosystem function experiments to argue for the importance of biodiversity conservation in nature.

Author contributions: M.V. and I.H.M.-S. designed research; M.V., L.B., I.H.M.-S., S.C.E., R.B., C.D.B., P.D.F., K.V., and S.W. performed research; L.B. and S.C.E. analyzed data; and M.V., L.B., I.H.M.-S., and S.C.E. wrote the paper.

The authors declare no conflict of interest.



This article is a PNAS Direct Submission.


Data deposition: Our database is provided as Dataset S1.

See Commentary on page 19187.

To whom correspondence should be addressed. E-mail: mark.vellend@usherbrooke.ca.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1312779110/-DCSupplemental.





No net insect abundance and diversity declines across US Long Term Ecological Research sites





Figure 3: The global Living Planet Index (1970 to 2018)

The average change in relative abundance of 31,821 populations, representing 5,230 species monitored across the globe, was a decline of 69%. The white line shows the index values and the shaded areas represent the statistical certainty surrounding the trend (95% statistical certainty, range 63% to 75%).
Source: WWF/ZSL (2022)¹⁸⁴.



THIS REPORT
HAS BEEN
PRODUCED IN
COLLABORATION
WITH

ZSL
FOR LIFE
EVERYWHERE



LIVING PLANET REPORT 2022

BUILDING A NATURE-POSITIVE SOCIETY




Figure 6: The Freshwater Living Planet Index: 1970 to 2016

The average abundance of 3,741 freshwater populations, representing 944 species, monitored across the globe declined by 84% on average. The white line shows the index values and the shaded areas represent the statistical certainty surrounding the trend (range: -89% to -77%)¹⁰⁷.

Key

-  Freshwater Living Planet Index
-  Confidence limits





Fig. 9: TERRESTRIAL LIVING PLANET INDEX, 1970–2005




Fig. 10: MARINE LIVING PLANET INDEX, 1970–2005





Fig. 11: FRESHWATER LIVING PLANET INDEX, 1970–2005





Living Planet Index

Home About Data Portal Projects Publications My Account

Keyword Search

Keyword

Refine Search

Advanced Search

Search Result: 3 record(s)

Map View List View

Species Name	Common Name	Location of Population	Countries	View
Mus musculus	House mouse	Residential area in Anxiang County, Hunan	China	View
Mus musculus	House mouse	Zhumadian Township of Nanxun County, Hunan	China	View
Mus musculus	House mouse	Pacific Valley, Sumnerland, British Columbia	Canada	View
Mus musculus	House mouse	Lachay National Reserve, Northern Lima, Peru	Peru	View
Mus musculus	House mouse	New Island South, West Falkland	Falkland Islands (Malvinas)	View

[Download CSV](#)

Home | Contact Us | Data Use Policy | My Account | Log Out

© Zoological Society of London and WWF 2014

Living Planet Index

Home About Data Portal Projects Publications My Account

Keyword Search

Keyword

Refine Search

Advanced Search

Search Result: 50 record(s)

Map View List View

Species Name	Common Name	Location of Population	Countries	View
Panthera tigris	Tiger	Fujian province and adjacent areas, China	China	View
Panthera tigris	Tiger	Malay Peninsula, Malaysia	Malaysia	View
Panthera tigris	Tiger	Sumatra, Indonesia	Indonesia	View
Panthera tigris	Tiger	India, all States	India	View
Panthera tigris	Tiger	Sikhote Alin Zapovednik, Russia	Russian Federation	View
Panthera tigris	Tiger	Russian Far East	Russian Federation	View
Panthera tigris	Tiger	Upland Thung Yai Naresuan Wildlife Sanctuary	Thailand	View
Panthera tigris	Tiger	Bardia National Park, Nepal	Nepal	View
Panthera tigris	Tiger	Chitla Range, Rajaji National Park, India	India	View
Panthera tigris	Tiger	Luzovskiy Zapovednik, Russian Far East	Russian Federation	View
Panthera tigris	Tiger	Luzovskiy Raton, Russian Far East	Russian Federation	View
Panthera tigris	Tiger	Ussuriskiy Zapovednik, Russian Far East	Russian Federation	View
Panthera tigris	Tiger	Iman, Russian Far East	Russian Federation	View
Panthera tigris	Tiger	Bikin, Russian Far East	Russian Federation	View
Panthera tigris	Tiger	Borisovskoye Plateau, Russian Far East	Russian Federation	View
Panthera tigris	Tiger	Sandagino, Russian Far East	Russian Federation	View
Panthera tigris	Tiger	Khor, Russian Far East	Russian Federation	View
Panthera tigris	Tiger	Bozhenskoye Zapovednik, Russian Far East	Russian Federation	View
Panthera tigris	Tiger	Bozhokhetskoye Zapovednik, Russian Far East	Russian Federation	View
Panthera tigris	Tiger	Tigres Dom, Russian Far East	Russian Federation	View
Panthera tigris	Tiger	Matanski Zakumsk, Russian Far East	Russian Federation	View
Panthera tigris	Tiger	Ussuriski Raton, Russian Far East	Russian Federation	View
Panthera tigris	Tiger	Sikhote Alin Zapovednik, Russian Far East	Russian Federation	View
Panthera tigris	Tiger	Sineysk, Russian Far East	Russian Federation	View
Panthera tigris	Tiger	Ternoy Hunting Lease, Russian Far East	Russian Federation	View
Panthera tigris	Tiger	Nasip, Uttaranchal, India	India	View
Panthera tigris	Tiger	Corbett, Uttaranchal, India	India	View
Panthera tigris	Tiger	Dudhwa-Khori-Pilibhit, Uttar Pradesh, India	India	View
Panthera tigris	Tiger	Sahibga, Uttar Pradesh, India	India	View
Panthera tigris	Tiger	Sahibgaon-Vaishali, Uttar Pradesh, Bihar	India	View
Panthera tigris	Tiger	Nayabhan, India	India	View
Panthera tigris	Tiger	Madhya Pradesh, India	India	View
Panthera tigris	Tiger	Maharashtra, India	India	View
Panthera tigris	Tiger	Chhatargarh, India	India	View
Panthera tigris	Tiger	Orissa, India	India	View
Panthera tigris	Tiger	Andhra Pradesh, India	India	View
Panthera tigris	Tiger	Karnataka, India	India	View
Panthera tigris	Tiger	Tamil Nadu	India	View
Panthera tigris	Tiger	Kerala, India	India	View
Panthera tigris	Tiger	Assam, India	India	View
Panthera tigris	Tiger	Mizoram, India	India	View
Panthera tigris	Tiger	Huay Kha Kaeng Wildlife Sanctuaries	Thailand	View
Panthera tigris	Tiger	Sumatra, Indonesia	Indonesia	View
Panthera tigris	Tiger	Russian Far East	Russian Federation	View
Panthera tigris	Tiger	Western sector, Chitwan National Park, Nepal	Nepal	View
Panthera tigris	Tiger	Hongjiang Province	China	View
Panthera tigris	Tiger	Nagarahole reserve, Karnataka State, India	India	View
Panthera tigris	Tiger	Panna tiger reserve, India	India	View
Panthera tigris	Tiger	Bukit Barisan Sultan National Park	Indonesia	View
Panthera tigris	Tiger	Hanchuan National Nature Reserve, eastern Sichuan Province, China	China	View

[Download CSV](#)

Home | Contact Us | Data Use Policy | My Account | Log Out

© Zoological Society of London and WWF 2014

Panthera tigris =>

Living Planet Index



[Home](#) [About](#) [Data Portal](#) [Projects](#) [Publications](#) [My Account](#)

Keyword Search

Keyword

[Search](#)

[Clear Search](#)

[Load Search](#)

[Save Search](#)

[Refine Search](#)

Advanced Search

Search Result: 51 record(s)



[My Account](#) [Log Out](#)

© Zoological Society of London and WWF 2014

Living Planet Index

[Home](#) [About](#) [Data Portal](#) [Projects](#) [Publications](#) [My Account](#)

Keyword Search

Keyword

[Search](#)

[Clear Search](#)

[Load Search](#)

[Save Search](#)

[Refine Search](#)

Advanced Search

Search Result: 3 record(s)

[Map View](#) [List View](#)



[Home](#) [Contact Us](#) [Data Use Policy](#) [My Account](#) [Log Out](#)

© Zoological Society of London and WWF 2014

Rattus norvegicus



O GLOBO

Um só planeta


ASSINE

Um só planeta

Raias gigantes ameaçadas de extinção vivem na Baía de Guanabara e quantidade em meio a poluição impressiona biólogos

Ainda não existe explicação para tamanha concentração numa área tão castigada por décadas de poluição e descaso.







Indicators of the national population status of all regularly occurring native bird species in Canada and eight selected subgroups.



Abundância (= biomassa)





nature communications



Article


<https://doi.org/10.1038/s41467-023-37127-2>


Regional occupancy increases for wide-spread species but decreases for narrowly distributed species in metacommunity time series

We found that on average, large-ranged species tended to increase in occupancy through time, whereas small-ranged species tended to decrease.


Our findings provide evidence for systematic decreases in occupancy of small-ranged species, and that habitat protection could mitigate these losses in the face of environmental change.

Red List Index of species survival











13%

21%



10%



25%



6%



<1%



2%



40%



2%




<1%



4%



<1%




INSTITUTO CHICO MENDES
DE CONSERVAÇÃO DA BIODIVERSIDADE



LIVRO VERMELHO
DA FAUNA BRASILEIRA AMEAÇADA DE EXTINÇÃO

2018



	Grupo taxonômico	Táxons avaliados	Táxons ameaçados
Vertebrados	Mammalia	732	111 (15,1%)
	Aves	1.979	236 (11,9%)
	Reptilia	732	85 (11,6%)
	Amphibia	973	41 (4,2%)
	Actinopterygii (Marinhos)	1.201	43 (3,6%)
	Actinopterygii (Continentais)	3.130	311 (9,9%)
	Dipneusti	1	0
	Elasmobranchii (Marinhos)	146	54 (36,9%)
	Elasmobranchii (Continentais)	17	1 (5,8%)
	Holocephali	6	0
Invertebrados	Myxini	5	1 (20%)
	Total vertebrados	8.922	883 (9,9%)
	Marinhos	661	29
	Águas continentais	248	37
	Terrestres	2.423	233
	Total invertebrados	3.332	299
	TOTAL	12.254	1.182



Cycloramphus diringshofeni Bokermann, 1957

Categoria de risco de extinção e critérios

Criticamente em Perigo (CR) B1ab(iii)

Justificativa

Cycloramphus diringshofeni é endêmica do Brasil, conhecida apenas da localidade-tipo, no município de São Bento do Sul, estado de Santa Catarina. Não há registros de coleta da espécie nos últimos 50 anos. Sua extensão de ocorrência estimada é de 4 km². Na área onde a espécie ocorre há declínio continuado da qualidade do *habitat* devido, principalmente, à ocupação urbana e à conversão de áreas naturais em plantações de *Pinus* spp. e banana. Por essas razões, *Cycloramphus diringshofeni* foi categorizada como Criticamente em Perigo (CR) pelo critério B1ab(iii).

Outras avaliações

Avaliação nacional anterior ¹¹⁶	DD
Justificativa para mudança de categoria	Novas ou melhores informações disponíveis
Listas estaduais de espécies ameaçadas	Não consta
Avaliação global ⁹⁴	DD

Distribuição geográfica

Cycloramphus diringshofeni é endêmica do Brasil e apresenta distribuição restrita a localidade-tipo, município de São Bento do Sul, estado de Santa Catarina¹¹². Sua extensão de ocorrência é de 4 km², calculada pelo método do esquadramento, utilizando-se uma *grid* de 2 km de lado.





Species

Our Work

+ [The IUCN Red List of Threatened Species](#)

+ [Influencing policy](#)

+ [Action on the ground](#)

+ [Sustainable use and trade](#)

[Amphibians](#)

[Birds](#)

+ [Freshwater biodiversity](#)

[Fungi](#)

– [Invasive species](#)

+ [The Honolulu Challenge on Invasive Alien Species](#)

[EICAT](#)

[Mammals](#)

[Marine](#)

+ [Plants](#)

Invasive species

What is an Invasive Alien Species?

An alien species is a species introduced by humans – either intentionally or accidentally - outside of its natural past or present distribution, however not all alien species have negative impacts, and it is estimated that between 5% and 20% of all alien species become problematic. It is these species that are termed ‘*invasive alien species*’ (IAS).

“An invasive alien species (IAS) is a species that is established outside of its natural past or present distribution, whose introduction and/or spread threaten biological diversity” Convention on Biological Diversity.

What are their impacts?

Invasive alien species are a major driver of biodiversity loss. In fact, an analysis of the IUCN Red List shows that they are the second most common threat associated with species that have gone completely extinct, and are the most common threat associated with extinctions of amphibians, reptiles and mammals.

An example is the Micronesian Kingfisher *Todiramphus cinnamominus* which was endemic to the island of Guam, but following predation by invasive alien snakes (Brown tree snake *Boiga irregularis*) it became Extinct in the Wild in 1986 when the last remaining wild birds were taken into captivity for captive breeding. In fact the Brown tree snake has caused many extinctions on Guam, including the local extinction of over half of Guam’s native bird and lizard species as well as two out of three of Guam’s native bat species.

Links

[Invasive Species Specialist Group](#)

[Invasive Species of the Week](#)

[The Global Invasive Species Database](#)


[The Global Register of Introduced and Invasive Species](#)

[The IUCN Red List of Threatened Species](#)



Escala local: espécies exóticas não se extinguem?



[illegible]



A CONTROVERSIAL LEGACY COLOMBIA'S HIPPOS

4 – Desmistificar conceitos fixistas que permeiam discurso **ambientalista**

Outras duas perguntas:

1: Por que consideramos a “6ª extinção” algo a ser contido?


2: O que significa conter a “perda” de biodiversidade?



Conserving the diversity of life


Outras duas perguntas:

1: Por que seria a “6ª extinção” algo a ser contido?





Outras duas perguntas:

1: Por que seria a “6ª extinção” algo a ser contido?




Outras duas perguntas:

1: Por que seria a “6ª extinção” algo a ser contido?









**A fumaça das chaminés
é o hálito da Rússia soviética**

Homo sum, humani a mi nihil alienum

Terêncio








Conserving the diversity of life



Conserving the diversity of life





4 – Desmistificar conceitos fixistas que permeiam discurso **ambientalista**

Outras duas perguntas:

2: O que significa conter a perda da biodiversidade?





LUIS FERNANDO VERÍSSIMO

Albert Speer fala
com as flores



















COUNTDOWN
2010
SAVE BIODIVERSITY







Biodiversity

WE ARE ALL IN THIS TOGETHER

