

Congenital Adrenal Hyperplasia

New M, Lekarev O, Lin-Su K, et al.

INTRODUCTION

Congenital adrenal hyperplasia (CAH) refers to a group of disorders that arise from defective steroidogenesis. The production of cortisol in the zona fasciculata of the adrenal cortex occurs in five major enzyme-mediated steps. CAH results from deficiency in any one of these enzymes. Impaired cortisol synthesis leads to chronic elevations of ACTH via the negative feedback system, causing overstimulation of the adrenal cortex and resulting in hyperplasia and oversecretion of the precursors to the enzymatic defect. The five forms of CAH are summarized in **Table 1**. Impaired enzyme function at each step of adrenal cortisol biosynthesis leads to a unique combination of elevated precursors and deficient products. The most common enzyme deficiency that accounts for more than 90% of all CAH cases is 21-hydroxylase deficiency [1].

Epidemiology

Data from close to 6.5 million newborn screenings worldwide indicate that classical CAH occurs in 1:13,000 to 1:15,000 live births [2]. It is estimated that 75% of patients have the salt-wasting phenotype [3]. Non-classical 21OHD CAH (NC21OHD) is more common. The incidence in the heterogeneous population of New York City is 1 in 100, making NC21OHD the most frequent autosomal recessive disorder in humans. NC21OHD is particularly prevalent in certain populations, showing a high ethnic specificity. In the Ashkenazi Jewish population, 1 in 3 are carriers of the allele, and 1 in 27 are affected with the disorder [4, 5, 6]. CAH owing to 11 β -hydroxylase deficiency (11 β -OHD) is the second most common cause of CAH, accounting for 5-8% of all cases [7]. It occurs in 1 of every 100,000 live births in the general population [8] and is more common in some populations of North African origin [9]. In Moroccan Jews, for example, the disease incidence was initially estimated to be 1 in 5,000 live births [10]; subsequently, it was shown to occur less frequently [11], but remains more common than in other populations. The other forms of CAH are considered rare diseases and the incidence is unknown in the general population.

Pathophysiology

Adrenal steroidogenesis occurs in three major pathways: **glucocorticoids, mineralocorticoids, and sex steroids** as shown in **Figure 1**. The adrenal gland secretions suggest that the adrenal acts as three separate glands: **zona glomerulosa, zona fasciculata, zona reticularis**. [12, 13]. The hypothalamic-pituitary-adrenal feedback system is mediated through the circulating level of plasma cortisol by negative feedback of cortisol on CRF and ACTH secretion. Therefore, **any CAH condition that results in a decrease in cortisol secretion** leads to increased ACTH production, which in turn stimulates (1) excessive synthesis of adrenal products in those pathways unimpaired by the enzyme deficiency and (2) a build-up of precursor molecules in pathways blocked by the enzyme deficiency.




Figure 1

illustrates adrenal steroidogenesis: Five enzymatic steps necessary for cortisol production are shown in numbers. 1= 20, 22 desmolase, 2= 17 hydroxylase (17-OH), 3=3β-hydroxysteroid dehydrogenase (3β HSD), 4=21 hydroxylase (21-OHD), 5=11β hydroxylase (11-OH) In the first step of adrenal stereoidgeneis, cholesterol enters mitochondria via a carrier protein called StAR. ACTH stimulates cholesterol cleavage, the rate limiting step of adrenal steroidogenesis.

The clinical symptoms of the five different forms of CAH result from the particular hormones that are deficient and those that are produced in excess as outlined in **Table 1**. In the most common form 21OHD-CAH, the function of 21-hydroxylating cytochrome P450 is inadequate, creating a block in P450 cortisol production pathway. This leads to an accumulation of 17-hydroxyprogesterone (17-OHP), a precursor adjacent to the 21-hydroxylation step. Excess 17-OHP is then shunted into the intact androgen pathway, where the 17,20-lyase enzyme converts the 17-OHP to Δ4-androstenedionea major adrenal androgen. Mineralocorticoid deficiency is a feature of the most severe form of the disease called salt wasting CAH. The enzyme defect in the non-classical form of 21OHD CAH is mild and salt wasting in this mild form of the disease is not evident. The analogy of all other enzyme deficiencies in terms of precursor retention and product deficiencies are shown in **Table 1**.

Table 1
Summary of the Clinical, Hormonal, and Genetic Features of Steroidogenic Defects [1].

Condition	Class	Abnormality	Genotype	Mineralocorticoid	Glucocorticoid	Sex
Liquid CAH	Congenital	20,22 desmolase	autosomal recessive	absent	absent	male
Liquid CAH	Congenital	17-OH	autosomal recessive	absent	absent	male
21-OHD	Congenital	21-OH	autosomal recessive	absent	absent	male
11-OH	Congenital	11-OH	autosomal recessive	absent	absent	male
17βHSD	Congenital	17βHSD	autosomal recessive	absent	absent	male
17,20 lyase	Congenital	17,20 lyase	autosomal recessive	absent	absent	male
3βHSD	Congenital	3βHSD	autosomal recessive	absent	absent	male
StAR	Congenital	StAR	autosomal recessive	absent	absent	male
17αHSD	Congenital	17αHSD	autosomal recessive	absent	absent	male
17,20 lyase	Congenital	17,20 lyase	autosomal recessive	absent	absent	male
17βHSD	Congenital	17βHSD	autosomal recessive	absent	absent	male
17,20 lyase	Congenital	17,20 lyase	autosomal recessive	absent	absent	male
3βHSD	Congenital	3βHSD	autosomal recessive	absent	absent	male
StAR	Congenital	StAR	autosomal recessive	absent	absent	male
17αHSD	Congenital	17αHSD	autosomal recessive	absent	absent	male
17,20 lyase	Congenital	17,20 lyase	autosomal recessive	absent	absent	male
17βHSD	Congenital	17βHSD	autosomal recessive	absent	absent	male
17,20 lyase	Congenital	17,20 lyase	autosomal recessive	absent	absent	male
3βHSD	Congenital	3βHSD	autosomal recessive	absent	absent	male
StAR	Congenital	StAR	autosomal recessive	absent	absent	male
17αHSD	Congenital	17αHSD	autosomal recessive	absent	absent	male
17,20 lyase	Congenital	17,20 lyase	autosomal recessive	absent	absent	male
17βHSD	Congenital	17βHSD	autosomal recessive	absent	absent	male
17,20 lyase	Congenital	17,20 lyase	autosomal recessive	absent	absent	male
3βHSD	Congenital	3βHSD	autosomal recessive	absent	absent	male
StAR	Congenital	StAR	autosomal recessive	absent	absent	male
17αHSD	Congenital	17αHSD	autosomal recessive	absent	absent	male
17,20 lyase	Congenital	17,20 lyase	autosomal recessive	absent	absent	male
17βHSD	Congenital	17βHSD	autosomal recessive	absent	absent	male
17,20 lyase	Congenital	17,20 lyase	autosomal recessive	absent	absent	male
3βHSD	Congenital	3βHSD	autosomal recessive	absent	absent	male
StAR	Congenital	StAR	autosomal recessive	absent	absent	male
17αHSD	Congenital	17αHSD	autosomal recessive	absent	absent	male
17,20 lyase	Congenital	17,20 lyase	autosomal recessive	absent	absent	male
17βHSD	Congenital	17βHSD	autosomal recessive	absent	absent	male
17,20 lyase	Congenital	17,20 lyase	autosomal recessive	absent	absent	male
3βHSD	Congenital	3βHSD	autosomal recessive	absent	absent	male
StAR	Congenital	StAR	autosomal recessive	absent	absent	male
17αHSD	Congenital	17αHSD	autosomal recessive	absent	absent	male
17,20 lyase	Congenital	17,20 lyase	autosomal recessive	absent	absent	male
17βHSD	Congenital	17βHSD	autosomal recessive	absent	absent	male
17,20 lyase	Congenital	17,20 lyase	autosomal recessive	absent	absent	male
3βHSD	Congenital	3βHSD	autosomal recessive	absent	absent	male
StAR	Congenital	StAR	autosomal recessive	absent	absent	male
17αHSD	Congenital	17αHSD	autosomal recessive	absent	absent	male
17,20 lyase	Congenital	17,20 lyase	autosomal recessive	absent	absent	male
17βHSD	Congenital	17βHSD	autosomal recessive	absent	absent	male
17,20 lyase	Congenital	17,20 lyase	autosomal recessive	absent	absent	male
3βHSD	Congenital	3βHSD	autosomal recessive	absent	absent	male
StAR	Congenital	StAR	autosomal recessive	absent	absent	male
17αHSD	Congenital	17αHSD	autosomal recessive	absent	absent	male
17,20 lyase	Congenital	17,20 lyase	autosomal recessive	absent	absent	male
17βHSD	Congenital	17βHSD	autosomal recessive	absent	absent	male
17,20 lyase	Congenital	17,20 lyase	autosomal recessive	absent	absent	male
3βHSD	Congenital	3βHSD	autosomal recessive	absent	absent	male
StAR	Congenital	StAR	autosomal recessive	absent	absent	male
17αHSD	Congenital	17αHSD	autosomal recessive	absent	absent	male
17,20 lyase	Congenital	17,20 lyase	autosomal recessive	absent	absent	male
17βHSD	Congenital	17βHSD	autosomal recessive	absent	absent	male
17,20 lyase	Congenital	17,20 lyase	autosomal recessive	absent	absent	male
3βHSD	Congenital	3βHSD	autosomal recessive	absent	absent	male
StAR	Congenital	StAR	autosomal recessive	absent	absent	male
17αHSD	Congenital	17αHSD	autosomal recessive	absent	absent	male
17,20 lyase	Congenital	17,20 lyase	autosomal recessive	absent	absent	male
17βHSD	Congenital	17βHSD	autosomal recessive	absent	absent	male
17,20 lyase	Congenital	17,20 lyase	autosomal recessive	absent	absent	male
3βHSD	Congenital	3βHSD	autosomal recessive	absent	absent	male
StAR	Congenital	StAR	autosomal recessive	absent	absent	male
17αHSD	Congenital	17αHSD	autosomal recessive	absent	absent	male
17,20 lyase	Congenital	17,20 lyase	autosomal recessive	absent	absent	male
17βHSD	Congenital	17βHSD	autosomal recessive	absent	absent	male
17,20 lyase	Congenital	17,20 lyase	autosomal recessive	absent	absent	male
3βHSD	Congenital	3βHSD	autosomal recessive	absent	absent	male
StAR	Congenital	StAR	autosomal recessive	absent	absent	male
17αHSD	Congenital	17αHSD	autosomal recessive	absent	absent	male
17,20 lyase	Congenital	17,20 lyase	autosomal recessive	absent	absent	male
17βHSD	Congenital	17βHSD	autosomal recessive	absent	absent	male
17,20 lyase	Congenital	17,20 lyase	autosomal recessive	absent	absent	male
3βHSD	Congenital	3βHSD	autosomal recessive	absent	absent	male
StAR	Congenital	StAR	autosomal recessive	absent	absent	male
17αHSD	Congenital	17αHSD	autosomal recessive	absent	absent	male
17,20 lyase	Congenital	17,20 lyase	autosomal recessive	absent	absent	male
17βHSD	Congenital	17βHSD	autosomal recessive	absent	absent	male
17,20 lyase	Congenital	17,20 lyase	autosomal recessive	absent	absent	male
3βHSD	Congenital	3βHSD	autosomal recessive	absent	absent	male
StAR	Congenital	StAR	autosomal recessive	absent	absent	male
17αHSD	Congenital	17αHSD	autosomal recessive	absent	absent	male
17,20 lyase	Congenital	17,20 lyase	autosomal recessive	absent	absent	male
17βHSD	Congenital	17βHSD	autosomal recessive	absent	absent	male
17,20 lyase	Congenital	17,20 lyase	autosomal recessive	absent	absent	male
3βHSD	Congenital	3βHSD	autosomal recessive	absent	absent	male
StAR	Congenital	StAR	autosomal recessive	absent	absent	male
17αHSD	Congenital	17αHSD	autosomal recessive	absent	absent	male
17,20 lyase	Congenital	17,20 lyase	autosomal recessive	absent	absent	male
17βHSD	Congenital	17βHSD	autosomal recessive	absent	absent	male
17,20 lyase	Congenital	17,20 lyase	autosomal recessive	absent	absent	male
3βHSD	Congenital	3βHSD	autosomal recessive	absent	absent	male
StAR	Congenital	StAR	autosomal recessive	absent	absent	male
17αHSD	Congenital	17αHSD	autosomal recessive	absent	absent	male
17,20 lyase	Congenital	17,20 lyase	autosomal recessive	absent	absent	male
17βHSD	Congenital	17βHSD	autosomal recessive	absent	absent	male
17,20 lyase	Congenital	17,20 lyase	autosomal recessive	absent	absent	male
3βHSD	Congenital	3βHSD	autosomal recessive	absent	absent	male
StAR	Congenital	StAR	autosomal recessive	absent	absent	male
17αHSD	Congenital	17αHSD	autosomal recessive	absent	absent	male
17,20 lyase	Congenital	17,20 lyase	autosomal recessive	absent	absent	male
17βHSD	Congenital	17βHSD	autosomal recessive	absent	absent	male
17,20 lyase	Congenital	17,20 lyase	autosomal recessive	absent	absent	male
3βHSD	Congenital	3βHSD	autosomal recessive	absent	absent	male
StAR	Congenital	StAR	autosomal recessive	absent	absent	male
17αHSD	Congenital	17αHSD	autosomal recessive	absent	absent	male
17,20 lyase	Congenital	17,20 lyase	autosomal recessive	absent	absent	male
17βHSD	Congenital	17βHSD	autosomal recessive	absent	absent	male
17,20 lyase	Congenital	17,20 lyase	autosomal recessive	absent	absent	male
3βHSD	Congenital	3βHSD	autosomal recessive	absent	absent	male
StAR	Congenital	StAR	autosomal recessive	absent	absent	male
17αHSD	Congenital	17αHSD	autosomal recessive	absent	absent	male
17,20 lyase	Congenital	17,20 lyase	autosomal recessive	absent	absent	male
17βHSD	Congenital	17βHSD	autosomal recessive	absent	absent	male
17,20 lyase	Congenital	17,20 lyase	autosomal recessive	absent	absent	male
3βHSD	Congenital	3βHSD	autosomal recessive	absent	absent	male
StAR	Congenital	StAR	autosomal recessive	absent	absent	male
17αHSD	Congenital	17αHSD	autosomal recessive	absent	absent	male
17,20 lyase	Congenital	17,20 lyase	autosomal recessive	absent	absent	male
17βHSD	Congenital	17βHSD	autosomal recessive	absent	absent	male
17,20 lyase	Congenital	17,20 lyase	autosomal recessive	absent	absent	male
3βHSD	Congenital	3βHSD	autosomal recessive	absent	absent	male
StAR	Congenital	StAR	autosomal recessive	absent	absent	male
17αHSD	Congenital	17αHSD	autosomal recessive	absent	absent	male
17,20 lyase	Congenital	17,20 lyase	autosomal recessive	absent	absent	male
17βHSD	Congenital	17βHSD	autosomal recessive	absent	absent	male
17,20 lyase	Congenital	17,20 lyase	autosomal recessive	absent	absent	male
3βHSD	Congenital	3βHSD	autosomal recessive	absent	absent	male
StAR	Congenital	StAR	autosomal recessive	absent	absent	male
17αHSD	Congenital	17αHSD	autosomal recessive	absent	absent	male
17,20 lyase	Congenital	17,20 lyase	autosomal recessive	absent	absent	male
17βHSD	Congenital	17βHSD	autosomal recessive	absent	absent	male
17,20 lyase	Congenital	17,20 lyase	autosomal recessive	absent	absent	male
3βHSD	Congenital	3βHSD	autosomal recessive	absent	absent	male
StAR	Congenital	StAR	autosomal recessive	absent	absent	male
17αHSD	Congenital	17αHSD	autosomal recessive	absent	absent	male
17,20 lyase	Congenital	17,20 lyase	autosomal recessive	absent	absent	male
17βHSD	Congenital	17βHSD	autosomal recessive	absent	absent	male
17,20 lyase	Congenital	17,20 lyase	autosomal recessive	absent	absent	male
3βHSD	Congenital	3βHSD	autosomal recessive	absent	absent	male
StAR	Congenital	StAR	autosomal recessive	absent	absent	male
17αHSD	Congenital	17αHSD	autosomal recessive	absent	absent	male
17,20 lyase	Congenital	17,20 lyase	autosomal recessive	absent	absent	male
17βHSD	Congenital	17βHSD	autosomal recessive	absent	absent	male
17,20 lyase	Congenital	17,20 lyase	autosomal recessive	absent	absent	male
3βHSD	Congenital	3βHSD	autosomal recessive	absent	absent	male
StAR	Congenital	StAR	autosomal recessive	absent	absent	male
17αHSD	Congenital	17αHSD	autosomal recessive	absent	absent	male
17,20 lyase	Congenital	17,20 lyase	autosomal recessive	absent	absent	male
17βHSD	Congenital	17βHSD	autosomal recessive	absent	absent	male
17,20 lyase	Congenital	17,20 lyase	autosomal recessive	absent	absent	male
3βHSD	Congenital	3βHSD	autosomal recessive	absent	absent	male
StAR	Congenital	StAR	autosomal recessive	absent	absent	male
17αHSD	Congenital	17αHSD	autosomal recessive	absent	absent	male
17,20 lyase	Congenital	17,20				

Females with Classical 21OHD and 11 β -hydroxylase deficiency CAH generally present at birth with virilization of their genitalia. Adrenocortical function begins at the 7th week of gestation [14]; thus, a female fetus with classical CAH is exposed to adrenal androgens at the critical time of sexual differentiation (between 9 to 15 weeks gestational age). Androgens masculinize external female genitalia causing genital ambiguity including: clitoral enlargement, fusion and scrotalization of the labial folds, and rostral migration of the urethral/vaginal perineal orifice, placing the phallus in the male position. The degree of genital virilization may range from mild clitoral enlargement alone to, in rare cases, a penile urethra (Prader V genitalia). Degrees of genital virilization are classified into five Prader stage [15] (see Figure 2).




Figure 2 Different degrees of virilization according to the scale developed by Prader (15) Stage I: clitoromegaly without labial fusion Stage II: clitoromegaly and posterior labial fusion Stage III: greater degree of clitoromegaly, single perineal urogenital orifice, and almost complete labial fusion Stage IV: increasingly phallic clitoris, urethra-like urogenital sinus at base of clitoris, and complete labial fusion Stage V: penile clitoris, urethral meatus at tip of phallus, and scrotum-like labia (appear like males without palpable gonads)

Internal genitalia

In contrast to the virilization of the external genitalia, internal female genitalia, the uterus, fallopian tubes and ovaries, develop normally. Females with CAH do not have testicular tissue and therefore do not produce anti-Müllerian hormone (AMH), which is produced by the testicular Sertoli cells. Internal female structures are Müllerian derivatives and are not androgen responsive. Therefore, the affected female born with virilized external genitalia but normal female internal genitalia and have the possibility of normal fertility [1].

Postnatal effects and growth

Deficient postnatal treatment in boys and girls results in continued exposure to excessive androgens, causing progressive penile or clitoral enlargement, the development of premature pubic hair (pubarche), axillary hair, acne, and impaired fertility. Advanced somatic and epiphyseal development occurs with rapid growth during childhood. This rapid linear growth is usually accompanied by premature epiphyseal maturation and closure, resulting in a final adult height that is below that expected from parental heights (on average -1.1 to -1.5 SD below the midparental target height) [16, 17]. This is on average 10 cm below the mid-parental height [18]. On the other hand, poor growth can occur in patients with 21OHD as a result of excess glucocorticoid treatment. Short stature occurs even in patients with good hormonal adrenal control. A study of growth hormone therapy alone or in combination with a GnRH analog in CAH patients with compromised height prediction showed improvement in short- and long-term growth to reduce the height deficit [18].

Puberty

In the majority of patients treated adequately from early life, the onset of puberty in both girls and boys with classical 21OHD occurs at the expected chronological age. A careful study showed that the mean ages at onset of puberty in both males and females were somewhat younger than the general population, but did not differ significantly among the three forms of 21OHD [19].

In those who are inadequately treated, advanced epiphyseal development can lead to central precocious puberty. In those with advanced

body maturation at the initial presentation, such as in simple virilizing males, the exposure to elevated androgens followed by the suddenly decreased androgen levels after initiation of glucocorticoid treatment may cause an early activation of the hypothalamic-pituitary-gonadal axis. Studies suggest that excess adrenal androgens (aromatized to estrogens) inhibit the pubertal pattern of gonadotropin secretion by the hypothalamic-pituitary axis. This inhibition, via a negative feedback effect, can be reversed by glucocorticoid treatment [19, 20].

Following the onset of puberty, in a majority of successfully treated patients, the milestones of further development of secondary sex characteristics in general appear to be normal [19, 20]. In adolescents and adults, signs of hyperandrogenism may include male-pattern alopecia (temporal balding) and acne. Female patients may develop hirsutism and menstrual irregularities. Although the expected age of menarche may be delayed in females with classical CAH [21], when adequately treated many have regular menses after menarche [22]. Menstrual irregularity and secondary amenorrhea with or without hirsutism occur in a subset of postmenarchal females, especially those in poor hormonal control. Primary amenorrhea or delayed menarche may occur if a woman with classical CAH is untreated, inadequately treated, or over treated with glucocorticoids [23]. In addition, women with CAH may develop Polycystic Ovarian Syndrome (PCOS) [24].

Gender role behavior and cognition

Prenatal androgen exposure in females affected with the classic forms of 21OHD CAH not only has a masculinizing effect on the development of the external genitalia, but also on childhood behavior. Both physical and behavioral masculinization were related to genotype, indicating that behavioral masculinization in childhood is a consequence of prenatal androgen exposure. Further, changes in childhood play behavior is correlated with reduced female gender satisfaction and reduced heterosexual interest in adulthood. Prenatal androgen exposure is related to a decrease in self-reported femininity in dose response manner in adulthood [25]. Affected adult females are more likely to have gender dysphoria, and experience less heterosexual interest and reduced satisfaction with the assignment to the female sex. In contrast to females, males affected with CAH do not show a general alteration in childhood play behavior, core gender identity and sexual orientation [26]. The rates of bisexual and homosexual orientation were increased in women with all forms of 21OHD CAH. They were found to correlate with the degree of prenatal androgenization. Of interest, bisexual/homosexual orientation was correlated with global measures of masculinization of nonsexual behavior and predicted independently by the degree of both prenatal androgenization and masculinization of childhood behavior [27].

With regards to cognitive abilities, such as visuospatial/motor ability and handedness, the effect of prenatal androgen exposure continues to be elucidated. A recent study found males and females with CAH scored higher than their siblings of the same sex in measures of visual special processing suggesting that androgens affect spatial ability [28]. The effect of CAH on intelligence is controversial. A more recent careful study showed no evidence of intellectual deficit in either females or males with CAH. Intelligence was not significantly associated with disease characteristics [29].

Fertility

Difficulty with fertility in females with CAH may arise for various reasons, including anovulation, secondary polycystic ovarian syndrome, irregular menses, non-suppressible serum progesterone levels, or an inadequate introitus. Fertility is reduced in salt-wasting 21OHD. In a retrospective survey of fertility rates in a large group of females with classical CAH, simple virilizers were shown to be more likely to become pregnant and carry the pregnancy to term than salt-wasters. Adequate glucocorticoid therapy is an important variable with respect to fertility outcome. The development of PCOS in CAH patients is not uncommon and may be related to both prenatal and postnatal excess androgen exposure, which can affect the hypothalamic-pituitary-gonadal axis. An inadequate vaginal introitus can affect up to a third of classical CAH adult females. Since vaginal dilation is needed to maintain good patency, vaginoplasty is delayed until sexual intercourse is regular or when the patient can assume responsibility for vaginal dilatation [30].

Males with CAH, particularly if poorly treated, may have reduced sperm counts and low testosterone as a result of small testes due to suppression of gonadotropins and sometimes intra-testicular adrenal rests [31, 32]. All of these complications may result in diminished fertility. In male patients with classical CAH, several long-term studies indicate that those who have been adequately treated undergo normal pubertal development, have normal testicular function, and normal spermatogenesis and fertility [33, 34]. However, small testes and aspermia can occur in patients as a result of inadequately controlled disease [35, 36]. Testicular adrenal rest tumor can lead to end

stage damage of testicular parenchyma, most probably as a result of longstanding obstruction of the seminiferous tubules [37]. In contrast, some investigators have reported normal testicular maturation as well as normal spermatogenesis and fertility in patients who had never received glucocorticoid treatment [38].

Studies demonstrate that postpubertal males with inadequately treated CAH are at a very high risk to develop hyperplastic nodular testes. In one study, almost all these patients were found to have adenomatous adrenal rests within the testicular tissue, as indicated by the presence of specific 11 β -hydroxylated steroids in the blood from gonadal veins [39]. These tumors have been reported to be ACTH dependent and to regress following adequate steroid therapy [40, 41, 42, 43, 44]. These testicular adrenal rests are more frequent in males with salt-wasting CAH and are associated with an increased risk of infertility [31, 45]. Regular testicular examination and periodic testicular ultrasonography are recommended for early detection of testicular lesions.

Salt-wasting 21-hydroxylase deficiency

When the deficiency of 21-hydroxylase is severe, adrenal aldosterone secretion is not sufficient for sodium reabsorption by the distal renal tubules, and individuals suffer from salt wasting as well as cortisol deficiency and androgen excess. Infants with renal salt wasting have poor feeding, weight loss, failure to thrive, vomiting, dehydration, hypotension, hyponatremia, and hyperkalemic metabolic acidosis progressing to adrenal crisis (azotemia, vascular collapse, shock, and death). Adrenal crisis can occur as early as age one to four weeks. The salt wasting is presumed to result from inadequate secretion of salt-retaining steroids, primarily aldosterone. In addition, hormonal precursors of the 21-OH enzyme may act as antagonists to mineralocorticoid action in the sodium-conserving mechanism of the immature newborn renal tubule [46, 47, 48].

Affected males who are not detected in a newborn screening program are at high risk for a salt-wasting adrenal crisis because their normal male genitalia do not alert medical professionals to their condition. They are often discharged from the hospital after birth without diagnosis and experience a salt-wasting crisis at home. On the other hand, salt wasting females are born with ambiguous genitalia that trigger the diagnostic process and appropriate treatment. It is important to recognize that the extent of genital virilization may not differ among the two forms of classical CAH, the simple virilizing and the salt-wasting form. Thus, even a mildly virilized newborn with 21OHD should be observed carefully for signs of a potentially life-threatening crisis within the first few weeks of life.

The difference between salt-wasting and simple virilizing form of 21OHD is the quantitative difference in activity of the 21-hydroxylase enzyme, which results from specific mutations. In vitro expression studies show that as little as 1% of 21-hydroxylase activity is sufficient to synthesize enough aldosterone to prevent significant salt wasting [49]. It has been observed that an aldosterone biosynthetic defect apparent in infancy may ameliorate with age [50, 51]. Speiser et al. reported a spontaneous partial recovery of aldosterone biosynthesis in an adult patient with a homozygous deletion of the *CYP21A2* gene who had documented severe salt wasting in infancy [52]. Therefore, it is desirable to follow the sodium and mineralocorticoid requirements carefully by measuring PRA in patients who have been diagnosed in the neonatal period as salt wasters.

Simple-virilizing 21-hydroxylase deficiency

The salient features of classical simple virilizing 21OHD are prenatal virilization and progressive postnatal masculinization with rapid somatic growth and advanced epiphyseal maturation leading to early epiphyseal closure and likely short stature. There is no evidence of mineralocorticoid deficiency in this disorder.

Diagnosis at birth of a female with simple virilizing CAH is usually made immediately because of the apparent genital ambiguity. Since the external genitalia are not affected in newborn males, hyperpigmentation may be the only clue suggesting increased ACTH secretion and cortisol deficiency. Diagnosis at birth in males thus rests on prenatal or newborn screening. If a female is not begun on glucocorticoid replacement therapy post-natally, her genitalia may continue to virilize due to continued excess adrenal androgens, and pseudoprecocious puberty can occur. In patients with salt-wasting 21OHD, signs of hyperandrogenism in children affected with CAH include early onset of facial, axillary, and pubic hair, adult body odor, and rapid somatic growth and bone age advancement, leading to short stature in adulthood. The same issues as discussed above related to puberty, fertility, behavior and cognition apply to patients with simple-virilizing 21OHD [1].)

Non-classical 21-hydroxylase deficiency

Non-classical 21OHD (NC-21OHD), previously known as late-onset 21OHD, is much more common than the classical form, with an incidence as high as 1:27 in Ashkenazi Jews [4]. Individuals with the non-classical (NC) form of 21OHD have only mild to moderate enzyme deficiency and present postnatally, eventually developing signs of hyperandrogenism. Females with NC-CAH do not have virilized genitalia at birth.

NC-CAH may present at any age after birth with a variety of hyperandrogenic symptoms. This form of CAH results from a mild deficiency of the 21-hydroxylase enzyme. **Table 2** summarizes main clinical characteristics of all forms of 21OHD CAH. While serum cortisol concentration is typically low in patients with the classic form of the disease, it is usually normal in patients with NC 21OHD. Similar to classical CAH, NC-CAH may cause premature development of pubic hair, advanced bone age and accelerated linear growth velocity in both males and females. Severe cystic acne has also been attributed to NC-CAH [53, 54].

Enzyme	21-OH Deficiency	Classic	Non-Classic
Clinical features	Variable to absent	Absent	Absent
Growth and development	Normal	Delayed	Normal
Salt wasting	10% of all individuals	Absent	Absent

Table 2

Clinical Features in Individuals with Classic and Non-Classic 21-Hydroxylase Deficiency in the Untreated Form.

Women may present with a variety of symptoms of androgen excess which may be highly variable and organ-specific, including hirsutism, temporal baldness, acne and infertility. Menarche in females may be normal or delayed, and secondary amenorrhea is a frequent occurrence. Further masculinization may include hirsutism, male habitus, deepening of the voice, or male-pattern alopecia (temporal recession). Polycystic ovarian syndrome may also be seen as a secondary complication in these patients. Possible reasons for the development of PCOS include reprogramming of the hypothalamic-pituitary-gonadal axis from prenatal exposure to androgens, or chronic levels of excess adrenal androgens that disrupt gonadotropin release and have direct effects on the ovary, ultimately leading to the formation of cysts. Because of the overlap of hyperandrogenic symptoms, it is important to consider NC 21OHD in a patient diagnosed with PCO [55, 56].

In adult males, early balding, acne, or infertility may prompt the diagnosis of NC-CAH. A highly reliable constellation of physical signs of adrenal androgen excess is the presence of pubic hair, enlarged phallus, and relatively small testes. Males may have small testes compared to the phallus, which results from suppression of the hypothalamic-pituitary-gonadal axis from adrenal androgens. They may also develop intra-testicular adrenal rests, which can cause infertility, although some untreated men have been fertile [34]. Symptoms in adult males with NC-CAH may be limited to short stature or oligozoospermia and impaired fertility.

A subset of individuals with NC-21OHD are completely asymptomatic when detected (usually as part of a family study or evaluation for infertility), but it is thought, based on longitudinal follow-up of such patients, that symptoms of hyperandrogenism may wax and wane with time. The presence of 21OHD can also be discovered during the evaluation of an incidental adrenal mass [57]. One study showed that an increased incidence of adrenal incidentalomas has been found, which was reported as high as 82% in patients with 21OHD and up to 45% in subjects heterozygous for 21OHD mutations. This probably arises from hyperplastic tissue areas and does not require surgical intervention [58]. Overall, however, CAH is an uncommon cause of incidentalomas, accounting for less than 1% in one series [59, 60].

OTHER FORMS OF CONGENITAL ADRENAL HYPERPLASIA

11-β hydroxylase deficiency (1) Chapter 3D

Virilization and low renin hypertension are the prominent clinical features of 11β hydroxylase deficiency (11β-OHD) [61]. The virilizing signs and symptoms of this disorder are similar to classical 21OHD. Despite failure of aldosterone production, overproduction of DOC, *in vivo* a less potent mineralocorticoid, causes salt retention and hypertension. Elevated blood pressure is usually not identified until later in childhood or in adolescence, although its appearance in an infant 3 months of age has been documented [62]. In addition, hypertension correlates variably with biochemical values, and clinical signs of mineralocorticoid excess and the degree of virilization are not well correlated. Some severely virilized females are normotensive, whereas mildly virilized patients may experience severe hypertension leading to fatal vascular accidents [63, 64]. Complications of long standing uncontrolled hypertension, including

cardiomyopathy, retinal vein occlusion and blindness have been reported in 11 β -OHD patients [65, 66]. Potassium depletion develops concomitantly with sodium retention, but hypokalemia is variable. Renin production is suppressed secondary to mineralocorticoid-induced sodium retention and volume expansion. Aldosterone production is low secondary to low serum potassium and low plasma rennin [61]. (See chapter on Endocrine Hypertension in Childhood).

A mild non-classical form of 11 β -OHD CAH has been reported. Unlike the common non-classical form of 21OHD, this form is very rare. Non-classical 11 β -OHD has been diagnosed in normotensive children with mild virilization or precocious pubarche [67] and in adults with signs of hyperandrogenemia [68] as well as a woman with infertility [69]. Despite a hormonal profile consistent with 11 β -OHD, mutations in the *CYP11B1* gene may not always be present [68].

3- β hydroxysteroid dehydrogenase deficiency (1) Chapter 3F

There are two forms of the 3 β -hydroxysteroid dehydrogenase enzyme (3 β -HSD): type I and type II. Type II 3 β -HSD enzyme is expressed in the adrenal cortex and gonads and is responsible for conversion of Δ 5 (delta 5) to Δ 4 (delta 4) steroids [1]. This enzyme is essential for the formation of progesterone, which is the precursor for aldosterone, and 17-OHP, which is the precursor for cortisol in the adrenal cortex as well as for androstenedione, testosterone, and estrogen in the adrenal cortex and gonads [70, 71]. Therefore, deficiency of 3 β -HSD in the classic form of 3 β -HSD deficiency CAH results in insufficient cortisol synthesis, salt-wasting in the most severe form, and virilization of external genitalia in females due to androgen effect from the peripheral conversion of circulating Δ 5 precursors to active Δ 4 steroids. Simultaneous type II 3 β -HSD deficiency in the gonads results in incomplete virilization of the external genitalia in males. Thus, genital ambiguity can result in both sexes [72].

17 α -hydroxylase/17,20 lyase deficiency (1) Chapter 3G

Steroid 17 α -hydroxylase/17,20 lyase deficiency accounts for approximately 1% of all CAH cases and affects steroid synthesis in both the adrenals and gonads [73]. Patients have impaired cortisol synthesis, leading to ACTH oversecretion, which increases serum levels of deoxycorticosterone and especially corticosterone, resulting in low renin hypertension, hypokalemia, and metabolic alkalosis. Affected females are born with normal external genitalia. Affected males are also born with under-virilized genitalia due to their deficient gonadal testosterone production. 17 α -Hydroxylase/17,20 lyase deficiency is often recognized at puberty in female patients who fail to develop secondary sex characteristics [74].

Congenital lipid adrenal hyperplasia

Congenital lipid adrenal hyperplasia is an extremely rare and severe form of CAH which is caused by mutations in the steroidogenic acute regulatory protein (StAR). Both the adrenal glands and the gonads exhibit a severe defect in the conversion of cholesterol to pregnenolone [75, 76]. More specifically, StAR mediates the acute steroidogenic response by moving cholesterol from the outer to inner mitochondrial membrane (the rate-limiting step of steroidogenesis), and when this does not occur, cholesterol and cholesterol esters accumulate [77]. In the most severe form, males with congenital lipid hyperplasia are born with female-appearing external genitalia. Females have a normal genital phenotype at birth but remain sexually infantile without treatment. Salt wasting occurs in both males and females. If not detected and treated, the severe form of lipid CAH is usually fatal [78]. More recently, several cases have been reported that demonstrate that lipid CAH has a spectrum of clinical presentation, with varying degrees of genital ambiguity (including normal male genitalia in a 46, XY male) and adrenal insufficiency. Mutations in the StAR protein have been reported that retain partial protein function, leading to variable phenotype [79].

Cytochrome P450 oxidoreductase deficiency (1) Chapter 3H & 3I

Cytochrome P450 oxidoreductase deficiency is another rare form of CAH that is caused by a mutation on 7q11.2 [80, 81, 82]. It is reviewed in Chapter 3H and 3I in *Genetic Steroid Disorders* [1].

GENETICS

In general, all forms of CAH are transmitted in autosomal recessive mode of inheritance as a single gene disorder. However, there have

been reports of cases where none or only one mutation in the responsible gene was identified, including in cases of 21 OHD CAH [83, 84], StAR protein deficiency [85] and POR deficiency [80]. The genes responsible for each form of CAH are shown in **Table 1**.

21-hydroxylase deficiency

The gene encoding the enzyme 21-hydroxylase, *CYP21A2*, is a microsomal cytochrome P450 located on the short arm of chromosome 6 [86] in the human lymphocyte antigen (HLA) complex [87]. *CYP21A2* and its homologue, the pseudogene *CYP21P*, alternate with two genes called C4B and C4A [87, 88] that encode the two isoforms of the fourth component (C4) of serum complement [89]. *CYP21A2* and *CYP21P*, which each contain 10 exons, share 98% sequence homology in exons and approximately 96% sequence homology in introns [90, 91].

More than 140 mutations have been described including point mutations, small deletions, small insertions and complex rearrangements of the gene [92]. The most common mutations appear to be the result of either of two types of meiotic recombination events between *CYP21* and *CYP21P*: 1) misalignment and unequal crossing over, resulting in large-scale DNA deletions, and 2) apparent gene conversion events that result in the transfer to *CYP21A2* of smaller-scale deleterious mutations present in the *CYP21P* pseudogene [93].

Both classical and non-classical 21-hydroxylase deficiency are inherited in a recessive manner as allelic variants. Classical 21-hydroxylase deficiency tends to result from the presence of two severely affected alleles and non-classical 21-hydroxylase deficiency tends to result from the presence of either two mild 21-hydroxylase deficiency alleles or one severe and one mild allele (compound heterozygote). It is important to note, however, that the 10 most common mutations observed in *CYP21A2* cause variable phenotype effects and are not always concordant with genotype. A study by Finkelstein et al demonstrated that the genotype-phenotype concordance was as high as 90.5% for salt-wasting CAH, 85.1% for simple-virilizing CAH, and 97.8% for non-classical CAH [95]. In a recent study of 1,507 subjects with CAH by New et al, a direct genotype-phenotype correlation was noted in less than 50% of the genotypes studied. In the salt wasting and non-classical forms of 21OHD CAH, a phenotype was strongly correlated to a genotype [96]. Rocha et al in 2008 showed that CAG repeats in the androgen receptor has a great influence on variability in virilization of external genitalia of CAH women [97].




Figure 3

Common mutations in CYP21A2 gene and their related phenotypes. The numbers indicated exons of the gene.xcviiiFrom: Forest MG. Recent advances in the diagnosis and management of congenital adrenal hyperplasia due to 21-hydroxylase deficiency.Hum Reprod Update.2004 Nov-Dec;10(6):469-85. Permission obtained

DIAGNOSIS

Hormonal diagnosis

Potential diagnosis of CAH must be suspected in infants born with ambiguous genitalia. The physician is obliged to make the diagnosis as quickly as possible to initiate therapy. The diagnosis and rational decision of sex assignment must rely on the determination of genetic sex, the hormonal determination of the specific deficient enzyme, and an assessment of the patient's potential for future sexual activity and fertility. Physicians are urged to recognize the physical characteristics of CAH in newborns (e.g. ambiguous genitalia) and to refer such cases to appropriate clinics for full endocrine evaluation. As indicated in Table 1, each form of CAH has its own unique hormonal profile, consisting of elevated levels of precursors and elevated or diminished levels of adrenal steroid products. Traditionally, laboratories measured urinary excretion of adrenal hormones or their urinary metabolites (e.g. 17-ketosteroids). However, collection of 24 hour urine excretion is difficult, particularly in neonates. (99) Therefore, simple and reliable radioimmunoassays are utilized now for measuring circulating serum levels of adrenal steroids [100]. Alternatively, a non-invasive random urine collection in the first days of life for steroid hormone metabolites and precursor/product ratio assessments can be measured simultaneously. It can be used independently or in conjunction with serum steroid assays to increase accuracy and confidence in making the diagnosis and distinguishing the separate enzymatic forms of the disorder [101, 102].

Diagnosis of the 21OHD CAH can also be confirmed biochemically by a hormonal evaluation. In a randomly timed blood sample, a very high concentration of 17-hydroxyprogesterone (17-OHP), the precursor of the defective enzyme, is diagnostic of classical 21OHD. Such testing is the basis of the newborn-screening program developed to identify classically affected patients who are at risk for salt wasting crisis [103]. Only 20µl blood, obtained by heel prick and blotted on microfilter paper, is used for this purpose to provide a reliable diagnostic measurement of 17-OHP. The simplicity of the test and the ease of transporting microfilter paper specimens by mail have facilitated the implementation of CAH newborn screening programs worldwide. As of 2009, all 50 states in the United States and 12 other countries screen for CAH. False-positive results are, however, common with premature infants [104]. Appropriate references based on weight and gestational age are therefore in place in many screening programs [105]. The majority of screening programs use a single screening test without retesting of questionable 17-OHP concentrations. To improve efficacy, a small number of programs perform a second screening test of the initial sample to re-evaluate borderline cases identified by the first screening. Current immunoassay methods used in newborn screening programs yield a high false positive rate. In order to decrease this high rate, liquid-chromatography- tandem mass spectrometry measuring different hormones (17-OHP, Δ4-androstenedione, and cortisol) has been suggested as a second-tier method of analyzing positive results [106].

The gold standard for hormonal diagnosis is the corticotropin stimulation test (250 µg cosyntropin intravenously), measuring levels of 17-OHP and Δ4-androstenedione at baseline and 60 min. These values can then be plotted in the published nomogram (Figure 4) to ascertain disease severity [107]. ~~It is important to note that the corticotropin stimulation test should not be performed during the initial 24 hours of life as samples from this period are typically elevated in all infants and may yield false positive results.~~ The corticotropin stimulation test is crucial in establishing hormonal diagnosis of non-classical form of the disease since early-morning values of 17-OHP may not be sufficiently elevated to allow accurate diagnosis.

17OHP NOMOGRAM FOR THE DIAGNOSIS OF STEROID 21 – HYDROXYLASE DEFICIENCY 60 MINUTE CORTROSYN STIMULATION TEST




Figure 4

Nomogram relating baseline to ACTH-stimulated serum concentrations of 17-hydroxyprogesterone (17-OHP). The scales are logarithmic. A regression line for all data points is shown. Data points cluster as shown into three nonoverlapping groups: classic and non-classic forms of 21-hydroxylase deficiency are readily distinguished from each other and from those that are heterozygotes and unaffected. Distinguishing unaffected from heterozygotes is difficult.¹⁰⁷ Adapted from: New MI, Lorenzen F, Lerner AJ, et al. 1983 Genotyping steroid 21-hydroxylase deficiency: hormonal reference data. *J Clin Endocrinol Metab* 57:320-6. Permission obtained.

Prenatal Diagnosis and Prenatal Treatment

Prenatal Diagnosis of 21OHD

A number of approaches to prenatal identification of affected fetuses have been used. In 1965, Jeffcoate et al first reported a successful prenatal diagnosis of 21OHD, based on elevated levels of 17-ketosteroids and pregnanetriol in the amniotic fluid [108]. The hormonal diagnostic test for 21OHD is amniotic fluid 17-OHP. Hormonal diagnosis is currently only used when molecular diagnosis is unavailable.

Advances in genotyping of the *CYP21A2* gene have made molecular genetic studies of extracted fetal DNA the ideal method to diagnose 21OHD CAH in the fetus. Approximately 95% to 98% of the mutations causing 21OHD have been identified through a combination of molecular genetic techniques to study large gene rearrangement and arrays of point mutations [109, 110]. Of the currently available methods for prenatal diagnosis of CAH, CVS, rather than amniocentesis, with molecular genotyping is the preferred diagnostic method in use. Chorionic villus sampling is performed between the 9th and 11th week of gestation, while amniocentesis is usually performed in the second trimester. The timing of prenatal diagnosis is particularly important when deciding to treat the fetus at risk for CAH with dexamethasone prenatally to prevent virilization of the genitalia (see Prenatal Treatment below). As we only wish to treat affected females until term and only 1/8 of the fetuses will be affected and 1/2 will be males, 7 out of 8 fetuses do not require treatment. Thus, amniocentesis, which is performed later in gestation, results in treatment of unaffected fetuses for a longer period of time than CVS. However, amniocentesis can be used as a reliable alternative method of prenatal diagnosis when CVS is unavailable. In such instances, the supernatant is used for hormonal measurement and the cells are cultured to obtain a genotype through DNA analysis. The supernatant hormone measurements distinguish affected status from non affected status only in SW patients. Nonetheless, pitfalls do occur in a small percentage of the patients undergoing prenatal diagnosis utilizing genetic diagnosis, such as undetectable mutations [111], allele drop outs [112], or maternal DNA contamination. Determination of satellite markers may increase the accuracy of molecular genetic analysis [113].

Non-invasive prenatal diagnosis of CAH

Virilization of the genitalia in a female fetus affected with CAH owing to 21OHD and 11B-OHD can be treated prenatally with dexamethasone administered to the mother (see Prenatal Treatment below). Because CAH is an autosomal recessive disorder, the risk is 1/4 of the fetus being affected with the disease and 1/8 of the fetus being a female with ambiguous genitalia. Therefore, 7 out of 8 pregnancies will receive unnecessary treatment until the sex and the affection status of the fetus are known. Treatment with dexamethasone must begin before the 9th week of gestation, yet chorionic villous sampling can only be done at the 9-11th week, with karyotype and DNA results available 2-3 weeks later. Non-invasive prenatal diagnosis would eliminate unnecessary treatment and invasive procedures such as CVS and amniocentesis. Dennis Lo et al. in 1997 discovered the presence of fetal DNA in the maternal circulation [114]. Fetal DNA has been extracted and enriched with high accuracy and yield in fetal Rh factor identification [115], and monogenic disorders such as thalassemia and cystic fibrosis [116, 117]. Identification of the SRY sequence in maternal blood, performed in multiple academic centers and recently in commercial laboratories, has also achieved excellent accuracy in several studies [118, 119]. In non-invasive prenatal diagnosis of CAH, by extracting fetal DNA from the maternal blood at 6-7 weeks gestation, the SRY sequence can be identified, and if the fetus is female (SRY sequence not identified), DNA analysis on extracted fetal DNA is to be done. Thus far, there has been only one case reported of prenatal diagnosis in CAH [120]. However, studies are underway to further develop this innovative method.

Preimplantation diagnosis

Preimplantation genetic diagnosis (PGD) identifies genetic abnormalities in preimplantation embryos prior to embryo transfer, so only unaffected embryos established from IVF are transferred. The procedure has been utilized in many monogenic recessive disorders such as cystic fibrosis, hemoglobinopathies, spinal muscular atrophy and Tay Sach's disease. PGD is being used for a growing number of genetic diseases [121]. There is only one report of PGD utilized in a family whose offspring is at risk for CAH [122], however we know from experience that families are seeing PGD with greater frequency. It would be desirable to have further studies of preimplantation diagnosis in CAH families.

Prenatal Treatment

In 21OHD, prenatal treatment with dexamethasone was introduced in France in 1978 [123] and in the United States in 1986 [124].

Institution of therapy before the 9th week of gestation, prior to the onset of adrenal androgen secretion, effectively suppresses excessive adrenal androgen production and prevents virilization of external female genitalia. Dexamethasone is used because it binds minimally to cortisol-binding globulin (CBG) in the maternal blood, and unlike hydrocortisone, it escapes inactivation by the placental 11-dehydrogenase enzyme. Thus, dexamethasone crosses the placenta from the mother to the fetus and suppresses ACTH secretion with longer half life compared to other synthetic steroids [124].

When dexamethasone administration begins as early as the 8th week of gestation, the treatment is blind to the disease status and sex of the fetus. If the fetus is later determined to be a male upon karyotype or an unaffected female upon DNA analysis, treatment is discontinued. Otherwise, treatment is continued to term. A simplified algorithm of management of potentially affected pregnancies is shown in Figure 4. The optimal dosage is 20 µg/kg/day of dexamethasone per maternal pre-pregnancy body weight, in three divided daily doses [125]. It is recommended to start the treatment as soon as pregnancy is confirmed, and no later than 9 weeks after the last menstrual period [126 127]. The mother’s blood pressure, weight, glycosuria, HbA1C, symptoms of edema, striae and other possible adverse effects of dexamethasone treatment should be carefully observed throughout pregnancy [104].




Figure 5

Algorithm of treatment, diagnosis and decision-making for prenatal treatment of fetuses at risk for 21-hydroxylase deficiency congenital adrenal hyperplasia. Mercado AB, Wilson RC, Cheng KC, Wei JQ, New MI 1995 Extensive personal experience: Prenatal treatment and diagnosis of congenital adrenal hyperplasia owing to steroid 21-hydroxylase deficiency. J Clin Endocrinol Metab 80:2014-2020.

Permission obtained.

Outcome of prenatal treatment of 21OHD

Prenatal treatment of 21OHD has proven to be successful in significantly reducing genital masculinization in affected females. Our group has performed prenatal diagnosis in over 685 pregnancies at risk for 21OHD, and 59 affected female fetuses have been treated to term [128]. Treatment was highly effective in preventing genital ambiguity when the mother was compliant until term. In all the female fetuses treated to term, the degree of virilization was on average 1.69, as measured using the Prader scoring system (Figure 2). Late treatment as well as no treatment resulted in much greater virilization, and the average Prader score was 3.73 for those female fetuses not treated. 124 and unpublished data. Not only does prenatal treatment effectively minimize the degree of female genital masculinization in the patients, it also lessens the high-level androgen exposure of the brain during differentiation. The latter is thought to cause a higher tendency to gender ambiguity in some females with CAH [129, 130]. Genital virilization in female newborns with classical 21OHD CAH has potential adverse psychosocial implications that may be alleviated by prenatal treatment [125].

Prenatal dexamethasone treatment has recently been a subject of controversy and heated debate [131]. Some uncertainties and concerns have been expressed about the long-term safety of prenatal diagnosis and treatment [132, 133]. Concerns have been raised in regards to the glucocorticoid effects on the fetal brain, which arise from studies of other conditions rather than direct studies on prenatal treatment of 21OHD CAH. These include studies whereby much higher doses of dexamethasone were given to the human subjects at the later part of pregnancy [134] or to animals [135, 136] and therefore hold little relevance to using dexamethasone prenatally in CAH. In a small-sample study of children prenatally treated with dexamethasone, Lajic and her colleagues found no effects on intelligence, handedness, memory encoding, or long-term memory, but short-term treated CAH-affected children had significantly poorer performance than controls on a test of verbal working memory. These patients also had lower questionnaire scores in self-perceived scholastic competence and social anxiety [137]. However, parents described these children as more sociable than controls, without significant difference in psychopathology, school performance, adaptive functioning or behavioral problems [138]. A larger multi-center study (US and France) indicated no adverse cognitive effects of short-term prenatal DEX exposure, including no adverse influence on verbal working memory; a small sample of dexamethasone treated girls affected with CAH showed lower scores on two of eight neuropsychological tests, however given the variability of cognitive findings in dexamethasone unexposed CAH-affected patients, this result cannot be linked to dexamethasone with certainty. This indicates that further studies are needed [139].

Compelling data from large cohorts of pregnancies with prenatal diagnosis and treatment of 21OHD CAH prove its efficacy and safety [140, 141]. All of the mothers who received prenatal treatment (partial or full-term) stated that they would take dexamethasone again for a future pregnancy [142].) Rare adverse events have been reported in treated children, but no harmful effects have been documented that can be clearly attributed to the treatment [143]. Another long-term follow-up study in Scandinavia showed that 44 children who were variably treated prenatally demonstrated normal prenatal and postnatal growth compared to matched controls. Further, there was no observed increase in fetal abnormalities or fetal death [144]. Although some abnormalities in postnatal growth and behavior were observed among dexamethasone exposed offspring, none could logically be explained by the present knowledge of teratogenic effects of glucocorticoids.

Published studies of almost 600 pregnancies, 80 of which were prenatally treated until term and 27 who were male and received dexamethasone for a short period of time, the newborns in the dexamethasone treated group did not differ in weight, length or head circumference from untreated, unaffected siblings. No significant or enduring side effects were noted in either the mothers or the fetuses. Greater weight gain in treated versus untreated mothers did occur, as well as the presence of striae and edema. Excessive weight gain was lost after birth. No differences were found regarding gestational diabetes or hypertension [147]. No cases have been reported of cleft palate, placental degeneration or fetal death, which have been observed in the rodent model of *in utero* exposure to high-dose glucocorticoids [145]. One explanation for the safety of human versus rodent is that glucocorticoid receptor-ligand systems in human differ from that of rodents [146]. Most recently, a comprehensive long-term outcome study looking at 149 male and female patients 12 years of age and older, affected and unaffected with CAH, who were treated with dexamethasone partially or to term was conducted. To date, this is the largest study evaluating the long-term effects of dexamethasone. No adverse effects such as increased risk for cognitive defects, disorders of gender identity and behavior, sexual function in adulthood, hypertension, diabetes, and osteopenia were found [147].

Prenatal diagnosis and treatment of 11 β -OHD CAH

A number of approaches to prenatal identification by measuring steroid precursors in affected fetuses have been used [148, 149, 150]. Advances in genotyping of the *CYP11B1* gene have made molecular genetic studies of extracted fetal DNA, the ideal method to diagnose 11 β -OHD CAH in the fetus [151, 152]. The established protocol of prenatal diagnosis and treatment in 21OHD CAH can be applied to 11 β -OHD CAH. Successful results in prenatal diagnosis and treatment in 11 β -OHD CAH have been reported [151, 153].

TREATMENT

Hormone Replacement

The goal of therapy in CAH is to both correct the deficiency in cortisol secretion and to suppress ACTH overproduction. Proper treatment with glucocorticoid reduces stimulation of the androgen pathway, thus preventing further virilization and allowing normal growth and development. The usual requirement of hydrocortisone (or its equivalent) for the treatment of classical 21OHD form of CAH is about 10-15 mg/m²/day divided into 2 or 3 doses per day. Dosage requirements for patients with NC-21OHD CAH are typically less. Adults may be treated with the longer-acting dexamethasone or prednisone, alone or in combination with hydrocortisone. A small dose of dexamethasone at bedtime (0.25 to 0.5 mg) is usually adequate for androgen suppression in non-classical patients. Anti-androgen treatment may be useful as adjunctive therapy in adult women who continue to have hyperandrogenic signs despite good adrenal suppression. Females with concomitant PCOS may benefit from an oral contraceptive, though this treatment would not be appropriate for patients trying to get pregnant. Treatment of adult males with NC-21OHD may not be necessary, though our group has found that it may be helpful in preventing adrenal rest tumors and preserving fertility. Optimal corticosteroid therapy is determined by adequate suppression of adrenal hormones balanced against normal physiological parameters. ~~The goal of corticosteroid therapy is to give the lowest dose required for optimal control.~~ Adequate biochemical control is assessed by measuring serum levels 17-OHP and androstenedione; serum testosterone can be used in females and prepubertal males (but not in newborn males). It is recommended that hormone levels are measured at a consistent time in relation to medication dosing, usually 2 hours after the morning corticosteroid. Titration of the dose should be aimed at maintaining androgen levels at age and sex-appropriate levels and 17-OHP levels of <1000 ng/dL. Concurrently, over-treatment should be avoided because it can lead to Cushing syndrome. Depending on the degree of stress, stress dose coverage may require doses of up to 50-100 mg/m²/day [1].

Patients with salt-wasting CAH have elevated plasma renin in response to the sodium-deficient state, and they require treatment with the salt-retaining 9 α -fludrocortisone acetate. The average dose is 0.1 mg daily, ranging from 0.05 mg to 0.2 mg daily. Infants should also be started on salt supplementation, as sodium chloride, at 1-2 g daily, divided into several feedings. Although patients with the SV and NC form of CAH can make adequate aldosterone, the aldosterone to renin ratio (ARR) has been found to be lower than normal, though not to the degree seen in the salt-wasting form [154]. It has not been customary to supplement conventional glucocorticoid replacement therapy with the administration of salt-retaining steroids in the SV and NC forms of CAH, though there has been some suggestion that adding fludrocortisone to patients with elevated PRA may improve hormonal control of the disease [155]. The requirement for fludrocortisone appears to diminish with age, and over-suppression of the PRA should be avoided, to prevent complications from hypertension and excessive mineralocorticoid activity. Measurements of plasma renin and aldosterone are used to monitor the efficacy of mineralocorticoid therapy in all patients with the salt wasting form of the disease [1].

In managing 11 β -OHD, glucocorticoid administration provides cortisol replacement and decreases ACTH, as it does in 21OHD. This in turn removes the drive for oversecretion of deoxycorticosterone (DOC) and, in most cases, normalizes blood pressure. A thorough examination undertaken by endocrine challenge and suppression studies to evaluate zonal differences has shown that in 11 β -OHD CAH, the zona fasciculata exhibits reduced 11 β -hydroxylation and 18-hydroxylation, while both functions appear to be spared in the zona glomerulosa [156]. This demonstrates that the zona glomerulosa and the zona fasciculata function as two physiologically, and likely genetically, separate glands. Glucocorticoid treatment produces natriuresis and diuresis, normalizes plasma volume and thus increases the plasma renin to levels that stimulate aldosterone production in the zona glomerulosa. In addition to normalizing blood pressure, the goal of treatment is to replace deficient steroids and in turn minimize adrenal sex hormone excess, prevent virilization, optimize growth, and protect potential fertility. Serum DOC is thus the principal steroid index of the 11 β -OHD. Plasma renin activity is useful as a therapeutic index as well. In poorly controlled 11 β -OHD patients, DOC is elevated and plasma renin is suppressed; both are normal in well-controlled patients. As in patients with 21OHD, oral hydrocortisone at a dose of 10-15 mg/m² divided into two to three daily doses

is the preferred treatment. Long-acting glucocorticoids may be used at or near the completion of linear growth. In patients who have had ongoing hypertension for some time before diagnosis is made, adding spironolactone, calcium channel blockers or amiloride may be necessary [61].

In patients with 3 β -HSD deficiency, glucocorticoid administration also reduces the excess production of androgens. In addition, these patients have mineralocorticoid deficiency and require treatment with the salt-retaining 9 α -fludrocortisone acetate. Patients with the StAR protein deficiency (lipoid form of CAH) classically have severe adrenal insufficiency with mineralocorticoid deficiency and salt wasting; they require both glucocorticoid and mineralocorticoid replacement. Patients with 17 α -hydroxylase/17,20 lyase deficiency typically have excess DOC and low-renin hypertension, and treatment with a glucocorticoid should normalize serum DOC level and lead to normalization of blood pressure. In several conditions, such as StAR protein deficiency, 3 β -HSD, 17 α -hydroxylase/17,20 lyase deficiency and cytochrome P450 oxidoreductase deficiency, patients require sex steroid replacement. Sex steroids should be added at a developmentally appropriate time to allow patients to resemble their peers.

Because patients with CAH are at risk for short stature as adults, other adjunct therapies are being utilized. Two studies have demonstrated significant improvement in growth velocity, final adult height prediction [20] and final adult height [19] with the use of growth hormone in conjunction with a GnRH analogue.

In non-life-threatening periods of illness or physiologic stress, the corticosteroid dose should be increased to 2 or 3 times the maintenance dose for the duration of that period. Each family should be given injection kits of hydrocortisone for emergency use (25 mg for infants, 50 mg for children, and 100 mg for adults). In the event of a surgical procedure, a total of 5 to 10 times the daily maintenance dose may be required during the first 24–48 hours, which can then be tapered over the following days to the normal preoperative schedule. Stress doses of dexamethasone should not be given because of the delayed onset of action. It is not necessary for increased mineralocorticoid doses during these periods of stress [1].

It is imperative for all patients who are receiving corticosteroid replacement therapy, such as patients with CAH, to wear a Medic-Alert bracelet or medallion that will enable correct and appropriate therapy in case of emergencies. Additionally, all responsible family members should be trained in the intramuscular administration of hydrocortisone.

Bone Mineral Density

In order to adequately suppress androgen production in patients with CAH, the usual requirement of hydrocortisone is generally higher than the endogenous secretory rate of cortisol. Chronic therapy with glucocorticoids at supraphysiologic levels can result in diminished bone accrual and lead to osteopenia and osteoporosis. Glucocorticoid induced bone loss is a well-known phenomenon and is the most prevalent form of secondary osteoporosis [157, 158, 159].

Unlike other diseases treated with chronic glucocorticoid therapy, however, the effect of glucocorticoid replacement in CAH on BMD is unclear. Previous studies of patients with 21OHD have reported increased, normal, or decreased BMD [160, 161, 162, 163]. It has been postulated that the elevated androgens typically found in patients may have a protective effect on bone integrity, but the precise mechanism is unknown. The increased adrenal androgens, which are converted to estrogens, may counteract the detrimental effects of glucocorticoids on bone mass. This may explain why older CAH women, particularly those who are post-menopausal, are at higher risk for osteoporosis than younger CAH patients. It has been proposed that the inhibitory effect of corticosteroid therapy on bone formation is counteracted by estrogen's effect on bone resorption through the RANK-L/osteoprotegerin (OPG) system [164]. A study is currently underway to examine a potential mechanism by which certain CAH patients are protected against glucocorticoid induced bone loss.

Surgery

~~The aim of surgical repair in females with ambiguous genitalia caused by CAH, when the decision is made by parents or patients themselves, is generally to remove the redundant erectile tissue, preserve the sexually sensitive glans clitoris, and provide a normal vaginal orifice that functions adequately for menstruation, intromission, and delivery.~~ A medical indication for early surgery other than for sex assignment is recurrent urinary tract infections as a result of pooling of urine in the vagina or urogenital sinus. In the past, it was routine to recommend early corrective surgery for neonates born with ambiguous genitalia. However, in recent years, the implementation of early corrective surgery has become increasingly controversial due to lack of data on long-term functional outcome.

Recent data shows that genital sensitivity is impaired in areas where feminizing genital surgery had been done, leading to difficulties in sexual function [165]. Another recent study showed that patients with more severe mutations in the *CYP21A2* gene, i.e. those with the null genotype and thus those more severely virilized, had more surgical complications than those less severely virilized and were less satisfied with their sexual life [166]. Because of the scarcity of this data, the role of the parents in sex assignment becomes crucial in all aspects of the decision making process, and should include full discussion of the controversy and all possible therapeutic options for the intersex child, particularly early versus delayed surgery. Further, a multidisciplinary case-by-case approach, involving pediatric endocrinology, urology, genetics, and psychoendocrinology, is imperative when considering sex assignment and possible surgical repair.

Other Treatment Strategies

Glucocorticoid replacement has been an effective treatment for CAH for over 50 years and remains its primary therapy; however, the management of these patients presents a challenge because inadequate treatment as well as oversuppression can both cause complications.

Bilateral adrenalectomy is a radical measure that can be effective in some cases. A few patients who were extremely difficult to control with medical therapy alone showed improvement in their symptoms after bilateral adrenalectomy [167, 168]. Because this approach renders the patient completely adrenal insufficient, however, it should be reserved for extreme cases and is not a good treatment option for patients who have a history of poor compliance with medication.

CONCLUSION

The pathophysiology of the various types of CAH (the most common being 21OHD) can be traced to specific, inherited defects in the genes encoding enzymes for adrenal steroidogenesis. Clinical presentation of each form is distinctive and depends largely on the underlying enzyme defect, its precursor retention and deficient products. Treatment of CAH is targeted to replace the insufficient adrenal hormones and to suppress the excess precursors. With proper hormone replacement therapy, normal and healthy development may be expected. Glucocorticoid and, if necessary, mineralocorticoid replacement, has been the mainstay of treatment for CAH, but new treatment strategies continue to be developed and studied to improve care. The advent of genetic testing in the fetus and the affected living patient has made the prenatal diagnosis of CAH more accurate and secure.

References

1. Genetic Steroid Disorders. New MI, Lekarev O, Parsa A, Yuen T, O'Malley B, Hammer G eds. 2013. Elsevier, San Diego, CA.
2. Pang SY, Wallace MA, Hofman L, et al. 1988 Worldwide experience in newborn screening for classical congenital adrenal hyperplasia due to 21-hydroxylase deficiency. *Pediatrics* 81:866-74. [PubMed: 3259306]
3. NNSIS 2009 National Newborn Screening Information System. Available at <http://www2.uthscsa.edu/nnsis>.
4. Speiser PW, Dupont B, Rubinstein P, Piazza A, Kastelan A, New MI 1985 High frequency of nonclassical steroid 21-hydroxylase deficiency. *Am J Hum Genet* 37:650-67. [PMC free article: PMC1684620] [PubMed: 9556656]
5. Sherman SL, Aston CE, Morton NE, Speiser P, New MI 1988 A segregation and linkage study of classical and nonclassical 21-hydroxylase deficiency. *Am J Hum Genet* 42:830-838. [PMC free article: PMC1715213] [PubMed: 3259403]
6. Zerah M, Ueshiba H, Wood E, et al. 1990 Prevalence of nonclassical steroid 21-hydroxylase deficiency based on a morning salivary 17-hydroxyprogesterone screening test: a small sample study. *Journal of Clinical Endocrinology & Metabolism* 70:1662-7. [PubMed: 2347899]
7. Zachmann M, Tassinari D, Prader A 1983 Clinical and biochemical variability of congenital adrenal hyperplasia due to 11 beta-hydroxylase deficiency. *J Endocrinol Metab* 56:222-229. [PubMed: 6296182]
8. Curnow KM, Slutsker L, Vitek J, et al. 1993 Mutations in the *CYP11B1* gene causing congenital adrenal hyperplasia and hypertension cluster in exons 6, 7, and 8. *Proc Natl Acad Sci (USA)* 90:4552. [PMC free article: PMC46550] [PubMed: 8506298]
9. Khemiri M, Ridane H, Bou YO, Matoussi N, Khaldi F 2006. 11 beta hydroxylase deficiency: a clinical study of seven cases. *Tunis Med* 84:106-13. [PubMed: 16755975]
10. Rosler A, Leiberman E, Cohen T 1992 High frequency of congenital adrenal hyperplasia (classic 11 beta-hydroxylase deficiency) among Jews from Morocco. *Am J Med Genet* 42:827-34. [PubMed: 1554023]

11. Paperna T, Gershoni-Baruch R, Badarneh K, Kasinetz L, Hochberg Z 2005 Mutations in CYP11B1 and congenital adrenal hyperplasia in Moroccan Jews. *J Clin Endocrinol Metab* 90:5463-5. [PubMed: 16030166]
12. Ganong W 1963 The central nervous system and the synthesis and release of adrenocorticotrophic hormone. In: Nalbandov A (ed) *Advances in Neuroendocrinology*. University of Illinois Press, Urbana, p 92.
13. Guillemin R, Schally A 1963 Recent advances in the chemistry of neuroendocrine mediators originating in the central nervous system. In: Nalbandov A (ed) *Advances in Neuroendocrinology*. University of Illinois Press, Urbana, p 314.
14. Goto M, Piper Hanley K, Marcos J, et al. 2006 In humans, early cortisol biosynthesis provides a mechanism to safeguard female sexual development. *J Clin Invest* 116:953-60. [PMC free article: PMC1421344] [PubMed: 16585961]
15. Prader A, Gurtner HP 1955 The syndrome of male pseudohermaphroditism in congenital adrenocortical hyperplasia without overproduction of androgens (adrenal male pseudohermaphroditism). *Helv Paediatr Acta* 10:397-412. [PubMed: 13285832]
16. DiMartino-Nardi J, Stoner E, O'Connell A, New MI 1986 The effect of treatment of final height in classical congenital adrenal hyperplasia (CAH). *Acta Endocrinol Suppl (Copenh)*. 279:305-14. [PubMed: 3465172]
17. New MI, Gertner JM, Speiser PW et al. 1989 Growth and final height in classical and nonclassical 21-hydroxylase deficiency. *J Endocrinol Invest*. 12(8 Suppl 3):91-5. [PubMed: 2809100]
18. Lin-Su K, Harbison MD, Lekarev O, Vogiatzi MG, New MI 2011 Final adult height in children with congenital adrenal hyperplasia treated with growth hormone. *J Clin Endocrinol Metab* 96(6): 170-7. [PMC free article: PMC3206397] [PubMed: 21450983]
19. Trinh L, Nimkarn S, Obeid J, New M, Su K Growth and Pubertal Characteristics in Patients with Congenital Adrenal Hyperplasia Due to 21-Hydroxylase Deficiency. *J Pediatr Endocrinol Metab*. 2007 Aug;20(8):883-91. [PubMed: 17937061]
20. Klingensmith G, Garcia S, Jones H, Migeon C, Blizzard R 1977 Glucocorticoid treatment of girls with congenital adrenal hyperplasia: effects on height, sexual maturation, and fertility. *J Pediatr* 90:996-1004. [PubMed: 870661]
21. Helleday J, Siwers B, Ritzen EM, Carlstrom K. Subnormal androgen and elevated progesterone levels in women treated for congenital virilizing 21-hydroxylase deficiency. *J Clin Endocrinol Metab* 1993;76(4):933-6. [PubMed: 8473408]
22. Premawardhana L, Hughes I, Read G, Scanlon M. Longer term outcome in females with congenital adrenal hyperplasia (CAH): the Cardiff experience. *Clin Endocrinol (Oxf)* 1997;46(3):327-32. [PubMed: 9156043]
23. Richards G, Grumbach M, Kaplan S, Conte F. The effect of long acting glucocorticoids on menstrual abnormalities in patients with virilizing congenital adrenal hyperplasia. *J Clin Endocrinol Metab* 1978;47(6): pp 1208-15. [PubMed: 233693]
24. Barnes R, Rosenfield R. and Ehrmann D. et al., Ovarian hyperandrogenism as a result of congenital adrenal virilizing disorders: evidence for perinatal masculinization of neuroendocrine function in women, *J Clin Endocrinol Metab* 79 (1994), pp. 1328-1333. [PubMed: 7962325]
25. Long D., Wisniewski A. and Migeon C. Gender role across development in adult women with congenital adrenal hyperplasia due to 21-hydroxylase deficiency, *J Pediatr Endocrinol Metab* 17 (2004), pp. 1367-1373. [PubMed: 15526714]
26. Hines M., Brook C and Conway G. Androgen and psychosexual development: core gender identity, sexual orientation and recalled childhood gender role behavior in women and men with congenital adrenal hyperplasia (CAH), *J Sex Res* 41 (2004), pp. 75-81. [PubMed: 15216426]
27. Meyer-Bahlburg H., Dolezal C. and Baker S. et al., Sexual orientation in women with classical or non-classical congenital adrenal hyperplasia as a function of degree of prenatal androgen excess, *Arch Sex Behav* 37 (2008), pp. 85-99. [PubMed: 18157628]
28. Berenbaum SA, Bryk KL, Beltz AM 2012 Early androgen effects on spatial and mechanical abilities: evidence from congenital adrenal hyperplasia. *Behav Neurosci* 126(1):86-96. [PMC free article: PMC3270350] [PubMed: 22289044]
29. Berenbaum SA, Bryk KK, Duck SC 2010 Normal intelligence in female and male patients with congenital adrenal hyperplasia. *Int J Pediatr Endocrinol* 2010:853103. [PMC free article: PMC2952913] [PubMed: 20976263]
30. Mulaikal RM, Migeon CJ, Rock JA. Fertility rates in female patients with congenital adrenal hyperplasia due to 21-hydroxylase deficiency. *NEJM* 1987;316:178-82. [PubMed: 3491959]
31. Cabrera M, Vogiatzi M, New M 2001 Long term outcome in adult males with classic congenital adrenal hyperplasia. *J Clin Endocrinol Metab* 86:3070-3080. [PubMed: 11443169]
32. Bonaccorsi A, Adler I, Figueiredo J 1987 Male infertility due to congenital adrenal hyperplasia: testicular biopsy findings, hormonal evaluation, and therapeutic results in three patients. *Fertil Steril* 47: pp 664-70. [PubMed: 3032693]
33. Prader A: Vollkommen mannliche aussere Genitalentwicklung und Salzerlutsyndrom bei Madech mit Kongenitalem

- Androgenitalem Syndrom. *Helv Paediatr Act* 13:5-14, 1958. [PubMed: 13524803]
34. Urban M, Lee P, Migeon C 1978 Adult height and fertility in men with congenital virilizing adrenal hyperplasia. *New Engl J Med* 299:1392. [PubMed: 152409]
 35. Claahsen-van der Grinten HL, Sweep FC, Blickman JG, Hermus AR, Otten BJ 2007 Prevalence of testicular adrenal rest tumours in male children with congenital adrenal hyperplasia due to 21-hydroxylase deficiency. *Eur J Endocrinol* 157:339-44. [PubMed: 17766717]
 36. Molitor J, Chertow B, Fariss B 1973 Long-term follow-up of a patient with congenital adrenal hyperplasia and failure of testicular development. *Fertil Steril* 24:319. [PubMed: 4694507]
 37. Claahsen-van der Grinten HL, Otten BJ, Hermus AR, Sweep FC, Hulsbergen-van de Kaa CA 2008 Testicular adrenal rest tumors in patients with congenital adrenal hyperplasia can cause severe testicular damage. *Fertil Steril* 89:597-601. [PubMed: 17543962]
 38. Wilkins L, Crigler J, Silverman S, Gardner L, Migeon C 1952 Further studies on the treatment of congenital adrenal hyperplasia with cortisone. II. The effects of cortisone on sexual and somatic development, with an hypothesis concerning the mechanism of feminization. *J Clin Endocrinol Metab* 12:277. [PubMed: 14907858]
 39. Blumberg-Tick J, Boudou P, Nahoul K, Schaison G 1991 Testicular tumors in congenital adrenal hyperplasia: steroid measurements from adrenal and spermatic veins. *J Clin Endocrinol Metab* 73:1129-33. [PubMed: 1939529]
 40. Schoen E, DiRaimondo V, Dominguez O 1961 Bilateral testicular tumors complicating congenital adrenocortical hyperplasia. *J Clin Endocrinol Metab* 21:518. [PubMed: 13748349]
 41. Miller E, Murray H 1962 Congenital adrenocortical hyperplasia: case previously reported as “bilateral interstitial cell tumor of the testicle. *J Clin Endocrinol Metab* 22:655. [PubMed: 14473989]
 42. Glenn J, Boyce W 1963 Adrenogenitalism with testicular adrenal rests simulating interstitial cell tumor. *J Urol* 89:456. [PubMed: 13948452]
 43. Srikanth M, West B, Ishitani M, Isaacs HJ, Applebaum H, Costin G 1992 Benign testicular tumors in children with congenital adrenal hyperplasia. *J Pediatr Surg* 27:639-41. [PubMed: 1320675]
 44. Rutgers J, Young R, Scully R 1988 The testicular “tumor” of the adrenogenital syndrome. A report of six cases and review of the literature on testicular masses in patients with adrenocortical disorders. *Am J Surg Pathol* 12:503-13. [PubMed: 3291624]
 45. Sugino Y, Usui T, Okubo K, et al. 2006 Genotyping of congenital adrenal hyperplasia due to 21-hydroxylase deficiency presenting as male infertility: case report and literature review. *J Assist Reprod Genet* 23:377-80. [PMC free article: PMC3455103] [PubMed: 17033937]
 46. Klein R 1960 Evidence for and evidence against the existence of a salt-losing hormone. *J Pediatr* 57:452. [PubMed: 14409705]
 47. Kowarski A, Finkelstein JW, Spaulding JS, Holman GH, Migeon CJ 1965 Aldosterone Secretion Rate in Congenital Adrenal Hyperplasia. A Discussion of the Theories on the Pathogenesis of the Salt-Losing Form of the Syndrome. *J Clin Invest* 44:1505-13. [PMC free article: PMC292632] [PubMed: 14332163]
 48. Kuhnle U, Land M, Ulick S 1986 Evidence for the secretion of an antiminerlocorticoid in congenital adrenal hyperplasia. *J Clin Endocrinol Metab* 62:934-40. [PubMed: 3007560]
 49. Tusie-Luna M, Traktman P, White PC: Determination of functional effects of mutations in the steroid 21-hydroxylase gene (CYP21) using recombinant vaccinia virus. *J Bio Chem* 265:20916-20922, 1990. [PubMed: 2249999]
 50. Stoner E, Dimartino-Nardi J, Kuhnle U, Levine LS, Oberfield SE, New MI 1986 Is salt-wasting in congenital adrenal hyperplasia due to the same gene as the fasciculata defect? *Clin Endocrinol (Oxf)* 24:9-20. [PubMed: 3486728]
 51. Luetscher JA 1956 Studies of aldosterone in relation to water and electrolyte balance in man. *Recent Prog Horm Res* 12:175. [PubMed: 13401056]
 52. Speiser PW, Agdere L, Ueshiba H, White PC, New MI 1991 Aldosterone synthesis in salt-wasting congenital adrenal hyperplasia with complete absence of adrenal 21-hydroxylase. *New England Journal of Medicine* 324:145-9. [PubMed: 1984191]
 53. Rose L, Newmark S, Strauss J, Pochi P. Adrenocortical hydroxylase deficiencies in acne vulgaris. *J Invest Dermatol* 1976;66(5): pp 324-6. [PubMed: 178807]
 54. Lucky A, Rosenfield R, McGuire J, Rudy S, Helke J. Adrenal androgen hyperresponsiveness to adrenocorticotropin in women with acne and/or hirsutism: adrenal enzyme defects and exaggerated adrenarche. *J Clin Endocrinol Metab* 1986;62(5): pp 840-8. [PubMed: 3007557]
 55. Lin-Su, K, Nimkarn S, New MI. Congenital adrenal hyperplasia in adolescence: diagnosis and management. *Ann NY Acad Sci*.

- 2008;1135:95-8. [PubMed: 18574213]
56. Witchel SF and Azziz R. Nonclassic Congenital Adrenal Hyperplasia. *International Journal of Pediatric Endocrinology*. Volume 2010. pp 1-11.
 57. Mokshagundam S, Surks M 1993 Congenital adrenal hyperplasia diagnosed in a man during workup for bilateral adrenal masses. *Arch Intern Med* 153:1389-91. [PubMed: 8507129]
 58. Jaresch S, Kornely E, Kley H, Schlaghecke R 1992 Adrenal incidentaloma and patients with homozygous or heterozygous congenital adrenal hyperplasia. *J Clin Endocrinol Metab* 74:685-9. [PubMed: 1311000]
 59. Barzon L, Scaroni C, Sonino N et al. Incidentally discovered adrenal tumors: endocrine and scintographic correlates. 1998 *J Clin Endocrinol Metab* 74:685-689. [PubMed: 9435416]
 60. Nieman L. Approach to the patient with an adrenal incidentaloma. *J Clin Endocrinol Metab*, September 2010, 95(9):4106-4113. [PMC free article: PMC2936073] [PubMed: 20823463]
 61. Nimkarn S, New MI 2008 Steroid 11beta- hydroxylase deficiency congenital adrenal hyperplasia. *Trends Endocrinol Metab* 19:96-9. [PubMed: 18294861]
 62. Mimouni M, Kaufman H, Roitman A, Morag C, Sadan N 1985 Hypertension in a neonate with 11 beta-hydroxylase deficiency. *Eur J Pediatr* 143:231-3. [PubMed: 3872797]
 63. Globerman H, Rosler A, Theodor R et al. An inherited defect in aldosterone biosynthesis caused by a mutation in or near the gene for steroid 11-hydroxylase. *N Engl J Med* 319:1193-1197, 1988. [PubMed: 3262827]
 64. Rosler A, Leiberman E, Sack J 1982 Clinical variability of congenital adrenal hyperplasia due to 11B-hydroxylase deficiency. *Hormone Research* 16:133. [PubMed: 7049883]
 65. Hague WM, Honour JW 1983 Malignant hypertension in congenital adrenal hyperplasia due to 11 beta-hydroxylase deficiency. *Clin Endocrinol (Oxf)* 18:505-10. [PubMed: 6603291]
 66. Chabre O, Portrat-Doyen S, Chaffanjon P, et al. 2000 Bilateral laparoscopic adrenalectomy for congenital adrenal hyperplasia with severe hypertension, resulting from two novel mutations in splice donor sites of CYP11B1. *J Clin Endocrinol Metab* 85:4060-8. [PubMed: 11095433]
 67. Zachmann M, Tassinari D, Prader A. Clinical and biochemical variability of congenital adrenal hyperplasia due to 11beta-hydroxylase deficiency, *J. Clin. Endocrinol. Metab.* 56 (1983), pp. 222-229. [PubMed: 6296182]
 68. Joehrer K, Geley S, Strasser-Wozac E et al. CYP11B1 Mutations Causing Non-Classical Adrenal Hyperplasia due to 11β-Hydroxylase Deficiency. *Hum. Mol. Genet.* (1997) 6 (11): 1829-1834. [PubMed: 9302260]
 69. Peters CJ, Tugent T, Perry LA et al. Cosegregation of a Novel Homozygous CYP11B1 Mutation with the Phenotype of Non-Classical Congenital Adrenal Hyperplasia in a Consanguineous Family, *Horm. Res.* 67 (2007), pp. 189-193. [PubMed: 17124386]
 70. Simard J, Rheaume E, Mebarki F et al. Molecular basis of human 3 beta-hydroxysteroid dehydrogenase deficiency. *J Steroid Biochem Mol Biol.* 1995 Jun;53(1-6):127-38. [PubMed: 7626445]
 71. Labrie F, Simard J, Luu-The V, et al. 1992 Structure and tissue-specific expression of 3 beta-hydroxysteroid dehydrogenase/5-ene-4-ene isomerase genes in human and rat classical and peripheral steroidogenic tissues. *J Steroid Biochem Mol Biol* 41:421-35. [PubMed: 1562516]
 72. Pang S. Congenital adrenal hyperplasia owing to 3 beta-hydroxysteroid dehydrogenase deficiency. *Endocrinol Metab Clin North Am.* 2001 Mar;30(1):81-99, vi-vii. [PubMed: 11344940]
 73. Yanase T, Simpson E, Waterman M 1991 17 alpha-hydroxylase/17,20-lyase deficiency: from clinical investigation to molecular definition. *Endocr Rev* 12:91-108. [PubMed: 2026124]
 74. Miller W, Auchus R and Geller D. The regulation of 17, 20 lyase activity. *Steroids*, Volume 62, Issue 1, January 1997, Pages 133-142. [PubMed: 9029728]
 75. Bose H, Sugawara T, Strauss Jr, Miller W 1996 The pathophysiology and genetics of congenital lipid adrenal hyperplasia. *International Congenital Lipoid Adrenal Hyperplasia Consortium*. *N Engl J Med* 335: 1870-8. [PubMed: 8948562]
 76. Bose HS, Sato S, Aisenberg J, Shalev SA, Matsuo N, Miller WL. Mutations in the Steroidogenic Acute Regulatory Protein (StAR) in Six Patients with Congenital Lipoid Adrenal Hyperplasia. *J Clin Endocrinol Metab.* 2000 Oct;85(10):3636-9. [PubMed: 11061515]
 77. Miller WL. Mechanism of StARs regulation of mitochondrial cholesterol import. *Molecular and Cellular Endocrinology* 265-266 (2007), pp 46-50. [PubMed: 17207924]

78. Stocco D, Clark B 1997 The role of the steroidogenic acute regulatory protein in steroidogenesis. *Steroids* 62:29-36. [PubMed: 9029711]
79. Sahakitrungruang T, Soccio R, Lang-Muritano M, et al. Clinical, genetic and functional characterization of four patients carrying partial loss-of-function mutations in the steroidogenic acute regulatory protein (StAR). *J Clin Endocrinol Metab*, July 2010, 95(7), pp 1-7. [PMC free article: PMC2928910] [PubMed: 20444910]
80. Scott RR, Miller WL. Genetic and clinical features of P450 oxidoreductase deficiency. *Hor Res*. 2008;69(5). Pp. 266-75. [PubMed: 18259105]
81. Fluck CE, Pandey AV, Huang N et al. P450 oxidoreductase deficiency - a new form of congenital adrenal hyperplasia. *Endocr Dev*. 2008; 13:67-81. Review. [PubMed: 18493134]
82. Fluck CE, Tajima T, Pandey AV, et al. 2004 Mutant P450 oxidoreductase causes disordered steroidogenesis with and without Antley-Bixler syndrome. *Nat Genet* 36:228-30. [PubMed: 14758361]
83. Wilson RC, Mercado AB, Cheng KC, New MI 1995 Steroid 21-hydroxylase deficiency: genotype may not predict phenotype. *J Clin Endocrinol Metab* 80:2322-2329. [PubMed: 7629224]
84. Nimkarn S, Cerame BI, Wei JQ, et al. 1999 Congenital adrenal hyperplasia (21-hydroxylase deficiency) without demonstrable genetic mutations. *J Clin Endocrinol Metab* 84:378-81. [PubMed: 9920112]
85. Tajima T, Fujieda K, Kouda N, Nakae J, Miller W 2001 Heterozygous mutation in the cholesterol side chain cleavage enzyme (p450scc) gene in a patient with 46,XY sex reversal and adrenal insufficiency. *J Clin Endocrinol Metab* 86: 3820-3825. [PubMed: 11502818]
86. Nebert DW, Nelson DR, Coon MJ. The P450 superfamily: update on new sequences, gene mapping, and recommended nomenclature. *DNA Cell Biol* 1991;10:1-14. [PubMed: 1991046]
87. Dupont B, Oberfield SE, Smithwick EM, Lee TD, Levine LS. Close genetic linkage between HLA and congenital adrenal hyperplasia (21-hydroxylase deficiency). *Lancet* 1977;2(8052-8053):1309-12. [PubMed: 74726]
88. White PC, Grossberger D, Onufer BJ, et al. Two genes encoding steroid 21-hydroxylase are located near the genes encoding the fourth component of complement in man. *Proc Natl Acad Sci U S A* 1985;82(4):1089-93. [PMC free article: PMC397199] [PubMed: 2983330]
89. Carroll MC, Campbell RD, Porter RR. The mapping of 21-hydroxylase genes adjacent to complement component C4 genes in HLA, the major histocompatibility complex in man. *Proc Natl Acad Sci USA* 1985;82:521-5. [PMC free article: PMC397071] [PubMed: 3871526]
90. Belt KT, Carroll MC, Porter RR. The structural basis of the multiple forms of human complement component C4. *Cell* 1984;36(4):907-14. [PubMed: 6546707]
91. White PC, New MI, Dupont B. Structure of the human steroid 21-hydroxylase genes. *Proc Natl Acad Sci U S A* 1986;83(14):5111-5. [PMC free article: PMC323900] [PubMed: 3487786]
92. Higashi Y, Yoshioka H, Yamane M, Gotoh O, Fujii-Kuriyama Y. Complete nucleotide sequence of two steroid 21-hydroxylase genes tandemly arranged in human chromosome: a pseudogene and a genuine gene. *Proc Natl Acad Sci U S A* 1986;83(9): p2841-5. [PMC free article: PMC323402] [PubMed: 3486422]
93. Human Gene Mutation Database (HGMD): 2010 update.
94. Werkmeister JW , New MI , Dupont B , White PC . Frequent deletion and duplication of the steroid 21-hydroxylase genes. *Am J Hum Genet*. 1986 Oct;39(4):461-9. [PMC free article: PMC1683994] [PubMed: 3490178]
95. Finkelstein GP, Chen W, Mehta SP et al. Comprehensive genetic analysis of 182 unrelated families with congenital adrenal hyperplasia due to 21-hydroxylase deficiency. *J Clin Endocrinol Metab*. 2010 Oct 6 [Epub ahead of print]. [PMC free article: PMC3038490] [PubMed: 20926536]
96. Krone N, Braun A, Roscher A, Knorr D, Schwarz H. Predicting phenotype in steroid 21-hydroxylase deficiency? Comprehensive genotyping in 155 unrelated, well defined patients from southern Germany. *J Clin Endocrinol Metab* 2000;85(3): p1059-65. [PubMed: 10720040]
97. Rocha RO , Billerbeck AE , Pinto EM et al. The degree of external genitalia virilization in girls with 21-hydroxylase deficiency appears to be influenced by the CAG repeats in the androgen receptor gene. *Clin Endocrinol (Oxf)*. 2008 Feb;68(2):226-32. [PubMed: 17803691]
98. Forest MG. Recent advances in the diagnosis and management of congenital adrenal hyperplasia due to 21-hydroxylase deficiency.

- Hum Reprod Update. 2004 Nov-Dec;10(6):469-85. [PubMed: 15514016]
99. Pang S, Levine LS, Chow DM, Faiman C, New MI 1979 Serum androgen concentrations in neonates and young infants with congenital adrenal hyperplasia due to 21-hydroxylase deficiency. *Clin Endocrinol (Oxf)* 11:575-84. [PubMed: 161209]
 100. Korth-Schutz S, Virdis R, Saenger P, Chow DM, Levine LS, New MI 1978 Serum androgens as a continuing index of adequacy of treatment of congenital adrenal hyperplasia. *J Clin Endocrinol Metab* 46:452-8. [PubMed: 156192]
 101. Caulfield MP, Lynn T, Gottschalk ME, et al. 2002 The diagnosis of congenital adrenal hyperplasia in the newborn by gas chromatography/mass spectrometry analysis of random urine specimens. *J Clin Endocrinol Metab* 87:3682-90. [PubMed: 12161496]
 102. Shackleton CH. Profiling steroid hormones and urinary steroids. *J Chromatogr.* 1986 Jun 20;379:91-156. Review. [PubMed: 3525596]
 103. Pang S, Hotchkiss J, Drash AL, Levine LS, New MI 1977 Microfilter paper method for 17 alpha-hydroxyprogesterone radioimmunoassay: its application for rapid screening for congenital adrenal hyperplasia. *J Clin Endocrinol Metab* 45:1003-8. [PubMed: 925125]
 104. Speiser P, Azziz R, Baskin L, et al. Congenital Adrenal Hyperplasia due to 21-Hydroxylase Deficiency: An Endocrine Society Clinical Practice Guideline. *J Clin Endocrinol Metab*, September 2010, 95(9):4133-4160. [PMC free article: PMC2936060] [PubMed: 20823466]
 105. Gruneiro-Papendieck L, Prieto L, Chiesa A, Bengolea S, Bossi G, Bergada C 2001 Neonatal screening program for congenital adrenal hyperplasia: adjustments to the recall protocol. *Horm Res* 55:271-7. [PubMed: 11805430]
 106. Janzen N, Peter M, Sander S et al. Newborn screening for congenital adrenal hyperplasia: additional steroid profile using liquid chromatography-tandem mass spectrometry. *J Clin Endocrinol Metab.* 2007 Jul;92(7):2581-9. [PubMed: 17456574]
 107. New MI, Lorenzen F, Lerner AJ, et al. 1983 Genotyping steroid 21-hydroxylase deficiency: hormonal reference data. *J Clin Endocrinol Metab* 57:320-6. [PubMed: 6306039]
 108. Jeffcoate T, Fleigner J, Russell S, Davis J, Wade A 1965 Diagnosis of the adrenogenital syndrome before birth. *Lancet* 2:553. [PubMed: 4158268]
 109. Wilson RC, Wei J-Q, Cheng KC et al. Rapid deoxyribonucleic acid analysis by allele specific polymerase chain reaction for detection of mutations in the steroid 21-hydroxylase gene. *J Clin Endocrinol Metab* 80: 1635-1640, 1995. [PubMed: 7745011]
 110. Tukul T, Uyguner O, Wei J, et al. 2003 A Novel semiquantitative polymerase chain reaction/enzyme digestion-based method for detection of large scale deletions/conversions of the CYP21 gene and mutation screening in Turkish families with 21-hydroxylase deficiency. *J Clin Endocrinol Metab* 88:5893-5897. [PubMed: 14671187]
 111. Mao R, Nelson L, Kates R, et al. 2002 Prenatal diagnosis of 21-hydroxylase deficiency caused by gene conversion and rearrangements: pitfalls and molecular diagnostic solutions. *Prenat Diagn* 22:1171-6. [PubMed: 12478627]
 112. Day DJ, Speiser PW, Schulze E, et al. 1996 Identification of non-amplifying CYP21 genes when using PCR-based diagnosis of 21-hydroxylase deficiency in congenital adrenal hyperplasia (CAH) affected pedigrees. *Hum Mol Genet* 5:2039-48. [PubMed: 8968761]
 113. 2000 Technical report: congenital adrenal hyperplasia. Section on Endocrinology and Committee on Genetics. *Pediatrics* 106:1511-8. [PubMed: 11099616]
 114. Lo D, Corbetta N, Chamberlain P et al. Presence of fetal DNA in maternal plasma and serum. *Lancet* 1997; 350: 485-87. [PubMed: 9274585]
 115. Bischoff FZ, Nguyen DD, Marquez Do D, Moise KJ Jr, Simpson JL, Elias S. Noninvasive Determination of fetal RhD Statuses Using Fetal DNA in Maternal Serum and PCR. *J Soc Gynecol Invest* 199;6: 64-69. 1999. [PubMed: 10205775]
 116. Lam KW, Jiang P, Liao GJ, Chan KC, Leung TY, Chiu RW, Lo YM. Noninvasive prenatal diagnosis of monogenic diseases by targeted massively parallel sequencing of maternal plasma: application to beta-thalassemia. *Clin Chem.* 2012 58(10): 1467-75. [PubMed: 22896714]
 117. Gonzalez-Gonzalez MC, Garcia-Hoyos M, Trujillo MJ, Rodriguez de Alba M, Lorda-Sanchez I, Diaz-Recasens J, Gallardo E, Ayuso C, Ramos C. Prenatal detection of a cystic fibrosis mutation in fetal DNA from maternal plasma. *Prenat Diagn.* 2002 22:946-948. [PubMed: 12378583]
 118. Johnson KL, Dukes KA, Vidaver J, LeShane ES, Ramirez I, Weber WD, Bischoff FZ, Hahn S, Sharma A, Dang DX, Hire LM, Bianchi DW, Simpson JL, Holzgreve W, Elias S, and Klinger KW. Interlaboratory comparison of fetal male DNA detection from

- common maternal plasma samples by real time PCR . *Clin Chem*. 50 2004 516-21. [PubMed: 14718394]
119. Bischoff FZ, Lewis DE, and Simpson JL. Cell-free fetal DNA in maternal blood: kinetics, source and structure . *Hum Reprod Update*. 11 2005 59-67. [PubMed: 15569699]
 120. Chiu RW, Lau TK, Cheung PT et al. Noninvasive prenatal exclusion of congenital adrenal hyperplasia by maternal plasma analysis: a feasibility study . *Clin Chem*. 48 2002 778-80. [PubMed: 11978609]
 121. Simpson JL. Preimplantation genetic diagnosis at 20 years. *Prenat Diagn*. 2010 Jul;30(7):682-95. [PubMed: 20572111]
 122. Fiorentino F, Biricik A, Nuccitelli A, et al. 2006 Strategies and clinical outcome of 250 cycles of Preimplantation Genetic Diagnosis for single gene disorders. *Hum Reprod* 21:670-84. [PubMed: 16311287]
 123. David M, Forest MG 1984 Prenatal treatment of congenital adrenal hyperplasia resulting from 21-hydroxylase deficiency. *J Pediatr* 105:799-803. [PubMed: 6334149]
 124. New MI, Carlson A, Obeid J et al. 2001 Prenatal diagnosis for congenital adrenal hyperplasia in 532 pregnancies. *J Clin Endocrinol Metab* 86:5651-5657. [PubMed: 11739415]
 125. Forest MG, David M and Morel Y (1993) Prenatal diagnosis and treatment of 21-hydroxylase deficiency. *J Steroid Biochem Mol Biol* 45, 75-82. [PubMed: 8481354]
 126. Mercado AB, Wilson RC, Cheng KC, Wei JQ, New MI 1995 Extensive personal experience: Prenatal treatment and diagnosis of congenital adrenal hyperplasia owing to steroid 21-hydroxylase deficiency. *J Clin Endocrinol Metab* 80:2014-2020. [PubMed: 7608248]
 127. Carlson AD, Obeid JS, Kanellopoulou N, Wilson RC, New MI Congenital adrenal hyperplasia: update on prenatal diagnosis and treatment. *J Steroid Biochem Mol Biol*. 1999 Apr-Jun;69(1-6):19-29. Review. [PubMed: 10418977]
 128. New MI, Abraham M, Yuen T, Lekarev O. “An Update on Prenatal Diagnosis and Treatment of Congenital Adrenal Hyperplasia.” *Seminars in Reproductive Medicine*. 2012 October; 30(5):396-9. [PubMed: 23044876]
 129. Saenger P 1984 Abnormal sex differentiation. *J Pediatr* 104:1-17. [PubMed: 6361226]
 130. Meyer-Bahlburg HFL, Dolezal C, Baker SW, Ehrhardt AA, New MI 2006 Gender development in women with congenital adrenal hyperplasia as a function of disorder severity. *Archives of Sexual Behavior* 35:667-684. [PubMed: 16902816]
 131. *American Journal of Bioethics*. 10(9). Multiple articles in issue devoted to the subject.
 132. Seckl JR 2004 Prenatal glucocorticoids and long-term programming. *Eur J Endocrinol* 151 Suppl 3:U49-62. [PubMed: 15554887]
 133. Seckl JR, Miller WL 1997 How safe is long-term prenatal glucocorticoid treatment? *JAMA* 277:1077-9. [PubMed: 9091698]
 134. Yeh TF, Lin YJ, Lin HC, et al. 2004 Outcomes at school age after postnatal dexamethasone therapy for lung disease of prematurity. *N Engl J Med* 350:1304-13. [PubMed: 15044641]
 135. Slotkin TA, Zhang J, McCook EC, Seidler FJ 1998 Glucocorticoid administration alters nuclear transcription factors in fetal rat brain: implications for the use of antenatal steroids. *Brain Res Dev Brain Res* 111:11-24. [PubMed: 9804869]
 136. Uno H, Lohmiller L, Thieme C, et al. 1990 Brain damage induced by prenatal exposure to dexamethasone in fetal rhesus macaques. I. Hippocampus. *Brain Res Dev Brain Res* 53:157-67. [PubMed: 2357788]
 137. Hirvikoski T, Nordenström A, Lindholm T et al. 2007 Cognitive functions in children at risk for congenital adrenal hyperplasia treated prenatally with dexamethasone. *J Clin Endocrinol Metab* 92:542-548. [PubMed: 17148562]
 138. Hirvikoski T, Nordenström A, Lindholm T et al. 2008 Long-term follow-up of prenatally treated children at risk for congenital adrenal hyperplasia : does dexamethasone cause behavioral problems ? *Eur J Endocrinol* 159:309-316. [PubMed: 18579553]
 139. Meyer-Bahlburg H, Dolezal C, Haggerty R et al. Cognitive and educational outcome of offspring from dexamethasone-treated pregnancies at risk for congenital adrenal hyperplasia due to 21-hydroxylase deficiency. Submitted for publication. [PMC free article: PMC3383400] [PubMed: 22549088]
 140. New M, Carlson A, Obeid J, et al. 2003 Update: Prenatal diagnosis for congenital adrenal hyperplasia in 595 pregnancies. *The Endocrinologist* 13:233-239.
 141. Forest MG, Dorr HG 2003 Prenatal therapy in congenital adrenal hyperplasia due to 21-hydroxylase deficiency: Retrospective follow-up study of 253 treated pregnancies in 215 families. *Endocrinologist* 13:252-259.
 142. Trautman PD, Meyer-Bahlburg HFL, Postelnek J, New MI 1996 Mothers’ reactions to prenatal diagnostic procedures and dexamethasone treatment of CAH. *J Psycho Obst Gyn* 17:175-81. [PubMed: 8892164]
 143. Lajic S, Nordenstrom A, Ritzen EM, Wedell A 2004 Prenatal treatment of congenital adrenal hyperplasia. *Eur J Endocrinol* 151 Suppl 3:U63-9. [PubMed: 15554888]

144. Lajic S, Wedell A, Bui T, Ritzen E, Holst M 1998 Long-term somatic follow-up of prenatally treated children with congenital adrenal hyperplasia. *J Clin Endocrinol Metab* 83:3872-80. [PubMed: 9814461]
145. Goldman A, Sharpior B, Katsumata M 1978 Human foetal palatal corticoid receptors and teratogens for cleft palate. *Nature* 272:464-6. [PubMed: 634373]
146. Nandi J, Bern HA, Biglieri EG, Pieprzyk JK 1967 In vitro steroidogenesis by the adrenal glands of mice. *Endocrinology* 80:576-82. [PubMed: 5336765]
147. New MI. Long Range Outcome of Prenatal Treatment. Hormonal and Genetic Basis of Sexual Differentiation Disorders and Hot Topics in Endocrinology. 2nd World Conference. Miami, FL. January 2010.
148. Rosler A, Leiberman E, Rosenmann A 1979 Prenatal diagnosis of 11beta hydroxylase deficiency congenital adrenal hyperplasia. *J Clin Endocrinol Metab* 49:546. [PubMed: 314453]
149. Schumert Z, Rosenmann A, Landau H, Rosler A 1980 11-Deoxycortisol in amniotic fluid: Prenatal diagnosis of congenital adrenal hyperplasia due to 11 beta-hydroxylase deficiency. *Clin Endocrinol* 12:257. [PubMed: 6966984]
150. Sippell WG, Muller-Holve W, Dorr HG, Bidlingmaier F, Knorr D 1981 Concentrations of aldosterone, corticosterone, 11-deoxycorticosterone, progesterone, 17-hydroxyprogesterone, 11-deoxycortisol, cortisol, and cortisone determined simultaneously in human amniotic fluid throughout gestation. *J Clin Endocrinol Metab* 52:385-92. [PubMed: 7462398]
151. Cerase BI, Newfield RS, Pascoe L, et al. 1999 Prenatal diagnosis and treatment of 11beta-hydroxylase deficiency congenital adrenal hyperplasia resulting in normal female genitalia. *J Clin Endocrinol Metab* 84:3129-34. [PubMed: 10487675]
152. Geley S, Kapelari K, Johrer K, et al. 1996 CYP11B1 mutations causing congenital adrenal hyperplasia due to 11 beta-hydroxylase deficiency. *J Clin Endocrinol Metab* 81:2896-901. [PubMed: 8768848]
153. Motaghedi R, Betensky BP, Slowinska B, et al. 2005 Update on the prenatal diagnosis and treatment of congenital adrenal hyperplasia due to 11beta-hydroxylase deficiency. *J Pediatr Endocrinol Metab* 18:133-42. [PubMed: 15751602]
154. Nimkarn S, Lin-Su K, Berglind N, Wilson RC, New MI 2006 Aldosterone-to-Renin Ratio as a Marker for Disease Severity in 21-Hydroxylase Deficiency Congenital Adrenal Hyperplasia. *J Clin Endocrinol Metab*. [PubMed: 17032723]
155. Rosler A, Levine LS, Schneider B et al. The interrelationship of Sodium balance, plasma renin activity and ACTH in congenital adrenal hyperplasia. *Journal of Clin Endoc and Metab*, Spt 1977, Vol. 45, No. 3, pp 500-512. [PubMed: 198425]
156. New MI and Seaman MP. Secretion rates of cortisol and aldosterone precursors in various forms of congenital adrenal hyperplasia, *J. Clin. Endocrinol. Metab.* 30 (1970), pp. 361-371. [PubMed: 4312795]
157. Canalis E 1996 Clinical review 83: Mechanisms of glucocorticoid action in bone: implications to glucocorticoid-induced osteoporosis. *J Clin Endocrinol Metab* 81:3441-7. [PubMed: 8855781]
158. Canalis E, Bilezikian JP, Angeli A, Giustina A 2004 Perspectives on glucocorticoid-induced osteoporosis. *Bone* 34:593-8. [PubMed: 15050888]
159. Mazziotti G AA, Bilezikian JP, Canalis E, Giustina A 2006 Glucocorticoid-induced osteoporosis: an update. *Trend Endocrinol Metab* 17:144-9. [PubMed: 16678739]
160. Cameron F, Kaymakci B, Byrt E, Ebeling P, Warne G, Wark J 1995 Bone mineral density and body composition in congenital adrenal hyperplasia. *J Clin Endocrinol Metab* 80:2238-43. [PubMed: 7608286]
161. Guo C, Weetman A, Eastell R 1996 Bone turnover and bone mineral density in patients with congenital adrenal hyperplasia. *Clin Endocrinol (Oxf)* 45:535-41. [PubMed: 8977749]
162. de Almeida Freire PO, de Lemos-Marini SH, Maciel-Guerra AT, et al. 2003 Classical congenital adrenal hyperplasia due to 21-hydroxylase deficiency: a cross-sectional study of factors involved in bone mineral density. *J Bone Miner Metab* 21:396-401. [PubMed: 14586796]
163. King JA, Wisniewski AB, Bankowski BJ, Carson KA, Zacur HA, Migeon CJ 2006 Long-term corticosteroid replacement and bone mineral density in adult women with classical congenital adrenal hyperplasia. *J Clin Endocrinol Metab* 91:865-9. [PubMed: 16278269]
164. Lin-Su K and New MI. Effects of adrenal steroids on the bone metabolism of children with congenital adrenal hyperplasia. *Ann N Y Acad Sci.* 2007 Nov;1117:345-51. [PubMed: 18056052]
165. Crouch NS, Liao LM, Woodhouse CR, Conway GS, Creighton SM. Sexual function and genital sensitivity following feminizing genitoplasty for congenital adrenal hyperplasia. *J Urol.* 2008 Feb;179(2):634-8. Epub 2007 Dec 21. [PubMed: 18082214]
166. Nordenstrom A, Frisen L, Falhammar H et al. Sexual function and surgical outcome in women with congenital adrenal

hyperplasia due to CYP21A2 deficiency : clinical perspective and the patients' perception. *J Clin Endocrinol Metab* 95: 3633-3640, 2010. [PubMed: 20466782]

167. VanWyk JJ, Gunther DF, Ritzen M et al. The use of adrenalectomy as a treatment for congenital adrenal hyperplasia. *J Clin Endocrinol Metab* 81: 3180-3182, 1996. [PubMed: 8784066]
168. Gmyrek G, New M, Sosa R, Poppas D 2002 Bilateral laparoscopic adrenalectomy as a treatment for classic congenital adrenal hyperplasia attributable to 21-hydroxylase deficiency. *Pediatrics* 109: pE28. [PubMed: 11826238]

Publication Details

Author Information

Maria New, MD

Dr. Maria New is Professor of Pediatrics, Professor of Genetics and Genomic Sciences, and Director of the Adrenal Steroid Disorders Program at Mount Sinai School of Medicine in New York City. She is also serving as Associate Dean for Clinical Research at the Florida International University Herbert Wertheim College of Medicine.

Oksana Lekarev, DO.

Clinical Instructor of Pediatrics. Division of Pediatric Endocrinology. Mount Sinai School of Medicine, New York, NY

Karen Lin-Su, MD

Assistant Professor of Pediatrics. Division of Pediatric Endocrinology. Weill Cornell Medical College, New York, NY

Alan Parsa, M.D.

Research Fellow. Division of Pediatric Endocrinology. Mount Sinai School of Medicine, New York, NY

Ahmed Khattab, MD

Division of Endocrinology, Diabetes, and Bone Disease. Mount Sinai School of Medicine, New York, NY ,Division of Pediatric Endocrinology

Christian Pina

Division of Endocrinology, Diabetes, and Bone Disease. Mount Sinai School of Medicine, New York, NY ,Division of Pediatric Endocrinology

Tony Yuen, PhD

Division of Endocrinology, Diabetes, and Bone Disease. Mount Sinai School of Medicine, New York, NY ,Division of Pediatric Endocrinology

Mabel Yau, MD

Division of Endocrinology, Diabetes, and Bone Disease. Mount Sinai School of Medicine, New York, NY ,Division of Pediatric Endocrinology

Publication History

Last Update: October 28, 2013.

Copyright

Copyright © 2000-2015, MDText.com, Inc.

This electronic version has been made freely available under a Creative Commons (CC-BY-NC-ND) license. A copy of the license can be viewed at <http://creativecommons.org/licenses/by-nc-nd/2.0/>.

Publisher

MDText.com, Inc., South Dartmouth (MA)

NLM Citation

New M, Lekarev O, Lin-Su K, et al. Congenital Adrenal Hyperplasia. [Updated 2013 Oct 28]. In: De Groot LJ, Beck-Peccoz P, Chrousos G, et al., editors. *Endotext* [Internet]. South Dartmouth (MA): MDText.com, Inc.; 2000-.