

IOC Consensus Statement: Dietary Supplements and the High-Performance Athlete

Ronald J. Maughan
St Andrews University

Louise M. Burke
Australian Institute of Sport and
Mary MacKillop Institute
for Health Research

Jiri Dvorak
Schulthess Clinic

D. Enette Larson-Meyer
University of Wyoming

Peter Peeling
University of Western Australia and
Western Australian Institute of Sport

Stuart M. Phillips
McMaster University

Eric S. Rawson
Messiah College

Neil P. Walsh
Bangor University

Ina Garthe
Norwegian Olympic and Paralympic
Committee and Confederation of Sport

Hans Geyer
German Sport University

Romain Meeusen
Vrije Universiteit Brussel

Luc van Loon
Mary MacKillop Institute
for Health Research and
Maastricht University Medical Centre

Susan M. Shirreffs
St Andrews University

Lawrence L. Spriet
University of Guelph

Mark Stuart
BMJ, London

Alan Vernec
World Anti-Doping Agency (WADA)

Kevin Currell
English Institute of Sport,
Loughborough

**Vidya M. Ali and
Richard G.M. Budgett**
International Olympic Committee

Arne Ljungqvist
Anti-Doping Foundation

Margo Mountjoy
University of Guelph

**Yannis Pitsiladis,
Torbjørn Soligard, Uğur Erdener,
and Lars Engebretsen**
International Olympic Committee

Nutrition usually makes a small but potentially valuable contribution to successful performance in elite athletes, and dietary supplements can make a minor contribution to this nutrition program. Nonetheless, supplement use is widespread at all levels of sport. Products described as supplements target different issues, including the management of micronutrient deficiencies, supply of convenient forms of energy and macronutrients, and provision of direct benefits to performance or indirect benefits such as

Note. This article is co-published with the *British Journal of Sports Medicine* (April 2018, Vol. 52, Issue 7; doi:10.1136/bjsports-2018-099027). Maughan and Shirreffs are with St Andrews University, St Andrews, UK. Burke is with the Australian Institute of Sport, Canberra, Australia. Burke and van Loon are with the Mary MacKillop Institute for Health Research, Melbourne, Australia. Dvorak is with Schulthess Clinic, Zurich, Switzerland. Larson-Meyer is with the University of Wyoming, Laramie, WY. Peeling is with The University of Western Australia, Crawley, Australia, and with the Western Australian Institute of Sport, Mt Claremont, Australia. Phillips is with McMaster University, Hamilton, Ontario, Canada. Rawson is with Messiah College, Mechanicsburg, PA. Walsh is with Bangor University, Bangor, United Kingdom. Garthe is with the Norwegian Olympic and Paralympic Committee and Confederation of Sport, Oslo, Norway. Geyer is with German Sport University, Cologne, Germany. Meeusen is with Vrije Universiteit Brussel, Brussels, Belgium. van Loon is also with Maastricht University Medical Centre, Maastricht, The Netherlands. Spriet is with the University of Guelph, Guelph, Ontario, Canada. Stuart is with *BMJ*, London, United Kingdom. Vernec is with the World Anti-Doping Agency, Montreal, Quebec, Canada. Currell is with the English Institute of Sport, Loughborough, United Kingdom. Ali, Budgett, Pitsiladis, Soligard, Erdener, and Engebretsen are with the International Olympic Committee, Lausanne, Switzerland. Ljungqvist is with the The Professor Arne Ljungqvist Anti-Doping Foundation, Stockholm, Sweden. Mountjoy is with the University of Guelph, Guelph, Ontario, Canada. Address author correspondence to Susan M. Shirreffs at s.shirreffs@st-andrews.ac.uk.

- (b) For convenient provision of energy and nutrients around an exercise session
- (c) To achieve a specific and direct performance benefit in competition
- (d) To gain a performance improvement indirectly accrued from outcomes such as allowing more effective training (i.e., higher intensity, greater volume), better recovery from training sessions, optimizing mass and body composition, or reducing risks of injury and illness
- (e) For financial gain (sponsorship) or because products are provided free of charge
- (f) As a “just in case” insurance policy
- (g) Because they know or believe that other athletes/competitors are using the supplement(s)


Some supplements may be used for multiple functions. Zinc, for example, may be taken with the aim of promoting wound healing and tissue repair (Prasad, 2014) or reducing the severity and duration of the symptoms of an upper respiratory tract infection (Solomons, 1998). Carbohydrate supplements are used to enhance performance in many events via the provision of fuel substrate (Stellingwerff & Cox, 2014), to support the immune system (Peake et al., 2017), or to improve bioavailability of other supplements (e.g., creatine; Steenge et al., 2000). Similarly, creatine supplementation may directly enhance performance in strength and power events, and can assist in training harder, gaining lean body mass, or maintaining lean mass during periods of immobilization after injury (Branch, 2003; Gualano et al., 2012; Heaton et al., 2017). Decisions on supplement use therefore need to consider both the context of use and the specific protocol employed.

Assessing the Evidence Base for Supplement Use

Supplements target a range of scenarios of use, so different approaches are needed to assess their effectiveness. Supplements aimed at correcting nutrient deficiencies need to be judged on their ability to prevent or treat suboptimal nutrient status, with the benefit accruing from the removal of the associated impairment of health, training capacity, or performance. The effectiveness of sports foods might be hard to isolate when they are used within the general diet to meet everyday energy needs and nutrient targets. However, benefits may be more easily detected when they are specifically consumed before, during, or after an event or training session to provide nutrients that are limiting for performance (e.g., to provide fuel for the muscle or brain) or to defend homeostasis (e.g., by replacing water and salt losses). Performance-enhancing supplements which are claimed to achieve direct or indirect benefits pose a greater challenge in terms of a sound evidence base. With only a few exceptions, there is a scarcity of research, and many of the available studies are not of sufficient quality to warrant their application to elite athletes.

Substantiating the claims made about performance supplements and sports foods is difficult (Burke & Peeling, 2018). To various audiences, “proof” comes in different forms. Figure 1 provides a proposed hierarchical model of the relative strength of the evidence provided by different information sources. However, most of the information around supplement efficacy in sport comes from models with the lowest rigor: anecdotes/observations from athletes and scientific or mechanistic hypotheses that explain how a supplement might target a critical/limiting factor in performance, but with little to no evidence. Systematic reviews and

Hierarchy of Scientific Evidence



Evidence matrix for performance supplements




Figure 1 — Hierarchy of evidence used to establish good practice. Adapted from Burke and Peeling (2018).

Downloaded by on 10/04/18, Volume \$[article.issue.volume], Article Number \$[article.issue.issue]

Table 2 Summary of Common Sports Foods and Functional Foods Used by Athletes

Sports Food	Form	Typical Composition	Common Sports-Related Use
Sports drink	Powder or ready to drink liquid	5–8% CHO, 10–35 mmol/L sodium, 3–5 mmol/L potassium	Simultaneous delivery of fluid + CHO during exercise; postexercise rehydration and refueling
Energy drink	Ready to drink liquid or concentrated shot	Carbohydrate, especially in typical ready to drink varieties, caffeine <i>Note:</i> May contain taurine, B vitamins, and other ingredients with variable supporting evidence and some level of concern	Preexercise caffeine supplement; carbohydrate and caffeine intake during exercise
Sports gel or sports confectionery	Gel: 30–40 g sachets Confectionery: Jelly-type confectionery (generally in pouch of ~40–50 g)	~25 g CHO per sachet; ~5 g CHO per confectionery piece (some contain caffeine or electrolytes)	Carbohydrate intake during exercise
Electrolyte replacement supplements	Powder sachets or tablets	50–60 mmol/L sodium, 10–20 mmol/L potassium, typically low CHO (2–4 g/100 mL)	Rapid rehydration following dehydration undertaken for weight-making; replacement of large sodium losses during ultra-endurance activities; rapid postexercise rehydration following moderate to large fluid and sodium deficits
Protein supplement	Powder (mix with water or milk) or ready-to-drink liquid; protein-rich bar, usually low in CHO	Provides 20–50 g protein in a single serve from high-quality types of animal (whey, casein, milk, egg) or vegetable (e.g., soy) origin <i>Note:</i> May contain other ingredients, some of which are not evidence based and may increase the risk of contamination	Postexercise recovery following key training sessions or events where adaptation requiring protein synthesis is desired; achievement of increase in lean mass during growth or response to resistance training; portable nutrition for busy schedule or travel
Liquid meal supplement	Powder (mix with water or milk) or ready-to-drink liquid	1–1.5 kcal/mL: 15–20% protein and 50–70% CHO; low to moderate fat; vitamins/minerals: 500–1000 mL supplies RDI/RDAs	Supplement high-energy diet (especially during heavy training/competition or weight gain); low-bulk meal replacement (especially preevent meal); postexercise recovery (CHO and protein); portable nutrition for busy schedule or travel
Sports bar	Bar	40–50 g CHO, 5–10 g protein; usually low in fat and fiber; vitamins/minerals: 50–100% of RDA/RDIs <i>Note:</i> May contain other ingredients, some of which are not evidence-based and may increase the risk of contamination	CHO source during exercise; postexercise recovery (provides CHO, protein, and micronutrients); portable nutrition for busy schedule or travel
Protein-enhanced food	Milk, yogurt, ice cream, cereal bars, and other food forms	Increased protein content from normal food variety achieved by adding protein sources or filtration of water from product; typically allows normal portion to provide ~20 g protein to meet sports nutrition target	Value added food able to achieve protein target for postexercise use or to improve protein content of other meals and snacks in athlete's diet

Abbreviations: CHO = carbohydrate; RDA = recommended dietary allowance; RDI = reference daily intake.

Supplements That Directly Improve Sports Performance

A few performance-enhancing supplements might, at the present time, be considered to have an adequate level of support to suggest that marginal performance gains may be possible. These supplements include: caffeine, creatine (in the form of creatine monohydrate), nitrate, sodium bicarbonate, and possibly also β -alanine. The mechanisms of action, typical dose, potential performance benefits, and known side-effects of each of these

supplements are summarized in Table 3. Performance-enhancing supplements should be considered only where a strong evidence-base supports their use as safe, legal, and effective, and, ideally, after adequacy of sports nutrition dietary practices is ensured. Whenever possible, supplements should be trialed thoroughly by the athlete in training that mimics the competition milieu as closely as possible before committing to use in a competition setting. Athletes should do a careful risk analysis to see if the marginal gains would outweigh the risk of inadvertent doping due to contamination.

Table 3 Supplements With Good to Strong Evidence of Achieving Benefits to Performance When Used in Specific Scenarios (see Peeling et al., 2018 for further details)

Caffeine	
Overview	Caffeine is a stimulant that possesses well-established benefits for athletic performance across endurance-based situations, and short-term, supramaximal, and/or repeated sprint tasks.
Mechanism	Adenosine receptor antagonism; increased endorphin release; improved neuromuscular function; improved vigilance and alertness; reduce perception of exertion during exercise (Burke, 2008; Spriet, 2014)
Protocol of use	3–6 mg/kg of BM, in the form of anhydrous caffeine (i.e., pill or powder form), consumed ~60 min prior to exercise (Ganio et al., 2009). Lower caffeine doses (<3 mg/kg BM, ~200 mg), provided both before and during exercise; consumed with a CHO source (Spriet, 2014).
Performance impact	Improved endurance capacity such as exercise time to fatigue (French et al., 1991), and endurance-based TT activities of varying duration (5–150 min), across numerous exercise modalities (i.e., cycling, running, rowing, and others; Ganio et al., 2009). Low-doses of caffeine (100–300 mg) consumed during endurance exercise (after 15–80 min of activity) may enhance cycling TT performance by 3–7% (Paton et al., 2015; Talanian & Spriet, 2016). During short-term, supramaximal, and repeated sprint tasks, 3–6 mg/kg BM of caffeine taken 50–60 min before exercise results in performance gains of >3% for task completion time, mean power output, and peak power output during anaerobic activities of 1–2 min in duration (Wiles et al., 2006), and of 1–8% for total work output and repeat sprint performances during intermittent team game activity (Schneiker et al., 2006; Wellington et al., 2017).
Further considerations and potential side effects	Larger caffeine doses (≥ 9 mg/kg BM) do not appear to increase the performance benefit (Bruce et al., 2000), and are more likely to increase the risk of negative side effects, including nausea, anxiety, insomnia, and restlessness (Burke, 2008). Lower caffeine doses, variations in the timing of intake before and/or during exercise, and the need for (or lack thereof) a caffeine withdrawal period should be trialed in training prior to competition use. Caffeine consumption during activity should be considered concurrent with CHO intake for improved efficacy (Talanian & Spriet, 2016). Caffeine is a diuretic, promoting increased urine flow, but this effect is small at the doses that have been shown to enhance performance (Maughan & Griffin, 2003).
Creatine	
Overview	Creatine loading can acutely enhance the performance of sports involving repeated high-intensity exercise (e.g., team sports), as well as the chronic outcomes of training programs based on these characteristics (e.g., resistance or interval training), leading to greater gains in lean mass and muscular strength and power (Rawson & Persky, 2007; Volek & Rawson, 2004).
Mechanism	Supplementation increases muscle creatine stores, augmenting the rate of PCr resynthesis, thereby enhancing short-term, high-intensity exercise capacity (Buford et al., 2007) and the ability to perform repeated bouts of high-intensity effort.
Protocol of use	Loading-phase: ~20 g/day (divided into 4 equal daily doses), for 5–7 days (Lanthers et al., 2017) Maintenance-phase: 3–5 g/day (single dose) for the duration of the supplementation period (Hultman et al., 1996). <i>Note:</i> Concurrent consumption with a mixed protein/CHO source (~50 g of protein and CHO) may enhance muscle creatine uptake via insulin stimulation (Steenge et al., 2000).
Performance impact	Enhanced maximum isometric strength (Maganaris & Maughan, 1998) and the acute performance of single and repeated bouts of high-intensity exercise (<150 s duration); most pronounced effects evident during tasks <30 s (Branch, 2003; Lanthers et al., 2017). Chronic training adaptations include lean mass gains and improvements to muscular strength and power (Rawson & Persky, 2007; Volek & Rawson, 2004). Less common: Enhanced endurance performance resulting from increased/improved protein synthesis, glycogen storage, and thermoregulation (Cooper et al., 2012; Kreider et al., 2017). Potential anti-inflammatory and anti-oxidant effects are noted (Deminice et al., 2013).
Further considerations and potential side effects	No negative health effects are noted with long-term use (up to 4 years) when appropriate loading protocols are followed (Schilling et al., 2001). A potential 1–2 kg BM increase after creatine loading (primarily as a result of water retention; Deminice et al., 2013; Powers et al., 2003), may be detrimental for endurance performance or in events where the BM must be moved against gravity (e.g., high jump, pole vault) or where athletes must achieve a specific BM target.
Nitrate	
Overview	Dietary nitrate (NO ₃ ⁻) is a popular supplement that has been commonly investigated to assess any benefits for prolonged submaximal exercise (Bailey et al., 2009) and high-intensity, intermittent, short-duration efforts (Thompson et al., 2015; Wylie et al., 2016).
Mechanism	Enhances nitric oxide (NO) bioavailability via the NO ₃ -nitrite-NO pathway, playing an important role in the modulation of skeletal muscle function (Jones, 2014a). Nitrate augments exercise performance via an enhanced function of type II muscle fibers (Bailey et al., 2015); a reduced ATP cost of muscle force production; an increased efficiency of mitochondrial respiration; an increased blood flow to the muscle; a decrease in blood flow to VO ₂ heterogeneities (Bailey et al., 2010).
Protocol of use	High nitrate containing foods include leafy green and root vegetables, including spinach, rocket salad, celery, and beetroot. Acute performance benefits are generally seen within 2–3 hr following a NO ₃ ⁻ bolus of 5–9 mmol (310–560 mg) (Hoon et al., 2014). Prolonged periods of NO ₃ ⁻ intake (>3 days) also appears beneficial to performance (Thompson et al., 2015, 2016), and may be a positive strategy for highly-trained athletes, where performance gains from NO ₃ ⁻ supplementation appear harder to obtain (Jones, 2014b).
Performance impact	Supplementation has been associated with improvements of 4–25% in exercise time to exhaustion and of 1–3% in sport-specific TT performances lasting <40 min in duration (Bailey et al., 2015; McMahon et al., 2016). Supplementation is proposed to enhance type II muscle fiber function (Bailey et al., 2015), resulting in the improvement (3–5%) of high-intensity, intermittent, team-sport exercise of 12–40 min in duration (Thompson et al., 2015; Wylie et al., 2016). Evidence is equivocal for any benefit to exercise tasks lasting <12 min (Reynolds et al., 2016; Thompson et al., 2016).

(continued)

Table 3 (continued)

Further considerations and potential side effects	The available evidence suggests there appears to be few side effects or limitations to nitrate supplementation. There may exist the potential for GI upset in susceptible athletes, and should therefore be thoroughly trialed in training. There appears to be an upper limit to the benefits of consumption (i.e., no greater benefit from 16.8 mmol [1,041 mg] vs. 8.4 mmol [521 mg]; Wylie et al., 2013). Performance gains appear harder to obtain in highly-trained athletes (Jones, 2014b).
Beta-Alanine	
Overview	Beta-alanine augments intracellular buffering capacity, having potential beneficial effects on sustained high-intensity exercise performance.
Mechanism	A rate-limiting precursor to the endogenous intracellular (muscle) buffer, carnosine; the immediate defense against proton accumulation in the contracting musculature during exercise (Lancha Junior et al., 2015). Chronic, daily supplementation of beta-alanine increases skeletal muscle carnosine content (Saunders et al., 2016).
Protocol of use	Daily consumption of ~65 mg/kg BM, ingested via a split-dose regimen (i.e., 0.8–1.6 g every 3–4 hr) over an extended supplement time frame of 10–12 weeks (Saunders et al., 2016).
Performance Impact	Small but potentially meaningful performance benefits (~0.2–3%) during both continuous and intermittent exercise tasks of 30 s to 10 min in duration (Baguet et al., 2010 ; Chung et al., 2012 ; Saunders et al., 2016).
Further considerations and potential side effects	A positive correlation between the magnitude of muscle carnosine change and performance benefit remains to be established (Saunders et al., 2016). Large interindividual variations in muscle carnosine synthesis have been reported (Nassis et al., 2016). The supplement effectiveness appears harder to realize in well-trained athletes (Bellinger, 2014). There is a need for further investigation to establish the practical use in various sport-specific situations (Hobson et al., 2012 ; Saunders et al., 2016). Possible negative side effects include skin rashes and/or transient paresthesia.
Sodium Bicarbonate	
Overview	Sodium bicarbonate augments extracellular buffering capacity, having potential beneficial effects on sustained high-intensity exercise performance.
Mechanism	Acts as an extracellular (blood) buffer, aiding intracellular pH regulation by raising the extracellular pH, and HCO ₃ ⁻ concentrations (Katz et al., 1984 ; Lancha Junior et al., 2015). The resultant pH gradient between the intracellular and extracellular environments leads to efflux of H ⁺ and La ⁻ from the exercising muscle (Katz et al., 1984 ; Mainwood et al., 1975).
Protocol of use	Single acute NaHCO ₃ dose of 0.2–0.4 g/kg BM, consumed 60–150 min prior to exercise (Carr et al., 2011b ; Sieglar et al., 2012). Alternative strategies include: split doses (i.e., several smaller doses giving the same total intake) taken over a 30–180 min time period (Lambert et al., 1993); serial-loading with 3–4 smaller doses per day for 2–4 consecutive days prior to an event (Burke, 2013 ; Douroudos et al., 2006 ; Mc Naughton & Thompson, 2001).
Performance impact	Enhanced performance (~2%) of short-term, high-intensity sprints lasting ~60 s in duration, with a reduced efficacy as the effort duration exceeds 10 min (Carr et al., 2011b).
Further considerations and potential side effects	Well-established GI distress may be associated with this supplement. Strategies to minimize GI upset include: co-ingestion with a small, carbohydrate-rich meal (~1.5 g/kg BM carbohydrates) (Carr et al., 2011c); the use of sodium citrate as an alternative (Requena et al., 2005); split-dose or stacking strategies (Burke, 2013 ; Douroudos et al., 2006 ; Mc Naughton & Thompson, 2001). Given the high potential for GI distress, thorough investigation into the best individualized strategy is recommended prior to use in a competition setting.

Abbreviations: ATP=adenosine triphosphate; BM=body mass; CHO=carbohydrate; GI=gastrointestinal; TT=time trial.

Supplements That Improve Performance Indirectly

Many dietary supplements claim to enhance performance indirectly by supporting the athlete's health, body composition, and their ability to train hard, recover quickly, adapt optimally, avoid or recover from injury, and tolerate pain or soreness. Illness is a major problem for athletes if it interrupts training or occurs at a critical time, such as during a selection event or a major competition. Susceptibility to illness is increased in situations where athletes are involved in a high volume of training or competition and either intentionally or unintentionally experience deficits in energy intake (e.g., weight-loss diets), macronutrient intake (e.g. train-low or sleep-low-carbohydrate), and micronutrient status (e.g., vitamin D insufficiency in the winter) ([Bermon et al., 2017](#)). Athletes might benefit from nutritional supplements to support immunity in these scenarios and at other times when they are either susceptible to infection (e.g., during the common cold season and after long-haul travel) or suffering from an infection. Table 4 summarizes evidence for some of the commonly promoted "immune supportive"

supplements, noting that the most promising candidates to assist in the prevention or treatment of upper respiratory symptoms are vitamin D and probiotics. Vitamin C during periods of heavy exertion and zinc lozenges at the onset of symptoms may be useful, but high doses of single antioxidants, particularly vitamin C and E, may blunt exercise-induced training adaptations ([Nikolaidis et al., 2012](#); [Paulsen et al., 2014](#); [Powers et al., 2011](#)). Probiotic supplementation may reduce the incidence of travelers' diarrhea and gastrointestinal infection. Cochrane reviews have noted the low quality of many studies on nutritional supplements that are claimed to support immunity; specifically, small samples, poor controls, and unclear procedures for randomization and blinding were commonplace ([Hao et al., 2015](#); [Singh & Das, 2013](#)). Clearly, there is a pressing need for randomized controlled trials in high-level athletes with sufficient participant numbers, rigorous controls and procedures, appropriate supplementation regimens, and clinically meaningful measures of immunity.

Supplements that assist an athlete to train harder, recover more quickly, and prevent injury or accelerate return to play when injury does occur can obviously enhance the athlete's preparation and, indirectly, their competition outcomes. Many products claim to

Table 4 Nutritional Supplements for Immune Health in Athletes: Proposed Mechanism of Action and Evidence for Efficacy

Supplement	Proposed Mechanism of Action	Evidence for Efficacy
Vitamin D	An essential fat-soluble vitamin known to influence several aspects of immunity, particularly innate immunity (e.g., expression of antimicrobial proteins). Skin exposure to sunlight accounts for 90% of the source of vitamin D.	Moderate support. Evidence for deficiency in some athletes and soldiers, particularly in the winter (decreased skin sunlight exposure). Deficiency has been associated with increased URS. Recommend 1,000 IU/day D ₃ autumn–spring to maintain sufficiency. Further support required (He et al., 2016).
Probiotics	Probiotics are live microorganisms which, when administered orally for several weeks, can increase the numbers of beneficial bacteria in the gut. This has been associated with a range of potential benefits to gut health, as well as modulation of immune function.	Moderate support in athletes with daily dose of ~10 ¹⁰ live bacteria; Cochrane review of 12 studies (n = 3,720) shows ~50% decrease in URS incidence and ~2 day shortening of URS; minor side effects. More evidence is required supporting efficacy to reduce gastrointestinal distress and infection (e.g., in a traveling athlete; Gleeson et al., 2011; Hao et al., 2015).
Vitamin C	An essential water-soluble antioxidant vitamin that quenches ROS and augments immunity. Reduces interleukin-6 and cortisol responses to exercise in humans.	Moderate support for preventing URS. Cochrane review of 5 studies in heavy exercisers (n = 598) shows ~50% decrease in URS taking vitamin C (0.25–1.0 g/day). Further support required. Unclear if antioxidants blunt adaptation in well-trained. Relatively small effects on cortisol compared with carbohydrate; immune measures no different from placebo. No support for treating URS. Cochrane reviews show no benefit of initiating vitamin C supplementation (>200 mg/day) after onset of URS (Hemila & Chalker, 2013; Nieman et al., 2002).
Carbohydrate (drinks, gels)	Maintains blood glucose during exercise, lowers stress hormones, and thus counters immune dysfunction.	Low-moderate support. Ingestion of carbohydrate (30–60 g/hr) attenuates stress hormone and some, but not all, immune perturbations during exercise. Very limited evidence that this modifies infection risk in athletes (Bermon et al., 2017; Walsh et al., 2011).
Bovine colostrum	First milk of the cow that contains antibodies, growth factors, and cytokines. Claimed to improve mucosal immunity and increase resistance to infection.	Low-moderate support that bovine colostrum blunts the decrease in saliva antimicrobial proteins after heavy exercise. Some evidence in small numbers of participants that bovine colostrum decreases URS. Further support required (Brinkworth & Buckley, 2003; Davison & Diment, 2010).
Polyphenols (e.g., Quercetin)	Plant flavonoids. In vitro studies show strong anti-inflammatory, anti-oxidant, and anti-pathogenic effects. Animal data indicate an increase in mitochondrial biogenesis and endurance performance.	Low-moderate support. Human studies show some reduction in URS during short periods of intensified training and mild stimulation of mitochondrial biogenesis and endurance performance, albeit in small numbers of untrained subjects. Limited influence on markers of immunity. Putative anti-viral effect for Quercetin. Further support required (Gleeson, 2016; Nieman et al., 2007).
Zinc	An essential mineral that is claimed to reduce incidence and duration of colds. Zinc is required for DNA synthesis and as an enzyme cofactor for immune cells. Zinc deficiency results in impaired immunity (e.g., lymphoid atrophy), and zinc deficiency is not uncommon in athletes.	No support for preventing URS. High doses of zinc can decrease immune function and should be avoided. Moderate support for treating URS. Cochrane review shows benefit of zinc acetate lozenges (75 mg) to decrease duration of URS; however, zinc must be taken <24 hr after onset of URS for duration of cold only. Side effects include bad taste and nausea (Singh & Das, 2013).
Glutamine	Nonessential amino acid that is an important energy substrate for immune cells, particularly lymphocytes. Circulating glutamine is lowered after prolonged exercise and very heavy training.	Limited support. Supplementation before and after exercise does not alter immune perturbations. Some evidence of a reduction in URS after endurance events in competitors receiving glutamine supplementation (2 × 5 g). Mechanism for therapeutic effect requires investigation (Castell et al., 1996; Walsh et al., 1998).
Caffeine	Stimulant found in a variety of foods and drinks (e.g., coffee and sports drinks). Caffeine is an adenosine receptor antagonist and immune cells express adenosine receptors.	Limited support. Evidence that caffeine supplementation activates lymphocytes and attenuates the fall in neutrophil function after exercise. Efficacy for altering URS in athletes remains unknown (Dulson & Bishop, 2016; Walker et al., 2007).
Echinacea	Herbal extract claimed to enhance immunity via stimulatory effects on macrophages. There is some in vitro evidence for this.	Limited support. Early human studies indicated possible beneficial effects but more recent, larger scale, and better controlled studies indicate no effect of Echinacea on infection incidence or cold symptom severity (Karsch-Volk et al., 2015; Linde et al., 2006).
Omega-3 PUFAs	Found in fish oil. May influence immune function by acting as a fuel, in their role as membrane constituents or by regulating eicosanoid formation (e.g., prostaglandin). Prostaglandin is immunosuppressive. Claimed to exert anti-inflammatory effects postexercise.	Limited support for blunting inflammation and functional changes after muscle-damaging eccentric exercise in humans and no evidence of reducing URS in athletes (Jakeman et al., 2017; Mickleborough, 2013).

(continued)

Table 4 (continued)

Supplement	Proposed Mechanism of Action	Evidence for Efficacy
Vitamin E	An essential fat-soluble antioxidant vitamin that quenches exercise-induced ROS and augments immunity.	No support. Immune-enhancing effects in the frail elderly but no benefit in young, healthy humans. One study actually showed that vitamin E supplementation increased URS in those under heavy exertion. High doses may be pro-oxidative (Hemila et al., 2003; Meydani et al., 2004).
β-glucans	Polysaccharides derived from the cell walls of yeast, fungi, algae, and oats that stimulate innate immunity.	No support in humans. Effective in mice inoculated with influenza virus; however, human studies with athletes show no benefits (Nieman et al., 2008; Volman et al., 2008).

Abbreviations: PUFA = polyunsaturated fatty acids; ROS = reactive oxygen species; URS = upper respiratory symptoms.

provide benefits of this nature. Table 5 summarizes the evidence for some of the most popular compounds. Finally, the manipulation of body composition, including gaining lean (muscle) mass and reducing body fat levels, can contribute to performance in many events. This explains the large number of “weight gainers” and “fat burners” in the general and sports supplement market, though many of these are prohibited in sport. Protein is considered to be the premier ingredient in weight-gain-promoting supplements and evidence-based reviews conclude that protein is effective at promoting lean mass gain when combined with resistive exercise (Hector & Philips, 2018). Evidence of efficacy for “fat burning” supplements is far from conclusive, however, and there is a complete absence of evidence for the effectiveness of the vast majority of supplements marketed in this category. Table 6 summarizes the evidence for some of the most common ingredients or products of this type (Hector & Philips, 2018).

Adverse Effects

Adverse effects from the use of supplements may arise from a number of factors, including the safety and composition of the product per se and inappropriate patterns of use by athletes. Poor practices by athletes include the indiscriminate mixing and matching of many products without regard to total doses of some ingredients or problematic interactions between ingredients. Even commonly-used products may have negative side effects, especially when used outside the optimal protocol. For example, iron supplementation in those with already adequate iron stores can result in symptoms that may begin with vomiting, diarrhea, and abdominal pain, and develop to haemochromatosis and liver failure (Mettler & Zimmermann, 2010). Bicarbonate may cause gastrointestinal distress when ingested in amounts sufficient to enhance performance; this can impair rather than improve performance and may counteract the benefits of other supplements taken at the same time (Carr et al., 2011a). The “more is better” philosophy, when applied to caffeine, may result in side effects, including nausea, anxiety, accelerated heart rate, and insomnia, which outweigh the performance benefits (Peeling et al., 2018). Unwanted outcomes become more common with caffeine doses ≥ 9 mg/kg body mass, but maximal benefits are usually achieved with intakes of 3–6 mg/kg (Burke, 2008). The possibility of more serious outcomes is illustrated by adverse, and potentially fatal, responses in two separate incidents in which very large doses (up to 30 grams) of caffeine were administered to healthy volunteers participating in laboratory studies (Bodkin, 2017). These incidents were due to errors in the dose calculation—if this can happen in a university research environment with supposed oversight by experienced staff, the potential clearly exists for similar errors by athletes and coaches.

Athletes and members of their support team should be aware of the regulations that govern the manufacture and marketing of supplements. According to the Dietary Supplements Health and Education Act 1994 (DSHEA) passed by US Congress, nutritional supplements sold in the United States that do not claim to diagnose, prevent, or cure disease are not subject to regulation by the Food and Drug Administration (FDA). Similar regulations apply in most other countries, where supplements are regulated in the same way as food ingredients and are therefore not subject to the stringent regulations that are applied to the pharmaceutical industry. This means that there is no requirement to prove claimed benefits, no requirement to show safety with acute or chronic administration, no quality assurance of content, and no liberal labeling requirements. It is well recognized that there are problems with some of the dietary supplements on sale, but the options open to those responsible for food safety are limited by the legislation that applies. The FDA regularly uses its powers to recall products in breach of the regulations, though they fully admit that their resources are insufficient for comprehensive monitoring, and recalls generally occur only after many people are harmed (<https://www.fda.gov/food/recallsoutbreaksemergencies/recalls/default.htm>). The FDA have recently recalled supplement products containing excessive doses of vitamins A, D, B6, and selenium because of potentially toxic levels of these components. Examples of product complaints have included the presence of impurities, including lead, broken glass, and metal fragments, because of the failure of the producers to observe good manufacturing practice. The risk of gastrointestinal upset because of poor hygiene during the production and storage of products is also of concern. Although this may seem a minor inconvenience, and of similarity to food safety issues, the coincidence of problems around a crucial training period or competitive event may significantly interfere with an athlete’s performance goals. It should be noted, though, that all of these problems are also regularly reported in normal foods.

Some supplements may actually cause harm to health, but these can be difficult to identify, and products are usually withdrawn only after a significant number of adverse events have occurred. For example, a range of products containing hydroxycitric acid were withdrawn from sale, but only after they were linked with the death of one consumer and with a substantial number of other cases of liver toxicity, cardiovascular problems, and seizures (<https://www.fda.gov/NewsEvents/PublicHealthFocus/ucm155817.htm>). The extent of the problem is illustrated by the fact that, in the United States in 2015, approximately 23,000 emergency department visits annually are reported to be associated with dietary supplement use (Geller et al., 2015). This figure can be viewed as substantial, or it can be seen as small compared to the total number of adverse responses associated with the use of medications (Maughan et al., 2018). However,

Table 5 Supplements That May Assist With Training Capacity, Recovery, Muscle Soreness, and Injury Management (see Rawson et al., 2018 for further details)

Supplement	Proposed Mechanism of Action	Evidence for Efficacy
<p>Creatine monohydrate: Creatine is a naturally occurring nutrient, consumed in the diet, and synthesized in the body. Recommended supplement dose is 20 g/day for 5 days, followed by 3 to 5 g/day to increase and maintain elevated body creatine levels (Harris et al., 1992; Hultman et al., 1996).</p>	<p>Enhanced adaptive response to exercise via increased growth factor/ gene expression, increased intracellular water; reduced symptoms of or enhanced recovery from muscle-damaging exercise (e.g., DOMS); enhanced recovery from disuse or immobilization/ extreme inactivity; improved cognitive processing; decreased risk/enhanced recovery from mTBI</p>	<p>Many studies demonstrate improved training adaptations, such as increased lean mass or strength, indicating an enhanced adaptive response to exercise (Branch, 2003; Heaton et al., 2017; Rawson & Volek, 2003). Reduced symptoms of, or enhanced recovery from, muscle-damaging exercise (e.g., DOMS) have been reported in some, but not all studies (reviewed in Rawson et al., 2017). Enhanced recovery from disuse or immobilization/extreme inactivity has been reported in some, but not all studies (reviewed in Heaton et al., 2017). Improved cognitive processing is reported in most studies, especially when volunteers were fatigued by sleep deprivation or mental/physical tasks (reviewed in Gualano et al., 2012; 2016; Rae & Broer, 2015; Rawson & Venezia, 2011). The effects in athletes have not been well characterized, and only one group attempted to translate these effects into to athletic performance, albeit with a positive result. (Cook et al., 2011). Decreased damage and enhanced recovery from mTBI is supported by open label trials in children (Sakellaris et al., 2006, 2008) and using animal models (Sullivan et al., 2000). These data are not conclusive and more research is warranted. However, athletes at risk for concussion, who already ingest creatine supplements for performance or muscular benefits, may receive important brain benefits as well. A small increase in body mass is common with supplementation. This may be relevant for sports with weight classes/restrictions or where increased body mass may decrease performance.</p>
<p>Beta-hydroxy beta-methylbutyrate (HMB): HMB is a metabolite of the amino acid leucine. Manufacturer recommended dosage is 3 g/day.</p>	<p>Enhanced adaptive response to exercise via decreased protein breakdown increased protein synthesis; increased cholesterol synthesis, increased growth hormone and IGF I mRNA, increased proliferation and differentiation of satellite cells, and inhibited apoptosis (reviewed in Szczesniak et al., 2015).</p>	<p>Beneficial effects of HMB on strength and fat free mass are small, while the effects on muscle damage are unclear (Rowlands & Thomson, 2009). Recent reports of “steroid-like” gains in strength, power, and fat free mass, and reductions in muscle damage from HMB-free acid (HMB-FA) supplementation (Lowery et al., 2016; Wilson et al., 2013, 2014) have not been reproduced and seem unlikely (Phillips et al., 2017). Potential use for HMB during extreme inactivity/disuse or recovery from injury, but these effects have only been described in older adults following 10 days of bed rest (Deutz et al., 2013). Benefits of HMB supplementation could most likely be obtained from normal dietary protein or whole protein supplements (Wilkinson et al., 2013), so HMB supplements may not be more effective than adhering to the current protein intake recommendations</p>
<p>Omega 3-fatty acids (about 2 g/day)</p>	<p>Improved cognitive processing; decreased risk/enhanced recovery from mTBI; increased muscle protein synthesis; reduced symptoms of or enhanced recovery from muscle-damaging exercise (e.g., DOMS).</p>	<p>Improved cognitive processing following omega 3-fatty acid supplementation shown in healthy older adults those with mild or severe cognitive impairment (reviewed in Barrett et al., 2014). It is not known if these benefits would occur in young, healthy athletes, or how this would translate to athletic performance. Animal data show that the structural damage and cognitive decline associated with mTBI are reduced/attenuated with omega-3 fatty acid supplementation when ingested either before or after the injury (reviewed in Barrett et al., 2014; Erdman et al., 2011; Tipton, 2015). Two case studies support these findings (Lewis et al., 2013; Roberts et al., 2008) and large, double-blind, placebo-controlled trials are currently under way (clinicaltrials.gov NCT101903525, NCT01814527). In muscle, omega-3 fatty acid supplementation can increase muscle protein synthesis (Smith et al., 2011a, 2011b), but this may not occur when protein is ingested after exercise in recommended amounts (Smith et al., 2011a, 2011b). Anti-inflammatory effects of omega-3 fatty acid intake may reduce muscle damage or enhance recovery from intense, eccentric exercise (e.g., decrease DOMS), but this is not a consistent finding (Gray et al., 2014; Jouris et al., 2011). No indication that decreased omega 3-fatty acids in the body impair performance, and high-dose supplements can cause some adverse effects (reviewed in Erdman et al., 2011; Mickleborough, 2013), so the best recommendation may be to include rich sources of omega-3 fatty acids, such as fatty fish, in the diet instead of supplements. Low risk but unclear if supplementation should be pursued by athletes, in lieu of including fatty fish in the diet as a source of omega-3 fatty acids. Fish oil or omega-3 fatty acid supplement consumption could include heavy metal contaminants, or cause bleeding, digestive problems, and/ or increased LDL.</p>

(continued)

Table 5 (continued)

Supplement	Proposed Mechanism of Action	Evidence for Efficacy
Vitamin D: An essential fat-soluble vitamin. Skin exposure to sunlight normally accounts for 90% of the source of vitamin D.	Enhanced adaptive response to exercise; decreased stress fractures	Data on the effects of vitamin D supplementation on muscle function and recovery are equivocal, with discrepancies likely explained by differences in baseline vitamin D concentrations prior to supplementation (Close et al., 2013, 2016; Owens et al., 2014, 2015). Collectively, these data strongly suggest a role for adequate vitamin D in the adaptive process to stressful exercise. Low vitamin D status is associated with a 3.6×higher stress fracture risk in Finnish military recruits (Ruohola et al., 2006). US Naval recruits supplemented with 800 IU/day of vitamin D3 and 2,000 mg calcium reduced stress fracture incidence by 20% (Lappe et al., 2008). More data are needed, but it appears that vitamin D status, relates to stress fracture risk, and supplementation, when warranted, may reduce this risk.
Gelatin and vitamin C/ collagen: Recommended dose is 5 to 15 g gelatin with 50 mg vitamin C (Shaw et al., 2017). Collagen hydrolysate dose is about 10 g/day (Clark et al., 2008; McAlindon et al., 2011)	Increased collagen production; thickened cartilage; decreased joint pain	Gelatin and collagen supplements appear to be low risk. Few data available (Clark et al., 2008; McAlindon et al., 2011; Shaw et al., 2017) but increased collagen production and decreased pain seem possible. Functional benefits, recovery from injury, and effects in elite athletes are not known.
Anti-inflammatory supplements: Curcumin (a constituent of the spice turmeric) is often ingested for anti-inflammatory effects at a dose of about 5 g/day. Tart cherry juice at a dose of about 250–350 mL (30 mL if concentrate) twice daily for 4 to 5 day before an athletic event or for 2 to 3 days afterwards to promote recovery.	Anti-inflammatory effects; reduced symptoms of or enhanced recovery from muscle-damaging exercise (e.g., DOMS)	Decreases in inflammatory cytokines and/or indirect markers of muscle damage with anti-inflammatory supplements such as curcumin (McFarlin et al., 2016; Nicol et al., 2015; Sciberras et al., 2015) and tart cherry juice (reviewed in Bell et al., 2014; Coelho Rabello Lima et al., 2015) have been reported. Anti-inflammatory effects may be beneficial, although benefits may be sport/training specific. More research is needed before these compounds can be recommended to athletes.

Abbreviations: DOMS = delayed onset muscle soreness; mTBI = mild traumatic brain injury (concussion).

minor problems that do not require acute medical aid may still be sufficient to interrupt training or prevent participation, so this statistic probably underestimates the risk for athletes.

The biggest concern for athletes who compete under an anti-doping code (usually the World Anti-Doping Code, as published by the World Anti-Doping Agency [WADA]) is that supplements can contain prohibited substances that result in an anti-doping rule violation (ADRV). Athletes—and their support teams—may be at risk for an ADRV if there is evidence that they have used or attempted to use products containing ingredients on the Prohibited List (www.wada.ama.org). A common problem is the recording of an adverse analytical finding (AAF) of a prohibited substance in a urine sample (“positive drug test”) as a result of supplement use (Maughan, 2005). Millions of athletes may be subject to anti-doping testing, although these are mostly professional-, national-, or international-level athletes. For these athletes in particular, even if the ingestion of the prohibited substance was unintentional, the rules of strict liability within the World Anti-Doping Code mean that an AAF will be recorded, and may mean the loss of medals won or records set, as well as financial sanctions and/or temporary or permanent suspension from competition. It also damages the athlete’s reputation and may lead to loss of employment and income through failed sponsorship opportunities. Where there has been deliberate cheating or benefit accrued from the use of a prohibited substance, these penalties seem entirely appropriate, but it is undoubtedly true that some ADRVs can be attributed to the innocent ingestion of prohibited substances in dietary supplements, with catastrophic results for the athlete.

One cause of an AAF arising from supplement use relates to an athlete’s failure to read product labels to recognize the presence of

prohibited substances. Many athletes consider supplements to be “natural” or “regulated” and therefore safe. Other athletes are confused by the number of chemical names for some prohibited substances and thus fail to recognize them on the product label. However, the most worrying cause of an inadvertent AAF is the use of supplements that contain prohibited substances as an undeclared ingredient or contaminant. Since the publication of the seminal study on the presence of undeclared prohibited substances in supplements (Geyer et al., 2004), there have been numerous reports of supplement contamination (Maughan, 2005). Recent reviews suggest that this problem remains (LGC, 2017; Martinez-Sanz et al., 2017). It is difficult to gain a perspective of the true prevalence of supplement contamination. Although the original study reported that ~15% of more than 600 products acquired from around the world contained undeclared prohormones (Geyer et al., 2004), this and other investigations rarely include a truly random sample of the supplements and sports foods used by athletes. Some individual products or categories of products can be considered inherently more at risk of contamination due to the country of origin, the manufacturer, the type of product, and the range of declared ingredients (<https://www.usada.org/substances/supplement-411/>). Nevertheless, it should also be recognized that common supplements, including vitamin C, multivitamins, and minerals, have also been found, albeit rarely, to contain prohibited substances (Geyer et al., 2008). The range of prohibited substances found as undeclared ingredients in supplements now includes products from many sections of the WADA’s List of Prohibited Substances and Methods, including stimulants, anabolic agents, selective androgen receptor modulators, diuretics, anorectics, and β_2 agonists (Martinez-Sanz et al., 2017).

Table 6 Supplements Promoted to Assist With Physique Changes: Gain in Lean Mass and Loss of Body Fat Mass (see Hector & Phillips, 2018 for further details)

Supplement	Proposed Mechanism of Action	Evidence for Efficacy
Gaining lean body mass*		
Protein Usually comprised of isolated proteins from various sources (whey and soy most common) <i>Recommended daily dose:</i> 1.6 g protein/kg/day optimal (up to 2.2 g/kg/day with no adverse effects) <i>Recommended per-meal doses:</i> 0.3–0.5 g protein/kg (3–4 times per day and in close temporal proximity to exercise, with postexercise being consistently shown to be effective)	Enhances lean mass gains when ingested during programs of resistance training due to increased provision of building blocks (amino acids) and leucine as a trigger for a rise in muscle protein synthesis and suppression of muscle protein breakdown	Meta-analyses focusing on younger and older participants have shown positive effects enhancing gains in muscle mass (Cermak et al., 2012; Morton et al., 2017), but effects are not large
Leucine	Stimulates muscle protein synthesis and suppresses protein breakdown (possibly through insulin)	Short-term mechanistic data available (Wilkinson et al., 2013), but no long-term trials showing efficacy (Aguilar et al., 2017)
Losing fat mass**		
Protein From increased dietary sources or supplemental isolated proteins	Enhance fat mass loss and promotes retention of lean mass	Meta-analyses confirm small but significant effects of greater dietary protein in weight loss to enhance fat mass loss and promote lean mass retention (Krieger et al., 2006; Wycherley et al., 2012)
Pyruvate	No data	Small-to-trivial effect (Onakpoya et al., 2014a)
Chromium	Potentiates biological actions of insulin	No effect (Tian et al., 2013)
Green tea (polyphenol catechins and caffeine)	Thermogenic agent and/or lipolytic-enhancing agent	Small-to-trivial effect (Jurgens et al., 2012)
α -Lipoic acid	No clear role, but possible antioxidant	Small-to-trivial effect (Kucukgoncu et al., 2017)
Conjugated linoleic acid (CLA)	Changes membrane fluidity favoring enhanced fat oxidation	Small-to-trivial effect (Onakpoya et al., 2012)
Konjac fiber (glucomannan)	Water-soluble polysaccharide—dietary fiber	Small-to-trivial effect (Onakpoya et al., 2014b)
Omega-3 polyunsaturated fatty acids	No clear role, but possible appetite suppression, improved blood flow, and/or modulator of gene expression	Small-to-trivial effect (Zhang et al., 2017)
Chitosan	Lipid-binding agent to reduce lipid absorption	Small-to-trivial effect (Jull et al., 2008)

*In combination with progressive resistance exercise programs. **In combination with an exercise- and/or diet-induced energy deficit.

In some cases, the amount of the prohibited substance in a supplement may be high, even higher than the normal therapeutic dose. For example, Geyer et al. (2002) reported the analysis of metandienone (commonly known as methandrostenolone or Dianabol) in high amounts in a “body building” supplement from England. The recommended amount of the supplement would have supplied a dose of 10–43 mg; in comparison, the typical therapeutic dose of this drug was 2.5–5 mg/day (Goodman & Gilman, 1975), though its medical use has been discontinued in most countries for many years. This amount would certainly have a potent anabolic effect, but would likely produce serious side effects, including psychiatric and behavioral effects and significant damage to a range of body systems including the liver (Solimini et al., 2017). Unlike many of the earlier cases involving steroids related to nandrolone and testosterone, this is not a trivial level of contamination and raises the possibility of deliberate adulteration of the product with the intention of producing a measurable effect on muscle strength and muscle mass. Most reports of adverse health outcomes resulting from supplement use have focused on liver problems of varying

degrees of severity, but other organs are also affected. One epidemiological case-control study (Li et al., 2015) examined the association between use of muscle-building supplements and testicular germ cell cancer (TGCC) risk, with 356 TGCC cases and 513 controls from the Eastern United States. The odds ratio (OR) for the use of muscle-building supplements in relation to risk of TGCC was elevated (OR = 1.65, 95% confidence interval [CI]: 1.11–2.46), with significantly stronger associations for early users and longer periods of use.

Ironically, supplements that are contaminated with extremely small amounts of prohibited substances (too low to have any physiological effect) may still cause a positive doping outcome. For instance, ingestion of 19-norandrostenedione, a precursor of nandrolone, will result in the appearance in the urine of 19-norandrosterone, the diagnostic metabolite for nandrolone. If the urinary concentration of 19-norandrosterone exceeds 2 ng/mL, an AAF is recorded (Baume et al., 2004). The addition of as little as 2.5 μ g of 19-norandrostenedione to a supplement can result in a urinary concentration of 19-norandrosterone that exceeds this

threshold (Watson et al., 2009). These amounts are close to the limits of detection of the analytical methods currently applied to the analysis of dietary supplements, and are far below the levels of contamination deemed acceptable from a health and safety perspective.

Various efforts are being made to address the problems, including the use of third-party auditing activities to identify products that athletes may consider to be at “low risk” of containing prohibited substances. There can be no absolute guarantee that any




Figure 2 — Flow chart to guide informed decision making and reducing risk of ADRV during nutritional supplement use.

Downloaded by on 10/04/18, Volume {article.issue.volume}, Article Number {article.issue.issue}

product is entirely safe, but these schemes do help the athlete to manage the risk. Athletes contemplating the use of dietary supplements should consider very carefully whether the possible benefits outweigh the risks of a doping offense that might end their career.

Practical Implications and Decision Tree

Dietary supplements are an established part of the landscape of modern sport and are likely to remain so. Athletes who take




Figure 3 — Flow chart to guide informed decision making and reducing risk of ADRV during ergogenic supplement use.

Downloaded by on 10/04/18, Volume {article.issue.volume}, Article Number {article.issue.issue}

supplements often have no clear understanding of the potential effects of supplements they are using, but supplements should be used only after a careful cost-benefit analysis has been conducted. On one side of the decision tree are the rewards, the most obvious of which are correction of nutrient deficiencies, achievement of nutritional goals, or enhancement of one or another physiological/biochemical function to directly or indirectly improve performance. On the other side lie the costs—the possibility of using an ineffective supplement, the possible risks to health, and the potential for an ADRV. A flow of questions that could be posed in reaching an informed decision is shown in Figures 2 and 3.

In deciding whether to use a supplement, athletes should consider all aspects of their maturation in, and preparation for, their event to ensure that the supplement under consideration provides an advantage that no other strategy can address. Whether the supplement is practical to use should also be assessed: Is the product available, affordable, tolerated, and compatible with the athlete's other goals? The input of the athlete's coaching team and medical/science support network is important. Athletes who do not have regular access to such a network should consider decisions around supplement use as an important reason to consult an independent sports nutrition expert as well as a physician. Analysis of the evidence around the effectiveness of supplements and their safety is often difficult. A complete nutritional assessment may provide an appropriate justification for the specific use of nutritional supplements and sports foods. For a small number of sports supplements, there is good evidence of a performance effect or indirect benefit for some athletes in some specific situations with little or no risk of adverse outcomes (Peeling et al., 2018; Rawson et al., 2018). Professional advice is often important in ensuring that the athlete is sufficiently knowledgeable about the appropriate protocol for use of these supplements, but individual athletes may respond very differently to a given supplement, with some exhibiting a markedly beneficial effect while others experience no benefit or even a negative effect on performance. Furthermore, the situation in which the athlete wishes to use the supplement may differ in important ways from its substantiated use. Repeated trials may be necessary to establish whether a true effect, rather than just random variation, is seen in response to use of any novel intervention. Some trial and error may also be involved in fine tuning the supplement protocol to suit the needs of the specific situation of use or the individual athlete.

Evidence to support the effectiveness and safety of many of the supplements targeted at athletes, however, is largely absent. There seems to be little incentive for those selling supplements to invest the substantial sums needed to undertake detailed scientific evaluation of their products. Even where some evidence does exist, it may not be relevant to the high-performance athletes because of limitations in the study design (such as the specificity of the exercise tests), the study population, or the context of use. Failure to verify the composition of the supplements used may also give misleading results. It seems sensible to exercise caution when using supplements, as any compound that has the potential to enhance health or exercise performance by altering physiological function must also have the potential for adverse effects in some individuals. Athletes should see good evidence of a performance or other benefit, and should be confident that it will not be harmful to health, before accepting the financial cost and the health or performance risks associated with any supplement. Finally, the athlete should be sure, if supplements or sports foods are to be used, that they have undertaken due diligence to source products that are at low risk of containing prohibited substances.

Conclusion

Dietary supplements can play a small role in an athlete's sports nutrition plan, with products that include essential micronutrients, sports foods, performance supplements, and health supplements all potentially providing benefits. Some supplements, when used appropriately, may help athletes to meet sports nutrition goals, train hard, and stay healthy and injury free. A few supplements can directly enhance competition performance. However, it takes considerable effort and expert knowledge to identify which products are appropriate, how to integrate them into the athlete's sports nutrition plan, and how to ensure that any benefits outweigh the possible negative side effects, including the potential for an ADRV. A strict risk-benefit analysis involving a decision tree approach to the effectiveness, safety, and risks should identify the small number of products that may benefit the athlete. Such an analysis requires the input of a well-informed sports nutrition professional.

References

- Aguiar, A.F., Grala, A.P., da Silva, R.A., Soares-Caldeira, L.F., Pacagnelli, F.L., Ribeiro, A.S., . . . Balvedi, M.C.W. (2017). Free leucine supplementation during an 8-week resistance training program does not increase muscle mass and strength in untrained young adult subjects. *Amino Acids*, 49(7), 1255–1262. PubMed doi:10.1007/s00726-017-2427-0
- Baguet, A., Bourgois, J., Vanhee, L., Achten, E., & Derave, W. (2010). Important role of muscle carnosine in rowing performance. *Journal of Applied Physiology* (1985), 109(4), 1096–1101. doi:10.1152/jappphysiol.00141.2010
- Bailey, R.L., Gahche, J.J., Lentino, C.V., Dwyer, J.T., Engel, J.S., Thomas, P.R., . . . Picciano, M.F. (2011). Dietary supplement use in the United States, 2003–2006. *The Journal of Nutrition*, 141(2), 261–266. doi:10.3945/jn.110.133025
- Bailey, S.J., Fulford, J., Vanhatalo, A., Winyard, P.G., Blackwell, J.R., DiMenna, F.J., . . . Jones, A.M. (2010). Dietary nitrate supplementation enhances muscle contractile efficiency during knee-extensor exercise in humans. *Journal of Applied Physiology* (1985), 109(1), 135–148. doi:10.1152/jappphysiol.00046.2010
- Bailey, S.J., Varnham, R.L., DiMenna, F.J., Breese, B.C., Wylie, L.J., & Jones, A.M. (2015). Inorganic nitrate supplementation improves muscle oxygenation, O₂ uptake kinetics, and exercise tolerance at high but not low pedal rates. *Journal of Applied Physiology* (1985), 118(11), 1396–1405. doi:10.1152/jappphysiol.01141.2014
- Bailey, S.J., Winyard, P., Vanhatalo, A., Blackwell, J.R., DiMenna, F.J., Wilkerson, D.P., . . . Jones, A.M. (2009). Dietary nitrate supplementation reduces the O₂ cost of low-intensity exercise and enhances tolerance to high-intensity exercise in humans. *Journal of Applied Physiology* (1985), 107(4), 1144–1155. doi:10.1152/jappphysiol.00722.2009
- Barrett, E.C., McBurney, M.I., & Ciappio, E.D. (2014). Omega-3 fatty acid supplementation as a potential therapeutic aid for the recovery from mild traumatic brain injury/concussion. *Advances in Nutrition*, 5(3), 268–277. doi:10.3945/an.113.005280
- Baume, N., Avois, L., Schweizer, C., Cardis, C., Dvorak, J., Cauderay, M., . . . Saugy, M. (2004). [13C]Nandrolone excretion in trained athletes: Interindividual variability in metabolism. *Clinical Chemistry*, 50(2), 355–364. doi:10.1373/clinchem.2003.022848
- Bell, P.G., McHugh, M.P., Stevenson, E., & Howatson, G. (2014). The role of cherries in exercise and health. *Scandinavian Journal of Medicine & Sciences in Sports*, 24(3), 477–490. doi:10.1111/sms.12085

- Bellinger, P.M. (2014). beta-Alanine supplementation for athletic performance: An update. *Journal of Strength & Conditioning Research*, 28(6), 1751–1770. doi:10.1519/JSC.0000000000000327
- Bermon, S., Castell, L.M., Calder, P.C., Bishop, N.C., Blomstrand, E., Mooren, F.C., . . . Nagatomi, R. (2017). Consensus statement immunonutrition and exercise. *Exercise Immunology Review*, 23, 8–50. PubMed
- Bodkin, H. (2017). Students left fighting for lives after taking enough caffeine for 300 cups of coffee in botched university experiment. *The Telegraph*. Retrieved from <http://www.telegraph.co.uk/news/2017/01/25/university-fined-400k-students-taking-part-caffeine-experiment/>
- Branch, J.D. (2003). Effect of creatine supplementation on body composition and performance: A meta-analysis. *International Journal of Sport Nutrition and Exercise Metabolism*, 13(2), 198–226.
- Brinkworth, G.D., & Buckley, J.D. (2003). Concentrated bovine colostrum protein supplementation reduces the incidence of self-reported symptoms of upper respiratory tract infection in adult males. *European Journal of Nutrition*, 42(4), 228–232. doi:10.1007/s00394-003-0410-x
- Bruce, C.R., Anderson, M.E., Fraser, S.F., Stepto, N.K., Klein, R., Hopkins, W.G., & Hawley, J.A. (2000). Enhancement of 2000-m rowing performance after caffeine ingestion. *Medicine & Science in Sports & Exercise*, 32(11), 1958–1963.
- Buford, T.W., Kreider, R.B., Stout, J.R., Greenwood, M., Campbell, B., Spano, M., . . . Antonio, J. (2007). International Society of Sports Nutrition position stand: Creatine supplementation and exercise. *Journal of the International Society of Sports Nutrition*, 4, 6. PubMed doi:10.1186/1550-2783-4-6
- Burke, L.M. (2008). Caffeine and sports performance. *Applied Physiology, Nutrition, and Metabolism*, 33(6), 1319–1334. doi:10.1139/H08-130
- Burke, L.M. (2013). Practical considerations for bicarbonate loading and sports performance. *Nestle Nutrition Institute Workshop Series*, 75, 15–26. doi:10.1159/000345814
- Burke, L.M., & Peeling, P. (2018). Methodologies for investigating performance changes with supplement use. *International Journal of Sport Nutrition and Exercise Metabolism*, 28(2). doi:10.1123/ijnsnem.2017-0325
- Carr, A.J., Gore, C.J., & Dawson, B. (2011a). Induced alkalosis and caffeine supplementation: Effects on 2,000-m rowing performance. *International Journal of Sport Nutrition and Exercise Metabolism*, 21(5), 357–364.
- Carr, A.J., Hopkins, W.G., & Gore, C.J. (2011b). Effects of acute alkalosis and acidosis on performance: A meta-analysis. *Sports Medicine*, 41. doi:10.2165/11591440-000000000-00000
- Carr, A.J., Slater, G.J., Gore, C.J., Dawson, B., & Burke, L.M. (2011c). Effect of sodium bicarbonate on [HCO₃⁻], pH, and gastrointestinal symptoms. *International Journal of Sport Nutrition and Exercise Metabolism*, 21(3), 189–194.
- Castell, L.M., Poortmans, J.R., & Newsholme, E.A. (1996). Does glutamine have a role in reducing infections in athletes? *European Journal of Applied Physiology*, 73(5), 488–490.
- Cermak, N.M., Res, P.T., de Groot, L.C., Saris, W.H., & van Loon, L.J. (2012). Protein supplementation augments the adaptive response of skeletal muscle to resistance-type exercise training: A meta-analysis. *The American Journal of Clinical Nutrition*, 96(6), 1454–1464. PubMed
- Chung, W., Shaw, G., Anderson, M.E., Pyne, D.B., Saunders, P.U., Bishop, D.J., & Burke, L.M. (2012). Effect of 10 week beta-alanine supplementation on competition and training performance in elite swimmers. *Nutrients*, 4(10), 1441–1453. PubMed doi:10.3390/nu4101441
- Clark, A., & Mach, N. (2017). The crosstalk between the gut microbiota and mitochondria during exercise. *Frontiers in Physiology*, 8, 319. doi:10.3389/fphys.2017.00319
- Clark, K.L., Sebastianelli, W., Flechsenhar, K.R., Aukermann, D.F., Meza, F., Millard, R.L., . . . Albert, A. (2008). 24-Week study on the use of collagen hydrolysate as a dietary supplement in athletes with activity-related joint pain. *Current Medical Research and Opinion*, 24(5), 1485–1496. PubMed doi:10.1185/030079908X291967
- Close, G.L., Hamilton, D.L., Philp, A., Burke, L.M., & Morton, J.P. (2016). New strategies in sport nutrition to increase exercise performance. *Free Radical Biology & Medicine*, 98, 144–158. doi:10.1016/j.freeradbiomed.2016.01.016
- Close, G.L., Russell, J., Cobley, J.N., Owens, D.J., Wilson, G., Gregson, W., . . . Morton, J.P. (2013). Assessment of vitamin D concentration in non-supplemented professional athletes and healthy adults during the winter months in the UK: Implications for skeletal muscle function. *Journal of Sports Sciences*, 31(4), 344–353. doi:10.1080/02640414.2012.733822
- Coelho Rabello Lima, L., Oliveira Assumpcao, C., Prestes, J., & Sergio Denadai, B. (2015). Consumption of cherries as a strategy to attenuate exercise-induced muscle damage and inflammation in humans. *Nutricion Hospitalaria*, 32(5), 1885–1893. PubMed doi:10.3305/nh.2015.32.5.9709
- Cook, C.J., Crewther, B.T., Kilduff, L.P., Drawer, S., & Gaviglio, C.M. (2011). Skill execution and sleep deprivation: Effects of acute caffeine or creatine supplementation - a randomized placebo-controlled trial. *Journal of the International Society of Sports Nutrition*, 8, 2. PubMed doi:10.1186/1550-2783-8-2
- Cooper, R., Naclerio, F., Allgrove, J., & Jimenez, A. (2012). Creatine supplementation with specific view to exercise/sports performance: An update. *Journal of the International Society of Sports Nutrition*, 9(1), 33. PubMed doi:10.1186/1550-2783-9-33
- Davison, G., & Diment, B.C. (2010). Bovine colostrum supplementation attenuates the decrease of salivary lysozyme and enhances the recovery of neutrophil function after prolonged exercise. *British Journal of Nutrition*, 103(10), 1425–1432. PubMed doi:10.1017/S0007114509993503
- Deminice, R., Rosa, F.T., Franco, G.S., Jordao, A.A., & de Freitas, E.C. (2013). Effects of creatine supplementation on oxidative stress and inflammatory markers after repeated-sprint exercise in humans. *Nutrition*, 29(9), 1127–1132. PubMed doi:10.1016/j.nut.2013.03.003
- Deutz, N.E., Pereira, S.L., Hays, N.P., Oliver, J.S., Edens, N.K., Evans, C.M., & Wolfe, R.R. (2013). Effect of beta-hydroxy-beta-methylbutyrate (HMB) on lean body mass during 10 days of bed rest in older adults. *Clinical Nutrition*, 32(5), 704–712. doi:10.1016/j.clnu.2013.02.011
- Douroudos, I.I., Fatouros, I.G., Gourgoulis, V., Jamurtas, A.Z., Tsitsios, T., & Hatzinikolaou, A. (2006). Dose-related effects of prolonged NaHCO₃ ingestion during high-intensity exercise. *Medicine & Science in Sports & Exercise*, 38(10), 1746–1753. doi:10.1249/01.mss.0000230210.60957.67
- Dulson, D.K., & Bishop, N.C. (2016). Effect of a high and low dose of caffeine on human lymphocyte activation in response to antigen stimulation. *Applied Physiology, Nutrition, and Metabolism*, 41(2), 224–227. doi:10.1139/apnm-2015-0456
- Erdman, J., Oria, M., & Pillsbury, L. (2011). *Nutrition and traumatic brain injury: Improving acute and subacute health outcomes in military personnel*. Washington, DC: National Academies Press.
- Fennell, D. (2004). Determinants of supplement usage. *Preventive Medicine*, 39(5), 932–939. PubMed doi:10.1016/j.ypmed.2004.03.031
- French, C., McNaughton, L., Davies, P., & Tristram, S. (1991). Caffeine ingestion during exercise to exhaustion in elite distance runners. Revision. *The Journal of Sports Medicine and Physical Fitness*, 31(3), 425–432.

- Ganio, M.S., Klau, J.F., Casa, D.J., Armstrong, L.E., & Maresh, C.M. (2009). Effect of caffeine on sport-specific endurance performance: A systematic review. *The Journal of Strength & Conditioning Research*, 23(1), 315–324. [PubMed](#) doi:10.1519/JSC.0b013e31818b979a
- Garthe, I., & Maughan, R.J. (2018). Athletes and supplements: Prevalence and perspectives. *International Journal of Sport Nutrition and Exercise Metabolism*, 28(2). doi:10.1123/ijnsnem.2017-0429
- Geller, A.I., Shehab, N., Weidle, N.J., Lovegrove, M.C., Wolpert, B.J., Timbo, B.B., . . . Budnitz, D.S. (2015). Emergency department visits for adverse events related to dietary supplements. *The New England Journal of Medicine*, 373(16), 1531–1540. [PubMed](#) doi:10.1056/NEJMSa1504267
- Geyer, H., Bredehoff, M., Marek, U., Parr, M.K., & Schanzer, W. (2002). Hohe Dosen des Anabolikums Metandienon in Nahrungsergänzungsmitteln. *Deutsche Apotheke Zeitung*, 142, 29.
- Geyer, H., Parr, M.K., Koehler, K., Mareck, U., Schanzer, W., & Thevis, M. (2008). Nutritional supplements cross-contaminated and faked with doping substances. *Journal of Mass Spectrometry*, 43(7), 892–902. [PubMed](#) doi:10.1002/jms.1452
- Geyer, H., Parr, M.K., Mareck, U., Reinhart, U., Schrader, Y., & Schanzer, W. (2004). Analysis of non-hormonal nutritional supplements for anabolic-androgenic steroids - results of an international study. *International Journal of Sports Medicine*, 25(2), 124–129. [PubMed](#) doi:10.1055/s-2004-819955
- Gibson, R.S. (2005). *Principles of nutritional assessment* (2nd ed.). New York, NY: Oxford University Press.
- Gleeson, M. (2016). Immunological aspects of sport nutrition. *Immunology and Cell Biology*, 94, 117–123. [PubMed](#) doi:10.1038/icb.2015.109
- Gleeson, M., Bishop, N.C., Oliveira, M., & Tauler, P. (2011). Daily probiotic's (*Lactobacillus casei* Shirota) reduction of infection incidence in athletes. *International Journal of Sport Nutrition and Exercise Metabolism*, 21(1), 55–64.
- Goodman, L.S., & Gilman, A. (1975). *Pharmacological basis of therapeutics* (5th ed., p. 1462). London/Toronto: Macmillan Publishing Co.
- Gray, P., Chappell, A., Jenkinson, A.M., Thies, F., & Gray, S.R. (2014). Fish oil supplementation reduces markers of oxidative stress but not muscle soreness after eccentric exercise. *International Journal of Sport Nutrition and Exercise Metabolism*, 24(2), 206–214. doi:10.1123/ijnsnem.2013-0081
- Gualano, B., Rawson, E.S., Candow, D.G., & Chilibeck, P.D. (2016). Creatine supplementation in the aging population: Effects on skeletal muscle, bone and brain. *Amino Acids*, 48(8), 1793–1805. [PubMed](#) doi:10.1007/s00726-016-2239-7
- Gualano, B., Roschel, H., Lancha, A.H., Jr., Brightbill, C.E., & Rawson, E.S. (2012). In sickness and in health: The widespread application of creatine supplementation. *Amino Acids*, 43(2), 519–529. [PubMed](#) doi:10.1007/s00726-011-1132-7
- Hao, Q., Dong, B.R., & Wu, T. (2015). Probiotics for preventing acute upper respiratory tract infections. *The Cochrane Database of Systematic Reviews*, (2), CD006895. doi:10.1002/14651858.CD006895.pub3
- Harris, R.C., Söderlund, K., & Hultman, E. (1992). Elevation of creatine in resting and exercised muscle of normal subjects by creatine supplementation. *Clinical Sciences (Lond)*, 83(3), 367–374.
- He, C.S., Aw Yong, X.H., Walsh, N.P., & Gleeson, M. (2016). Is there an optimal vitamin D status for immunity in athletes and military personnel? *Exercise Immunology Review*, 22, 42–64.
- Heaney, R.P. (2008). Vitamin D: Criteria for safety and efficacy. *Nutrition Reviews*, 66(10 Suppl. 2), S178–S181. [PubMed](#) doi:10.1111/j.1753-4887.2008.00102.x
- Heaton, L.E., Davis, J.K., Rawson, E.S., Nuccio, R.P., Witard, O.C., Halson, S.L., . . . Carter, J.M. (2017). In season nutritional strategies to enhance recovery for team sport athletes. *Sports Medicine*, 47(11), 2201–2218. [PubMed](#)
- Hector, A.J., & Philips, S.M. (2018). Protein recommendations for weight loss in elite athletes: A focus on body composition and performance. *International Journal of Sport Nutrition and Exercise Metabolism*, 28(2). doi:10.1123/ijnsnem.2017-0273
- Hemila, H., & Chalker, E. (2013). Vitamin C for preventing and treating the common cold. *The Cochrane Database of Systematic Reviews*, (1), CD000980. doi:10.1002/14651858.CD000980.pub4
- Hemila, H., Virtamo, J., Albanes, D., & Kaprio, J. (2003). Physical activity and the common cold in men administered vitamin E and beta-carotene. *Medicine & Science in Sports & Exercise*, 35(11), 1815–1820. [PubMed](#) doi:10.1249/01.MSS.0000093616.60899.92
- Hobson, R.M., Saunders, B., Ball, G., Harris, R.C., & Sale, C. (2012). Effects of beta-alanine supplementation on exercise performance: A meta-analysis. *Amino Acids*, 43(1), 25–37. [PubMed](#) doi:10.1007/s00726-011-1200-z
- Hoon, M.W., Jones, A.M., Johnson, N.A., Blackwell, J.R., Broad, E.M., Lundy, B., . . . Burke, L.M. (2014). The effect of variable doses of inorganic nitrate-rich beetroot juice on simulated 2,000-m rowing performance in trained athletes. *International Journal of Sports Physiology and Performance*, 9(4), 615–620. [PubMed](#) doi:10.1123/ijsp.2013-0207
- Hossein-Nezhad, A., & Holick, M.F. (2013). Vitamin D for health: A global perspective. *Mayo Clinic Proceedings*, 88(7), 720–755. [PubMed](#) doi:10.1016/j.mayocp.2013.05.011
- Hultman, E., Soderlund, K., Timmons, J.A., Cederblad, G., & Greenhaff, P.L. (1996). Muscle creatine loading in men. *Journal of Applied Physiology* (1985), 81(1), 232–237.
- Jakeman, J.R., Lambrick, D.M., Wooley, B., Babraj, J.A., & Faulkner, J.A. (2017). Effect of an acute dose of omega-3 fish oil following exercise-induced muscle damage. *European Journal of Applied Physiology*, 117(3), 575–582. doi:10.1007/s00421-017-3543-y
- Jones, A.M. (2014a). Dietary nitrate supplementation and exercise performance. *Sports Medicine*, 44(Suppl. 1), S35–S45.
- Jones, A.M. (2014b). Influence of dietary nitrate on the physiological determinants of exercise performance: A critical review. *Applied Physiology, Nutrition, and Metabolism*, 39(9), 1019–1028.
- Jouris, K.B., McDaniel, J.L., & Weiss, E.P. (2011). The effect of omega-3 fatty acid supplementation on the inflammatory response to eccentric strength exercise. *Journal of Sports Sciences Medicine*, 10(3), 432–438.
- Jull, A.B., Ni Mhurchu, C., Bennett, D.A., Dunshea-Mooij, C.A., & Rodgers, A. (2008). Chitosan for overweight or obesity. *The Cochrane Database of Systematic Reviews*, (3), CD003892. doi:10.1002/14651858.CD003892.pub3
- Jurgens, T.M., Whelan, A.M., Killian, L., Doucette, S., Kirk, S., & Foy, E. (2012). Green tea for weight loss and weight maintenance in overweight or obese adults. *The Cochrane Database of Systematic Reviews*, (12), CD008650. doi:10.1002/14651858.CD008650.pub2
- Karsch-Volk, M., Barrett, B., & Linde, K. (2015). Echinacea for preventing and treating the common cold. *JAMA*, 313(6), 618–619. [PubMed](#) doi:10.1001/jama.2014.17145
- Katz, A., Costill, D.L., King, D.S., Hargreaves, M., & Fink, W.J. (1984). Maximal exercise tolerance after induced alkalosis. *International Journal of Sports Medicine*, 5(2), 107–110. [PubMed](#) doi:10.1055/s-2008-1025890
- Kreider, R.B., Kalman, D.S., Antonio, J., Ziegenfuss, T.N., Wildman, R., Collins, R., . . . Lopez, H.L. (2017). International Society of Sports Nutrition position stand: Safety and efficacy of creatine

- supplementation in exercise, sport, and medicine. *Journal of the International Society of Sports Nutrition*, 14, 18. PubMed doi:10.1186/s12970-017-0173-z
- Krieger, J.W., Sitren, H.S., Daniels, M.J., & Langkamp-Henken, B. (2006). Effects of variation in protein and carbohydrate intake on body mass and composition during energy restriction: A meta-regression. *American Journal of Clinical Nutrition*, 83(2), 260–274. PubMed
- Kucukgoncu, S., Zhou, E., Lucas, K.B., & Tek, C. (2017). Alpha-lipoic acid (ALA) as a supplementation for weight loss: Results from a meta-analysis of randomized controlled trials. *Obesity Reviews*, 18(5), 594–601. PubMed doi:10.1111/obr.12528
- Lambert, C.P., Greenhaff, P.L., Ball, D., & Maughan, R.J. (1993). Influence of sodium bicarbonate ingestion on plasma ammonia accumulation during incremental exercise in man. *European Journal of Applied Physiology and Occupational Physiology*, 66(1), 49–54. PubMed
- Lancha Junior, A.H., Painelli Vde, S., Saunders, B., & Artioli, G.G. (2015). Nutritional strategies to modulate intracellular and extracellular buffering capacity during high-intensity exercise. *Sports Medicine*, 45(Suppl. 1), S71–S81. PubMed doi:10.1007/s40279-015-0397-5
- Lanthers, C., Pereira, B., Naughton, G., Trousselard, M., Lesage, F.X., & Dutheil, F. (2017). Creatine supplementation and upper limb strength performance: A systematic review and meta-analysis. *Sports Medicine*, 47(1), 163–173. PubMed doi:10.1007/s40279-016-0571-4
- Lappe, J., Cullen, D., Haynatzki, G., Recker, R., Ahlf, R., & Thompson, K. (2008). Calcium and vitamin D supplementation decreases incidence of stress fractures in female navy recruits. *Journal of Bone and Mineral Research*, 23(5), 741–749. doi:10.1359/jbmr.080102
- Larson-Meyer, D.E., Woolf, K., & Burke, L.M., (2018). Assessment of nutrient status in athletes and the need for supplementation. *International Journal of Sport Nutrition and Exercise Metabolism*, 28(2). doi:10.1123/ijnsnem.2017-0338
- Larson-Meyer, D.E., & Willis, K.S. (2010). Vitamin D and athletes. *Current Sports Medicine Reports*, 9(4), 220–226. PubMed
- Lewis, M., Ghassemi, P., & Hibbeln, J. (2013). Therapeutic use of omega-3 fatty acids in severe head trauma. *American Journal of Emergency Medicine*, 31(1), 273.e5–273.e8. doi:10.1016/j.ajem.2012.05.014
- LGC. (2017). Australian supplement survey summary. Retrieved from <http://www2.lgcgroup.com/australiansupplementsurvey>
- Li, N., Hauser, R., Holford, T., Zhu, Y., Zhang, Y., Bassig, B.A., . . . Zheng, T. (2015). Muscle-building supplement use and increased risk of testicular germ cell cancer in men from Connecticut and Massachusetts. *British Journal of Cancer*, 112(7), 1247–1250. PubMed doi:10.1038/bjc.2015.26
- Linde, K., Barrett, B., Wolkart, K., Bauer, R., & Melchart, D. (2006). Echinacea for preventing and treating the common cold. *The Cochrane Database of Systematic Reviews*, (1), CD000530. doi:10.1002/14651858.CD000530.pub2
- Lowery, R.P., Joy, J.M., Rathmacher, J.A., Baier, S.M., Fuller, J.C., Jr., Shelley, M.C., . . . Wilson, J.M. (2016). Interaction of beta-hydroxy-beta-methylbutyrate free acid and adenosine triphosphate on muscle mass, strength, and power in resistance trained individuals. *The Journal of Strength & Conditioning Research*, 30(7), 1843–1854. PubMed
- Maganaris, C.N., & Maughan, R.J. (1998). Creatine supplementation enhances maximum voluntary isometric force and endurance capacity in resistance trained men. *Acta Physiologica Scandinavica*, 163(3), 279–287. PubMed doi:10.1046/j.1365-201x.1998.00395.x
- Mainwood, G.W., & Worsley-Brown, P. (1975). The effects of extracellular pH and buffer concentration on the efflux of lactate from frog sartorius muscle. *The Journal of Physiology*, 250(1), 1–22. PubMed
- Martinez-Sanz, J.M., Sospedra, I., Ortiz, C.M., Baladia, E., Gil-Izquierdo, A., & Ortiz-Moncada, R. (2017). Intended or unintended doping? A review of the presence of doping substances in dietary supplements used in sports. *Nutrients*, 9(10). doi:10.3390/nu9101093
- Maughan, R.J. (2005). Contamination of dietary supplements and positive drug tests in sport. *Journal of Sports Sciences*, 23(9), 883–889. doi:10.1080/02640410400023258
- Maughan, R.J., Depiesse, F., & Geyer, H. (2007). The use of dietary supplements by athletes. *Journal of Sports Sciences*, 25(Suppl. 1), S103–S113. doi:10.1080/02640410701607395
- Maughan, R.J., & Griffin, J. (2003). Caffeine ingestion and fluid balance: A review. *Journal of Human Nutrition and Dietetics*, 16(6), 411–420. PubMed
- Maughan, R.J., Shirreffs, S.M., & Verrec, A. (2018). Making decisions about supplement use. *International Journal of Sport Nutrition and Exercise Metabolism*, 28(2). doi:10.1123/ijnsnem.2018-0009
- McAlindon, T.E., Nuite, M., Krishnan, N., Ruthazer, R., Price, L.L., Burstein, D., . . . Flechsenhar, K. (2011). Change in knee osteoarthritis cartilage detected by delayed gadolinium enhanced magnetic resonance imaging following treatment with collagen hydrolysate: A pilot randomized controlled trial. *Osteoarthritis Cartilage*, 19(4), 399–405. PubMed doi:10.1016/j.joca.2011.01.001
- McFarlin, B.K., Venable, A.S., Henning, A.L., Sampson, J.N., Pennel, K., Vingren, J.L., & Hill, D.W. (2016). Reduced inflammatory and muscle damage biomarkers following oral supplementation with bioavailable curcumin. *BBA Clinical*, 5, 72–78. PubMed doi:10.1016/j.bbacli.2016.02.003
- McMahon, N.F., Leveritt, M.D., & Pavey, T.G. (2016). The effect of dietary nitrate supplementation on endurance exercise performance in healthy adults: A systematic review and meta-analysis. *Sports Medicine*, 47(4), 735–756. doi:10.1007/s40279-016-0617-7
- Mc Naughton, L., & Thompson, D. (2001). Acute versus chronic sodium bicarbonate ingestion and anaerobic work and power output. *The Journal of Sports Medicine and Physical Fitness*, 41(4), 456–462. PubMed
- Mettler, S., & Zimmermann, M.B. (2010). Iron excess in recreational marathon runners. *European Journal of Clinical Nutrition*, 64(5), 490–494. PubMed doi:10.1038/ejcn.2010.16
- Meydani, S.N., Han, S.N., & Hamer, D.H. (2004). Vitamin E and respiratory infection in the elderly. *Annals of the New York Academy of Sciences*, 1031, 214–222. PubMed doi:10.1196/annals.1331.021
- Mickleborough, T.D. (2013). Omega-3 polyunsaturated fatty acids in physical performance optimization. *International Journal of Sport Nutrition and Exercise Metabolism*, 23(1), 83–96. PubMed
- Morton, R.W., Murphy, K.T., McKellar, S.R., Schoenfeld, B.J., Henselmans, M., Helms, E., . . . Phillips, S.M. (2017). A systematic review, meta-analysis and meta-regression of the effect of protein supplementation on resistance training-induced gains in muscle mass and strength in healthy adults. *British Journal of Sports Medicine*. doi:10.1136/bjsports-2017-097608
- Nassis, G.P., Sporer, B., & Stathis, C.G. (2016). Beta-alanine efficacy for sports performance improvement: From science to practice. *British Journal of Sports Medicine*. doi:10.1136/bjsports-2016-097038
- Nicol, L.M., Rowlands, D.S., Fazakerly, R., & Kelllett, J. (2015). Curcumin supplementation likely attenuates delayed onset muscle soreness (DOMS). *European Journal of Applied Physiology*, 115(8), 1769–1777. doi:10.1007/s00421-015-3152-6
- Nieman, D.C., Henson, D.A., Gross, S.J., Jenkins, D.P., Davis, J.M., Murphy, E.A., . . . Mayer, E.P. (2007). Quercetin reduces illness but not immune perturbations after intensive exercise.

- Medicine & Science in Sports & Exercise*, 39(9), 1561–1569. doi:10.1249/mss.0b013e318076b566
- Nieman, D.C., Henson, D.A., McAnulty, S.R., McAnulty, L., Swick, N.S., Utter, A.C., . . . Morrow, J.D. (2002). Influence of vitamin C supplementation on oxidative and immune changes after an ultramarathon. *Journal of Applied Physiology*, 92(5), 1970–1977. doi:10.1152/jappphysiol.00961.2001
- Nieman, D.C., Henson, D.A., McMahon, M., Wrieden, J.L., Davis, J.M., Murphy, E.A., . . . Dumke, C.L. (2008). Beta-glucan, immune function, and upper respiratory tract infections in athletes. *Medicine & Science in Sports & Exercise*, 40(8), 1463–1471. doi:10.1249/MSS.0b013e31817057c2
- Nikolaidis, M.G., Kerksick, C.M., Lamprecht, M., & McAnulty, S.R. (2012). Does vitamin C and E supplementation impair the favorable adaptations of regular exercise? *Oxidative Medicine and Cellular Longevity*, 2012, 707941. PubMed doi:10.1155/2012/707941
- Office of Dietary Supplements, National Institutes of Health. (1994). Dietary supplement health and education act of 1994. Retrieved from https://ods.od.nih.gov/About/DSHEA_Wording.aspx
- Onakpoya, I., Hunt, K., Wider, B., & Ernst, E. (2014a). Pyruvate supplementation for weight loss: A systematic review and meta-analysis of randomized clinical trials. *Critical Reviews in Food Science and Nutrition*, 54(1), 17–23. doi:10.1080/10408398.2011.565890
- Onakpoya, I., Posadzki, P., & Ernst, E. (2014b). The efficacy of glucomannan supplementation in overweight and obesity: A systematic review and meta-analysis of randomized clinical trials. *The Journal of the American College of Nutrition*, 33(1), 70–78. doi:10.1080/07315724.2014.870013
- Onakpoya, I.J., Posadzki, P.P., Watson, L.K., Davies, L.A., & Ernst, E. (2012). The efficacy of long-term conjugated linoleic acid (CLA) supplementation on body composition in overweight and obese individuals: A systematic review and meta-analysis of randomized clinical trials. *European Journal of Nutrition*, 51(2), 127–134. doi:10.1007/s00394-011-0253-9
- Owens, D.J., Sharples, A.P., Polydorou, I., Alwan, N., Donovan, T., Tang, J., . . . Close, G.L. (2015). A systems-based investigation into vitamin D and skeletal muscle repair, regeneration, and hypertrophy. *American Journal of Physiology-Endocrinology and Metabolism*, 309(12), E1019–E1031. PubMed doi:10.1152/ajpendo.00375.2015
- Owens, D.J., Webber, D., Impey, S.G., Tang, J., Donovan, T.F., Fraser, W.D., . . . Close, G.L. (2014). Vitamin D supplementation does not improve human skeletal muscle contractile properties in insufficient young males. *European Journal of Applied Physiology*, 114(6), 1309–1320. doi:10.1007/s00421-014-2865-2
- Paton, C., Costa, V., & Guglielmo, L. (2015). Effects of caffeine chewing gum on race performance and physiology in male and female cyclists. *Journal of Sports Sciences*, 33(10), 1076–1083. doi:10.1080/02640414.2014.984752
- Paulsen, G., Cumming, K.T., Holden, G., Hallen, J., Ronnestad, B.R., Sveen, O., . . . Raastad, T. (2014). Vitamin C and E supplementation hampers cellular adaptation to endurance training in humans: A double-blind, randomised, controlled trial. *Journal of Physiology*, 592(8), 1887–1901. doi:10.1113/jphysiol.2013.267419
- Peake, J.M., Neubauer, O., Walsh, N.P., & Simpson, R.J. (2017). Recovery of the immune system after exercise. *Journal of Applied Physiology* (1985), 122(5), 1077–1087. doi:10.1152/jappphysiol.00622.2016
- Peeling, P., Binnie, M., Goods, P.S.R., Sim, M., & Burke, L.M. (2018). Evidence-based supplements for the enhancement of athletic performance. *International Journal of Sport Nutrition and Exercise Metabolism*, 28(2). doi:10.1123/ijnsnem.2017-0343
- Phillips, S.M., Aragon, A.A., Arciero, P.J., Arent, S.M., Close, G.L., Hamilton, D.L., . . . Willoughby, D.S. (2017). Changes in body composition and performance with supplemental HMB-FA + ATP. *Journal of Strength & Conditioning Research*, 31(5):e71–e72. doi:10.1519/jsc.0000000000001760
- Powers, M.E., Arnold, B.L., Weltman, A.L., Perrin, D.H., Mistry, D., Kahler, D.M., . . . Volek, J. (2003). Creatine supplementation increases total body water without altering fluid distribution. *Journal of Athletic Training*, 38(1), 44–50. PubMed
- Powers, S.K., Nelson, W.B., & Larson-Meyer, E. (2011). Antioxidant and Vitamin D supplements for athletes: sense or nonsense? *Journal of Sports Sciences*, 29(Suppl.1), S47–S55. doi:10.1080/02640414.2011.602098
- Prasad, A.S. (2014). Zinc is an antioxidant and anti-inflammatory agent: Its role in human health. *Frontiers in Nutrition*, 1, 14. doi:10.3389/fnut.2014.00014
- Rae, C.D., & Broer, S. (2015). Creatine as a booster for human brain function. How might it work? *Neurochemistry International*, 89, 249–259. doi:10.1016/j.neuint.2015.08.010
- Rawson, E.S., Clarkson, P.M., & Tarnopolsky, M.A. (2017). Perspectives on Exertional Rhabdomyolysis. *Sports Medicine*, 47(Suppl.1), 33–49. doi:10.1007/s40279-017-0689-z
- Rawson, E.S., Miles, M.P., & Larson-Meyer, D.E. (2018). Dietary supplements for health, adaptation, and recovery in athletes. *International Journal of Sport Nutrition and Exercise Metabolism*, 28(2). doi:10.1123/ijnsnem.2017-0340
- Rawson, E.S., & Persky, A.M. (2007). Mechanisms of muscular adaptations to creatine supplementation: Review article. *International SportMed Journal*, 8(2), 43–53.
- Rawson, E.S., & Venezia, A.C. (2011). Use of creatine in the elderly and evidence for effects on cognitive function in young and old. *Amino Acids*, 40(5), 1349–1362. PubMed
- Rawson, E.S., & Volek, J.S. (2003). Effects of creatine supplementation and resistance training on muscle strength and weightlifting performance. *Journal of Strength & Conditioning Research*, 17(4), 822–831.
- Requena, B., Zabala, M., Padial, P., & Ferliche, B. (2005). Sodium bicarbonate and sodium citrate: Ergogenic aids? *Journal of Strength & Conditioning Research*, 19(1), 213–224. doi:10.1519/13733.1
- Reynolds, C., Halpenny, C., Hughes, C., Jordan, S., Quinn, A., & Egan, B. (2016). Acute ingestion of beetroot juice does not improve repeated sprint performance in male team sport athletes. *Proceedings of the Nutrition Society*, 75(OCE3), E97. doi:10.1017/S0029665116001129
- Ribeiro, I.F., Miranda-Vilela, A.L., Klautau-Guimaraes Mde, N., & Grisolia, C.K. (2013). The influence of erythropoietin (EPO T → G) and alpha-actinin-3 (ACTN3 R577X) polymorphisms on runners' responses to the dietary ingestion of antioxidant supplementation based on pequi oil (Caryocar brasiliense Camb.): A before-after study. *Journal of Nutrigenetics and Nutrigenomics*, 6(6), 283–304. doi:10.1159/000357947
- Roberts, L., Bailes, J., Dedhia, H., Zikos, A., Singh, A., McDowell, D., . . . Carpenter, J. (2008). Surviving a mine explosion. *Journal of the American College of Surgeons*, 207(2), 276–283. doi:10.1016/j.jamcollsurg.2008.02.015
- Rowlands, D.S., & Thomson, J.S. (2009). Effects of beta-hydroxy-beta-methylbutyrate supplementation during resistance training on strength, body composition, and muscle damage in trained and untrained young men: A meta-analysis. *Journal of Strength & Conditioning Research*, 23(3), 836–846.
- Ruohola, J.P., Laaksi, I., Ylikomi, T., Haataja, R., Mattila, V.M., Sahi, T., . . . Pihlajamaki, H. (2006). Association between serum 25(OH)D

- concentrations and bone stress fractures in Finnish young men. *Journal of Bone and Mineral Research*, 21(9), 1483–1488. doi:10.1359/jbmr.060607
- Sakellaris, G., Kotsiou, M., Tamiolaki, M., Kalostos, G., Tsapaki, E., Spanaki, M., . . . Evangelidou, A. (2006). Prevention of complications related to traumatic brain injury in children and adolescents with creatine administration: An open label randomized pilot study. *The Journal of Trauma*, 61(2), 322–329. doi:10.1097/01.ta.0000230269.46108.d5
- Sakellaris, G., Nasis, G., Kotsiou, M., Tamiolaki, M., Charissis, G., & Evangelidou, A. (2008). Prevention of traumatic headache, dizziness and fatigue with creatine administration. A pilot study. *Acta Paediatrica*, 97(1), 31–34. PubMed doi:10.1111/j.1651-2227.2007.00529.x
- Saunders, B., Elliott-Sale, K., Artioli, G.G., Swinton, P.A., Dolan, E., Roschel, H., . . . Gualano, B. (2016). Beta-alanine supplementation to improve exercise capacity and performance: A systematic review and meta-analysis. *British Journal of Sports Medicine*. doi:10.1136/bjsports-2016-096396
- Schilling, B.K., Stone, M.H., Utter, A., Kearney, J.T., Johnson, M., Coglianese, R., . . . Stone, M.E. (2001). Creatine supplementation and health variables: A retrospective study. *Medicine & Science in Sports & Exercise*, 33(2), 183–188.
- Schneiker, K.T., Bishop, D., Dawson, B., & Hackett, L.P. (2006). Effects of caffeine on prolonged intermittent-sprint ability in team-sport athletes. *Medicine & Science in Sports & Exercise*, 38(3), 578–585. doi:10.1249/01.mss.0000188449.18968.62
- Schrier, S.L., & Auerbach, M. (2017). Treatment of iron deficiency anemia in adults. Retrieved from <https://www.uptodate.com/contents/treatment-of-iron-deficiency-anemia-in-adults>
- Sciberras, J.N., Galloway, S.D., Fenech, A., Grech, G., Farrugia, C., Duca, D., & Mifsud, J. (2015). The effect of turmeric (Curcumin) supplementation on cytokine and inflammatory marker responses following 2 hours of endurance cycling. *Journal of the International Society of Sports Nutrition*, 12(1), 5. doi:10.1186/s12970-014-0066-3
- Shaw, G., Lee-Barthel, A., Ross, M.L., Wang, B., & Baar, K. (2017). Vitamin C-enriched gelatin supplementation before intermittent activity augments collagen synthesis. *The American Journal of Clinical Nutrition*, 105(1), 136–143. doi:10.3945/ajcn.116.138594
- Siegler, J.C., Marshall, P.W.M., Bray, J., & Towlson, C. (2012). Sodium bicarbonate supplementation and ingestion timing: Does it matter? *Journal of Strength & Conditioning Research*, 26, 1953–1958. doi:10.1519/JSC.0b013e3182392960
- Singh, M., & Das, R.R. (2013). Zinc for the common cold. *The Cochrane Database of Systematic Reviews*, (6), CD001364. doi:10.1002/14651858.CD001364.pub4
- Smith, G.I., Atherton, P., Reeds, D.N., Mohammed, B.S., Rankin, D., Rennie, M.J., & Mittendorfer, B. (2011a). Dietary omega-3 fatty acid supplementation increases the rate of muscle protein synthesis in older adults: A randomized controlled trial. *The American Journal of Clinical Nutrition*, 93(2), 402–412. doi:10.3945/ajcn.110.005611
- Smith, G.I., Atherton, P., Reeds, D.N., Mohammed, B.S., Rankin, D., Rennie, M.J., & Mittendorfer, B. (2011b). Omega-3 polyunsaturated fatty acids augment the muscle protein anabolic response to hyperinsulinaemia-hyperaminoacidaemia in healthy young and middle-aged men and women. *Clinical Science (Lond)*, 121(6), 267–278.
- Solimini, R., Rotolo, M.C., Mastrobattista, L., Mortali, C., Minutillo, A., Pichini, S., . . . Palmi, I. (2017). Hepatotoxicity associated with illicit use of anabolic androgenic steroids in doping. *European Review for Medical and Pharmacological Sciences*, 21(Suppl. 1), 7–16. PubMed
- Solomons, N.W. (1998). Mild human zinc deficiency produces an imbalance between cell-mediated and humoral immunity. *Nutrition Reviews*, 56(1 Pt 1), 27–28.
- Spriet, L.L. (2014). Exercise and sport performance with low doses of caffeine. *Sports Medicine*, 44(Suppl. 2), S175–S184. doi:10.1007/s40279-014-0257-8
- Steenge, G.R., Simpson, E.J., & Greenhaff, P.L. (2000). Protein- and carbohydrate-induced augmentation of whole body creatine retention in humans. *Journal of Applied Physiology* (1985), 89(3), 1165–1171.
- Stellingwerff, T., & Cox, G.R. (2014). Systematic review: Carbohydrate supplementation on exercise performance or capacity of varying durations. *Applied Physiology, Nutrition, and Metabolism*, 39(9), 998–1011. doi:10.1139/apnm-2014-0027
- Sullivan, P.G., Geiger, J.D., Mattson, M.P., & Scheff, S.W. (2000). Dietary supplement creatine protects against traumatic brain injury. *Annals of Neurology*, 48(5), 723–729. PubMed
- Szczesniak, K.A., Ostaszewski, P., Fuller, J.C., Jr., Ciecierska, A., & Sadkowski, T. (2015). Dietary supplementation of beta-hydroxy-beta-methylbutyrate in animals – a review. *Journal of Animal Physiology and Animal Nutrition (Berl)*, 99(3), 405–417. doi:10.1111/jpn.12234
- Talanian, J.L., & Spriet, L.L. (2016). Low and moderate doses of caffeine late in exercise improve performance in trained cyclists. *Applied Physiology, Nutrition, and Metabolism*, 41(8), 850–855. doi:10.1139/apnm-2016-0053
- Thomas, D.T., Erdman, K.A., & Burke, L.M. (2016). American College of Sports Medicine Joint Position Statement. Nutrition and Athletic Performance. *Medicine & Science in Sports & Exercise*, 48(3), 543–568. doi:10.1249/MSS.0000000000000852
- Thompson, C., Vanhatalo, A., Jell, H., Fulford, J., Carter, J., Nyman, L., . . . Jones, A.M. (2016). Dietary nitrate supplementation improves sprint and high-intensity intermittent running performance. *Nitric Oxide*, 61, 55–61. PubMed doi:10.1016/j.niox.2016.10.006
- Thompson, C., Wylie, L.J., Fulford, J., Kelly, J., Black, M.I., McDonagh, S.T., . . . Jones, A.M. (2015). Dietary nitrate improves sprint performance and cognitive function during prolonged intermittent exercise. *European Journal of Applied Physiology*, 115(9), 1825–1834. doi:10.1007/s00421-015-3166-0
- Tian, H., Guo, X., Wang, X., He, Z., Sun, R., Ge, S., & Zhang, Z. (2013). Chromium picolinate supplementation for overweight or obese adults. *The Cochrane Database of Systematic Reviews*, (11), CD010063. doi:10.1002/14651858.CD010063.pub2
- Tipton, K.D. (2015). Nutritional support for exercise-induced injuries. *Sports Medicine*, 45(Suppl. 1), S93–S104. doi:10.1007/s40279-015-0398-4
- Volek, J.S., & Rawson, E.S. (2004). Scientific basis and practical aspects of creatine supplementation for athletes. *Nutrition*, 20(7–8), 609–614. PubMed doi:10.1016/j.nut.2004.04.014
- Volman, J.J., Ramakers, J.D., & Plat, J. (2008). Dietary modulation of immune function by beta-glucans. *Physiology & Behavior*, 94(2), 276–284. doi:10.1016/j.physbeh.2007.11.045
- Walker, G.J., Finlay, O., Griffiths, H., Sylvester, J., Williams, M., & Bishop, N.C. (2007). Immunoendocrine response to cycling following ingestion of caffeine and carbohydrate. *Medicine & Science in Sports & Exercise*, 39(9), 1554–1560. doi:10.1249/mss.0b013e3180a74228
- Walsh, N.P., Blannin, A.K., Robson, P.J., & Gleeson, M. (1998). Glutamine, exercise and immune function. Links and possible mechanisms. *Sports Medicine*, 26(3), 177–191. PubMed
- Walsh, N.P., Gleeson, M., Shephard, R.J., Gleeson, M., Woods, J.A., Bishop, N.C., . . . Simon, P. (2011). Position statement. Part one: Immune function and exercise. *Exercise Immunology Review*, 17, 6–63.

- Watson, P., Judkins, C., Houghton, E., Russell, C., & Maughan, R.J. (2009). Urinary nandrolone metabolite detection after ingestion of a nandrolone precursor. *Medicine & Science in Sports & Exercise*, *41*(4), 766–772. doi:[10.1249/MSS.0b013e31818edaeb](https://doi.org/10.1249/MSS.0b013e31818edaeb)
- Wellington, B.M., Leveritt, M.D., & Kelly, V.G. (2017). The effect of caffeine on repeat high intensity effort performance in rugby league players. *International Journal of Sports Physiology and Performance*, *12*(2), 206–210. doi:[10.1123/ijsp.2015-0689](https://doi.org/10.1123/ijsp.2015-0689)
- Wiles, J.D., Coleman, D., Tegerdine, M., & Swaine, I.L. (2006). The effects of caffeine ingestion on performance time, speed and power during a laboratory-based 1 km cycling time-trial. *Journal of Sports Sciences*, *24*(11), 1165–1171. doi:[10.1080/02640410500457687](https://doi.org/10.1080/02640410500457687)
- Wilkinson, D.J., Hossain, T., Hill, D.S., Phillips, B.E., Crossland, H., Williams, J., . . . Atherton, P.J. (2013). Effects of leucine and its metabolite beta-hydroxy-beta-methylbutyrate on human skeletal muscle protein metabolism. *The Journal of Physiology*, *591*(Pt 11), 2911–2923.
- Wilson, J.M., Lowery, R.P., Joy, J.M., Andersen, J.C., Wilson, S.M., Stout, J.R., . . . Rathmacher, J. (2014). The effects of 12 weeks of beta-hydroxy-beta-methylbutyrate free acid supplementation on muscle mass, strength, and power in resistance-trained individuals: A randomized, double-blind, placebo-controlled study. *European Journal of Applied Physiology*, *114*(6), 1217–1227.
- Wilson, J.M., Lowery, R.P., Joy, J.M., Walters, J.A., Baier, S.M., Fuller, J.C., Jr., . . . Rathmacher, J. (2013). Beta-Hydroxy-beta-methylbutyrate free acid reduces markers of exercise-induced muscle damage and improves recovery in resistance-trained men. *The British Journal of Nutrition*, *110*(3), 538–544.
- Wycherley, T.P., Moran, L.J., Clifton, P.M., Noakes, M., & Brinkworth, G.D. (2012). Effects of energy-restricted high-protein, low-fat compared with standard-protein, low-fat diets: A meta-analysis of randomized controlled trials. *The American Journal of Clinical Nutrition*, *96*(6), 1281–1298.
- Wylie, L.J., Bailey, S.J., Kelly, J., Blackwell, J.R., Vanhatalo, A., & Jones, A.M. (2016). Influence of beetroot juice supplementation on intermittent exercise performance. *European Journal of Applied Physiology*, *116*(2), 415–425. doi:[10.1007/s00421-015-3296-4](https://doi.org/10.1007/s00421-015-3296-4)
- Wylie, L.J., Kelly, J., Bailey, S.J., Blackwell, J.R., Skiba, P.F., Winyard, P.G., . . . Jones, A.M. (2013). Beetroot juice and exercise: Pharmacodynamic and dose-response relationships. *Journal of Applied Physiology* (1985), *115*(3), 325–336. doi:[10.1152/jappphysiol.00372.2013](https://doi.org/10.1152/jappphysiol.00372.2013)
- Zhang, Y.Y., Liu, W., Zhao, T.Y., & Tian, H.M. (2017). Efficacy of omega-3 polyunsaturated fatty acids supplementation in managing overweight and obesity: A meta-analysis of randomized clinical trials. *The Journal of Nutrition, Health & Aging*, *21*(2), 187–192. doi:[10.1007/s12603-016-0755-5](https://doi.org/10.1007/s12603-016-0755-5)