



Fungicide Efficacy for Control of Wheat Diseases

The **North Central Regional Committee on Management of Small Grain Diseases (NCERA-184)** has developed the following information about fungicide efficacy for the control of certain foliar diseases of wheat for use by the grain production industry in the United States.

The efficacy ratings for each fungicide listed in the table were determined by field testing the materials over multiple years and locations by the members of the committee. Efficacy is based on proper application timing to achieve optimum effectiveness of the fungicide as determined by labeled instructions and overall level of disease in the field at the time of application. Differences in efficacy among fungicide products were determined by direct comparisons among products in field tests and are based on a single application of the labeled rate as listed in the table.

The table includes most widely marketed products, and is not intended to be a list of all labeled products.

Many products have specific use restrictions. Restrictions may be present on the amount of active ingredient that can be applied within a period of time or on the number of sequential applications that can occur.

Read and follow all use restrictions before applying any fungicide.

Efficacy of fungicides for wheat disease control based on appropriate application timing

Class	Fungicide(s)		Rate/A (fl. oz)	Powdery Mildew	Stagonospora Leaf/Glume Blotch	Septoria Leaf Blotch	Tan Spot	Stripe Rust	Leaf Rust	Stem Rust	Head Scab ⁴	Harvest Restriction
	Active Ingredient	Product										
Strobilurins	picoxystrobin 22.5%	Aproach SC [®]	6.0-12.0	G ¹	VG	VG ²	VG	E ³	VG	VG	NL	Feekes 10.5
	pyraclostrobin 23.6%	Headline SC [®]	6.0-9.0	G	VG	VG ²	E	E ³	E	G	NL	Feekes 10.5
Triazoles	metconazole 8.6%	Caramba 0.75SL [®]	10.0-17.0	VG	VG	--	VG	E	E	E	G	30 days
	tebuconazole 38.7%	Folicur 3.6F ^{®5}	4.0	NL	NL	NL	NL	E	E	E	F	30 days
	prothioconazole 41%	Proline 480SC [®]	5.0-5.7	--	VG	VG	VG	VG	VG	VG	G	30 days
	prothioconazole 19%	Prostaro 4215C [®]	6.5-8.2	G	VG	VG	VG	E	E	E	G	30 days
	tebuconazole 19%											
	propiconazole 41.8%	Tilt 3.6EC ^{®5}	4.0	VG	VG	VG	VG	VG	VG	VG	P	Feekes 10.5.4
	tebuconazole 22.6%	Absolute Maxx SC [®]	5.0	G	VG	VG	VG	VG	E	VG	NL	35 days
	trifloxystrobin 22.6%											
	cyproconazole 7.17%	Aproach Prima SC [®]	3.4-6.8	VG	VG	VG	VG	E	VG	VG	--	45 days
	picoxystrobin 17.94%											
Mixed modes of action ⁶	prothioconazole 16.0%	Delaro 325SC [®]	8.0	G	VG	VG	VG	VG	VG	VG	NL	Feekes 10.5 35 days
	trifloxystrobin 13.7%											
	pydiflumetofen 13.7%	Miravis Ace SE [®]	13.7	VG	VG	VG	VG	VG	VG	VG	G ⁷	Feekes 10.5.4
	propiconazole 11.4%											
	flupyroxad 2.8%	Nexicor EC [®]	7.0-13.0	VG	VG	E	E	E	E	VG	NL	Feekes 10.5
	pyraclostrobin 18.7%											
	propiconazole 11.7%	Preemptor SC [®]	4.0-6.0	--	--	VG	VG	E	VG	--	NL	Feekes 10.5 and 40 days
	fluoxastrobin 14.8%											
	flutriafol 19.3%	Priaxor [®]	4.0-8.0	G	VG	VG	E	VG	VG	G	NL	Feekes 10.5
	fluxapyroxad 14.3%											
	pyraclostrobin 28.6%	Quilt Xcel 2.2SE ^{®5}	10.5-14.0	VG	VG	VG	VG	E	E	VG	NL	Feekes 10.5.4
	propiconazole 11.7%											
azoxystrobin 13.5%	Stratego YLD [®]	4.0	G	VG	VG	VG	VG	VG	VG	NL	Feekes 10.5 35 days	
prothioconazole 10.8%												
trifloxystrobin 32.3%	Trivapro SE [®]	9.4-13.7	VG	VG	VG	VG	E	E	VG	NL	Feekes 10.5.4	
benzovindiflupyr 2.9%												
propiconazole 11.9%												
azoxystrobin 10.5%	Topguard EQ [®]	4.0-7.0	VG	NL	VG	VG	E	E	VG	NL	Feekes 10.5.4 30 days	
flutriafol 18.63%												
azoxystrobin 25.30%												

¹ Efficacy categories: NL=Not Labeled; NR=Not Recommended; P=Poor; F=Fair; G=Good; VG=Very Good; E=Excellent; --=Insufficient data to make statement about efficacy of this product.
² Product efficacy may be reduced in areas with fungal populations that are resistant to strobilurin fungicides.
³ Efficacy may be significantly reduced if solo strobilurin products are applied after stripe rust infection has occurred.
⁴ Application of products containing strobilurin fungicides may result in elevated levels of the mycotoxin Deoxynivalenol (DON) in grain damaged by head scab.
⁵ Multiple generic products containing the same active ingredients also may be labeled in some states.
⁶ Products with mixed modes of action generally combine triazole and strobilurin active ingredients. Miravis Ace[®], Nexicor[®], Priaxor[®], and Trivapro[®] include carboxamide active ingredients.
⁷ Based on application timing at the beginning of anthesis (Feekes 10.5.1).

Find Out More

The Crop Protection Network (CPN) is a multi-state and international collaboration of university and provincial extension specialists, and public and private professionals who provide unbiased, research-based information to farmers and agricultural personnel. Our goal is to communicate relevant information that will help professionals identify and manage field crop diseases.

Find more resources at CropProtectionNetwork.org.



We Are Extension



This publication was developed and published by Erick DeWolf (Kansas State University) and members of the NCERA-184.

The information in this publication is only a guide, and the authors assume no liability for practices implemented based on this information. Reference to products in this publication is not intended to be an endorsement to the exclusion of others that may be similar. Individuals using such products assume responsibility for their use in accordance with current directions of the manufacturer.

May 2020

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or a part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotope, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, S.W., Washington, D.C. 20250-9410 or call (800) 795-3272 (voice) or (202) 720-6382 (TDD). USDA is an equal opportunity provider and employer.

©2020 by the Crop Protection Network. All rights reserved.