

Simulation design matters

Citation for published version (APA):

Pinto de Melo, B. C. (2018). Simulation design matters: improving obstetrics training outcomes. Maastricht: Maastricht University. <https://doi.org/10.26481/dis.20181212bm>

Document status and date:

Published: 01/01/2018

DOI:

[10.26481/dis.20181212bm](https://doi.org/10.26481/dis.20181212bm)

Document Version:

Publisher's PDF, also known as Version of record

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.



Simulation Design Matters

Improving Obstetrics Training Outcomes

Brena Carvalho Pinto de Melo

The research reported here was carried out at:



in the School of Health Professions Education:



To Juliano e Bernardo

ISBN: 978-94-6323-438-2

The copyright of articles that have been published has been transferred to the respective journals.

The research reported here was conducted at the School of Health Professions Education, Maastricht University, Maastricht, Netherlands, at the Instituto de Medicina Integral Prof. Fernando Figueira (IMIP), Recife, Brasil and Faculdade Pernambucana de Saúde (FPS), Recife, Brasil.

Copyright © Brena Melo, Recife

Layout: Carla Teixeira

Cover design: Breno Carvalho

Printed by: Gildeprint – www.gildeprint.nl

Simulation Design Matters

Improving Obstetrics Training Outcomes

DISSERTATION

to obtain the degree of Doctor at Maastricht University,
on the authority of the Rector Magnificus,
Prof. dr. Rianne M. Letschert
in accordance with the decision of the Board of Deans,
to be defended in public
on Wednesday 12th December 2018, at 12:45 hours

by

Brena Carvalho Pinto de Melo

Supervisors:

Prof. dr. J.J.G. van Merriënboer
Prof. dr. C.P.M. van der Vleuten

Co-Supervisors:

Prof. dr. A.R. Falbo, IMIP, Brasil

Assessment Committee:

Prof. dr. J.J. Rethans (chair)
Prof. dr. W.N.K.A. van Mook
Dr. A.G. van der Niet
Prof. dr. F. Scheele, VU University Amsterdam
Dr. R. Tio, Catharina Ziekenhuis Eindhoven

Table of Contents

Chapter 1	9
Introduction	
Chapter 2	21
The limited use of instructional design guidelines in healthcare simulation scenarios: an expert appraisal	
Chapter 3	43
The use of instructional design guidelines to increase effectiveness of postpartum hemorrhage simulation training	
Chapter 4	59
Self-perceived long-term transfer of learning after postpartum hemorrhage simulation training	
Chapter 5	79
Effects of an in situ instructional design based postpartum hemorrhage simulation training on patient outcomes: an uncontrolled before-and-after study	
Chapter 6	99
General Discussion	
Summary	114
Samenvatting	120
Valorization	126
Acknowledgment	132
Curriculum Vitae	138
SHE Dissertations Series	140





Chapter 1

Introduction

Introduction

Healthcare simulation training has widely spread throughout international medical schools in recent decades.¹ The concurrent financial investment it demands is supported by the legitimate argument of improving patient outcome. Evidence, however, on the effectiveness of this type of training format is still being sought.²

Why simulate?

The concept of simulation training was originally imported into the healthcare field from the war and airway industries, mainly to present itself as an alternative to an “error safe environment”.³ Its dissemination was propelled by both the goals of increasing patient safety and providing an alternative to supervised practice opportunity to residents after the adoption of duty hour’s restriction by residency programs.

Patient safety issues received great attention from the public and scientific communities after the American National Institute of Medicine (NIH) report, “To err is human”, in 1999. According to this report, each year, 98,000 people died in US hospitals as a result of preventable medical errors with recent updates on this analysis presenting even more worrisome results with estimations at least twice as large.^{4,5}

Residents’ duty restriction hours’ policy was another reason for great public and scientific commotion and was a consequence of the Libby Zion case. In 1984, Libby Zion, an 18-year-old girl, was admitted in to the emergency room in Boston, USA, with fever and earache. She was pronounced dead after 6 hours. She had been examined and taken care of by two overworked young doctors. Further, a senior clinician was available only over the phone for supervision. Her father, a New York Times journalist, requested an investigation into her death and a grand jury investigation lead to no criminal charges, but unveiled the strenuous workload of residents in training. As a result, this investigation led to a series of recommendations on limiting residents’ working hours, which were adopted by the regulating boards. Critiques to this

policy, however, argued that a relevant contributing problem to the case and to other potentially preventable adverse outcomes was lack of supervision.⁶ Another downside argument to residents' duty restriction hours was the consequent diminished opportunity to practice, in particular for rare events.^{7,8} Simulation training arose as the promising solution to all these problems: patient safety, practice restriction and a lack of supervision.

After the initial boom of this type of training format, a concern regarding its effectiveness began to emerge, particularly due to the evolving technological and budget demands.⁹ Systematic reviews analysing both the impact of technology and instructional features on simulation trainings described large study inconsistencies and poor description of instructional features use. As a conclusion, the reviewers recommend adopting instructional design guidelines when devising simulation trainings.^{2,10}

Instructional design (ID) guidelines were developed based on sound cognitive psychological theories on how people learn.¹¹ They have a unique role in helping learners reach complex learning. Complex learning is the integrated acquisition of knowledge, skills and attitudes along with the ability to properly coordinate the qualitatively different constituent skills.¹² Moreover, ID guidelines aim at optimizing long-term knowledge transferability by presenting authentic, relevant, every day real-world problems, while simultaneously emphasizing practice at different levels of complexity and providing feedback and diminished instructional support through training.¹³ Two examples of models providing instructional design guidelines are Merrill's First Principles of Instruction¹⁴ and the 4C/ID model.¹⁵ According to Merrill, learning is better promoted when the following five principles are present: (1) learners are engaged in solving real life problems, (2) their previous knowledge is activated as a foundation to new knowledge, (3) the new knowledge is demonstrated to them, (4) they have the opportunity to apply the new knowledge, and (5) they have the opportunity to integrate it into their world. On its turn, the 4C/ID model claims that four interrelated components are essential for complex learning: (1) learning tasks, (2) supportive information, (3) procedural information, and (4) part-task practice.^{12,16} Healthcare personnel are expected to achieve complex learning, particularly when managing high-

risk situations with their demanding simultaneous, multidisciplinary, and multitasking tasks.^{16,17}

An example of a healthcare high-risk situation is postpartum hemorrhage (PPH). PPH is commonly defined as a blood loss of 500ml or more within 24 hours after birth. It is the leading cause of maternal mortality worldwide with most related deaths occurring within these 24 hours.¹⁸⁻²⁰ Concerns were raised due to a recent documented rise in its frequency in several high-income countries, along with an increase in US maternal mortality rates.²¹ Furthermore, it is frequently occurring, unpredictable and can happen to either high-risk women or women with no prior disease.²² Additionally, to reinforce the reason for such concern, PPH related deaths are described as preventable in some studies in up to 100% of cases, and attributable to management flaws.²²⁻²⁶ Simulation training has, therefore, been advocated as the healthcare personnel training strategy to strike this scenery.^{18,20,24,26}

The path in reaching optimal simulation training

Demonstrating simulation training effectiveness became of paramount relevance with such high stakes involved: improving patient safety, providing training opportunities for both novices and experienced personnel– either for daily recurrent tasks or rare high-risk events – and enabling for a supervised “error safe” environment. In spite of the expanding number of publications in the field, many of these concerned replication studies, the so-called “me too” studies, and comparison studies between simulation and no intervention.²⁷ Therefore, sound evidence on the effectiveness of this type of training format is still needed.²⁸

When considering training effectiveness, particularly for a complex environment such as healthcare, different domains must be taken into account: the learners’ context, work environment related elements and/or patient outcome. According to the Kirkpatrick training evaluation model, there are four different levels: learners’ satisfaction, learning outcomes, behavior, and patient outcomes, which may be considered stepping-stones for training evaluation.²⁹ Critiques of this model underscore it as an oversimplified model, which does not acknowledge moderating variables, such as trainee’s motivation and work environment and that failure to reach transfer would be solely attributable to training design.³⁰⁻³²

These shortcomings, however, may be compensated by adopting complementary models, such as the training transfer Baldwin and Ford model, when assessing training evaluation, particularly for the behavior/training transfer stage.³³ Moreover, additional studies regarding simulation training effectiveness should also aim to further explore related patient outcomes with detailed description. For instance, they should provide data beyond procedural outcomes, but also report on survival, duration of hospitalization, complications and preferably describe be designed to have patient outcomes as the study primary outcome. A systematic review on patient outcomes in simulation-based medical education, however, highlights these topics as poorly described, reporting large studies' inconsistencies, high prevalence of statistical errors, procedural-solely described and lack of prior definition of primary outcomes.³⁴ When addressing the need to further promote simulation training translational outcome, the literature again advises exploring the use of instructional design guidelines.³⁵

The use of instructional design guidelines for simulation training may be explored from a multitude of perspectives, as exemplified in the following four: use of ID features per se, cognitive impact, influence on work environment and behavior, and the ultimate patient outcome impact.^{2,34,36} First, on the use of instructional features per se, it may be worth exploring how they are currently used for specific contents either individually, on the amount of exposure, as well as its intensity. For instance, the debate on a proper definition of authenticity for the simulation training field includes a discussion about simulator fidelity and simulation training environments (simulation center or in situ).³⁷⁻³⁹ Second, exploring the cognitive impact of the use of each of the instructional features or their combination in training complex tasks to promote complex learning seems as the long-demanded advance in the frontier of knowledge on simulation training educational research.⁴⁰ Third, preliminary data begins to emerge concerning training transfer for simulation training, but still mostly on short term transfer.^{41,42} Finally, the complex field of simulation training patient outcome starts to present its first results but still with the methodological limitations previously cited.⁴³

The recent epidemiological commotion about PPH alarming mortality rates and its specific clinical management characteristics sets it up as an ideal

content for researching simulation training from an educational perspective. Its optimal clinical management requires complex learning to be reassured by the execution of simultaneous tasks from a multitude of different domains such as: communication, teamwork, situational awareness, systematic approach to problem solving, while providing opportunity to supervised practice of either recurrent and non-recurrent skills.

Bridging the simulation training educational research beyond

In summary, improving patient safety by providing supervised practice opportunity in a safe environment is complex and concerns multiple variables. In addition, one of its basic premises is ensuring effective healthcare personnel training with long-term transferability. The use of instructional design guidelines in simulation training may contribute in reaching this goal. Postpartum hemorrhage due to its omnipresence, unpredictability, and the multidisciplinary care it requires allows multiple learning dimensions to be studied.⁴⁴

We therefore aim to explore the use of instructional design guidelines for postpartum hemorrhage simulation training.

We intend to reach this aim by answering the following questions:

- 1)** How are instructional design guidelines described and used in the current postpartum hemorrhage healthcare simulation literature?
- 2)** What are the different learning outcomes when two instructionally different postpartum hemorrhage simulation training programmes are compared, one based on instructional design guidelines and the other on current best practices?
- 3)** Is there long-term transfer after these two different training formats? Is there a difference in learning outcomes according to the attended training format? What instructional design features may have facilitated and/or hindered long-term transfer?
- 4)** What is the impact on patient outcome at a teaching hospital after Obstetrics and Gynecology residents attending an instructional design based postpartum hemorrhage simulation training programme format?

Structure of the thesis

Our first study, presented in Chapter 2, provides an overview of the currently adopted Instructional Design Guidelines in published articles at which PPH simulation trainings were described. Invited raters (N=40) analyzed 32 articles with regards to: Were they applied to the trainings designs? If so, how? Were the used ID features described in details at the analyzed articles' methods section to allow for reproducibility? A rating scale was elaborated for the detailed analysis. It was based on Merrill's First Principles of Instructions and assessed training description elements such as: authenticity, activation of prior knowledge, demonstration, application and integration.

The study presented on Chapter 3 compared the learning outcomes of two instructionally different PPH simulation training: one based on instructional design guidelines and the other on a prior instructional "best practice". In order to select this to-be-compared "best PPH training" seven experts (five educationalists and two from a healthcare training background) rated previously identified articles from the literature with regards to their instructional perspective. For the quasi-randomized comparison, 20 teams were distributed accordingly: 13 teams attended the ID based training and seven, the BP based. Learning outcomes were compared with regards to the following expected-to-be-executed tasks subscales: communication, teamwork, vital signs, laboratory evaluation, drug management, mechanical management and surgical management.

On Chapter 4, we present the residents perception on the long-term transfer after attending either of the training formats. We analyzed semi-structured interviews, using thematic analyzed. We explored: (1) Whether the residents perceived training transfer on the long term, (2) if there was a difference in long-term training transfer perception according to the attended format and (3) whether they reported any influence of specific instructional design features on their long-term training transfer perception.

The PPH simulation training patient outcome study is described at chapter 5. Data on clinical outcome of women delivering at a high risk teaching maternity was collected at Recife, Brazil, for a six months period in total. To avoid potential seasonality bias, data was collected for the same

three months period one year before the PPH simulation training and for the same three months immediately after the training. During training, active management of the third stage of labor (PPH prophylaxis) was part of the content. Primary outcomes were therapeutic uterotonics use rates within 24h of birth (TUW24h) and blood transfusion rates (BTR). Secondary outcome measures included uterotonic use rates and dosages for each drug, postpartum hemoglobin <6g/dL, B-Lynch suture, uterine artery embolization, hysterectomy, renal insufficiency, obstetric intensive care unit (ICU) admission, mechanical ventilation, length of hospital stay, near-miss maternal mortality and maternal death.

Chapter 6 presents a general discussion on the findings of the previously described studies. The strengths and limitations are discussed along with the implications for practice and for research in simulation training with regards to the use of ID guidelines. The discussion chapter is followed by a summary in English and Dutch.


References

1. Gaba DM. The future vision of simulation in health care. *Qual Saf Health Care* 2004;13:i2:i10.
2. Cook DA, Hamstra SJ, Brydges R, et al. Comparative effectiveness of instructional design features in simulation-based education: systematic review and meta-analysis. *Med Teach* 2013;35:e867-98.
3. Rosen KR. The history of medical simulation. *J Crit Care* 2008;23:157-66.
4. Kohn LT, Corrigan JM, Donaldson MS. *To err is human: building a safer health system*. United States of America: Library of Congress Cataloging-in-Publication Data; 2000.
5. James JT. A new, evidence-based estimate of patient harms associated with hospital care. *J Patient Saf* 2013;9:122-8.
6. Patel N. Learning lessons: the Libby Zion case revisited. *J Am Coll Cardiol* 2014;64:2802-4.
7. Jamal MH, Wong S, Whalen TV. Effects of the reduction of surgical residents' work hours and implications for surgical residency programs: a narrative review. *BMC Med Educ* 2014;14(Suppl 1).
8. Rybock JD. Residents' duty hours and professionalism. *N Engl J Med* 2009;361:930-1.

9. McGaghie WC, Issenberg SB, Petrusa ER, Scalese RJ. A critical review of simulation-based medical education research: 2003-2009. *Med Educ* 2010;44:50-63.
10. Cook DA, Hatala R, Brydges R, et al. Technology-enhanced simulation for health professions education: a systematic review and meta-analysis. *JAMA* 2011;306:978.
11. Paas F, Renkl A, Sweller J. Cognitive load theory and instructional design: recent developments. *Educ Psychol* 2003;38:1-4.
12. van Merriënboer JGG, Kirschner PA. *Ten Steps to Complex Learning*. 3rd ed. New York, NY: Routledge; 2018.
13. van Merriënboer JGG, Kester L, Paas F. Teaching complex rather than simple tasks: Balancing intrinsic and germane load to enhance transfer of learning. *Appl Cogn Psychol* 2006;20:343-52.
14. Merrill MD. *First Principles of Instruction*. *ETR&D* 2002;50:43-59.
15. van Merriënboer JGG, Clark RE, de Croock MBM. Blueprints for complex learning: The 4C/ID-Model. *ETR&D* 2002;50:39-61.
16. Van Merriënboer JGG, Sluijsmans DMA. Toward a synthesis of cognitive load theory, Four-Component Instructional Design, and self-directed learning. *Educ Psychol Rev* 2009;21:55-66.
17. Bradley P. The history of simulation in medical education and possible future directions. *Med Educ* 2006;40:254-62.
18. World Health Organization. WHO recommendations for the prevention and treatment of postpartum haemorrhage. Italy, 2012.
19. GBD 2015 Maternal Mortality Collaborators. Global, regional, and national levels of maternal mortality, 1990–2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet* 2016;388:1775-812.
20. Antony KM, Dildy GA, 3rd. Postpartum hemorrhage: The role of the Maternal-Fetal Medicine specialist in enhancing quality and patient safety. *Semin Perinatol* 2013;37:246-56.
21. Agrawal P. Maternal mortality and morbidity in the United States of America. *Bull World Health Organ* 2013;93:135.
22. Driessen M, Bouvier-Colle MH, Dupont C, et al. Postpartum hemorrhage resulting from uterine atony after vaginal delivery: factors associated with severity. *Obstet Gynecol* 2011;117:21-31.
23. Bohlmann MK, Rath W. Medical prevention and treatment of postpartum hemorrhage: a comparison of different guidelines. *Arch Gynecol Obstet* 2014;289:555-67.

24. Knight M, Callaghan WM, Berg C, et al. Trends in postpartum hemorrhage in high resource countries: a review and recommendations from the International Postpartum Hemorrhage Collaborative Group. *BMC Pregnancy Childbirth* 2009;9:55.
25. Say L, Chou D, Gemmill A, et al. Global causes of maternal death - a WHO systematic analysis. *Lancet* 2014:e323-e33.
26. Wong CA, Scott S, Jones RL, Walzer J, Geller S. The state of Illinois obstetric hemorrhage project: pre-project and post-training examination scores. *J Matern Fetal Neonatal Med* 2016;29:845-9.
27. Cook DA. If you teach them, they will learn: why medical education needs comparative effectiveness research. *Adv Health Sci Educ Theory Pract* 2012;17:305-10.
28. Cook DA. How much evidence does it take? A cumulative meta-analysis of outcomes of simulation-based education. *Med Educ* 2014;48:750-60.
29. Kirkpatrick D. Great ideas revisited. Techniques for evaluating training programs. Revisiting Kirkpatrick's Four-Level Model. *Train Dev J* 1996;50:54-9.
30. Holton III EF. The flawed Four-Level evaluation model. *Hum Resour Dev Q* 1996;7,5-21.
31. Holton III EF. Holton's Evaluation Model: new evidence and construct elaborations. *Adv Dev Hum Resour* 2005;7:37-54.
32. Yardley S, Dornan T. Kirkpatrick's levels and education 'evidence'. *Med Educ* 2012;46:97-106.
33. Baldwin TT, Ford JK. Transfer of training: A review and directions for future research. *Pers Psychol* 1988;41:63-105.
34. Zendejas B, Brydges R, Wang AT, Cook DA. Patient outcomes in simulation-based medical education: A systematic review. *J Gen Intern Med* 2013;28:1078-89.
35. McGaghie WC, Draycott TJ, Dunn WF, Lopez CM, Stefanidis D. Evaluating the impact of simulation on translational patient outcomes. *Simul Healthc* 2011;6 Suppl:S42-7.
36. Issenberg SB, McGaghie WC, Petrusa ER, Lee Gordon D, Scalese RJ. Features and uses of high-fidelity medical simulations that lead to effective learning: a BEME systematic review. *Med Teach* 2005;27:10-28.
37. Norman G, Dore K, Grierson L. The minimal relationship between simulation fidelity and transfer of learning. *Med Educ* 2012;46:636-47.
38. Grierson LE. Information processing, specificity of practice, and the transfer of learning: Considerations for reconsidering fidelity. *Adv Health Sci Educ Theory Pract* 2014;19:281-9.

39. Sorensen JL, Ostergaard D, LeBlanc V, et al. Design of simulation-based medical education and advantages and disadvantages of in situ simulation versus off-site simulation. *BMC Med Educ* 2017;17:20.
40. Haji FA, Cheung JJ, Woods N, Regehr G, de Ribaupierre S, Dubrowski A. Thrive or overload? The effect of task complexity on novices' simulation-based learning. *Med Educ* 2016;50:955-68.
41. Mema B, Harris I. The Barriers and facilitators to transfer of ultrasound-guided central venous line skills from simulation to practice: exploring perceptions of learners and supervisors. *Teach Learn Med* 2016;28:115-24.
42. Evans LV, Dodge KL, Shah TD, et al. Simulation training in central venous catheter insertion: improved performance in clinical practice. *Acad Med* 2010;85:1462-9.
43. Markova V, Sorensen JL, Holm C, Norgaard A, Langhoff-Roos J. Evaluation of multi-professional obstetric skills training for postpartum hemorrhage. *Acta Obstet Gynecol Scand* 2012;91:346-52.
44. Eichbaum Q. Collaboration and teamwork in the health professions: Rethinking the role of conflict. *Acad Med* 2018;93:574-580.



This chapter is embargoed at request

Chapter 2

The limited use of instructional design guidelines in healthcare simulation scenarios: an expert appraisal¹

¹ This chapter is based on a manuscript submitted for publication: De Melo BCP, Falbo AR, Souza E, Muijtjens AMM, van Merriënboer JJG, & van der Vleuten CPM. (submitted). The limited use of instructional design guidelines in healthcare simulation scenarios: an expert appraisal.



Chapter 3

The use of instructional design guidelines to increase effectiveness of postpartum hemorrhage simulation training²

² This chapter is based on the published article: De Melo BCP, Falbo AR, Muijtjens AMM, van der Vleuten CPM, & van Merriënboer, JG. (2017). The use of instructional design guidelines to increase effectiveness of postpartum hemorrhage simulation training. *International Journal of Gynaecology and Obstetrics: The Official Organ of the International Federation of Gynaecology and Obstetrics*, 137(1), 99-105. DOI: 10.1002/ijgo.12084

Abstract

Introduction

Simulation training has become widespread for the past decade. The search for optimal training effectiveness has raised awareness with regards to simulation design. The use of instructional design guidelines has been recommended.

Objective

To compare learning outcomes of postpartum hemorrhage simulation training based on either instructional design guidelines or best practice.

Methods

A pretest–post-test non-equivalent groups study was conducted among obstetrics and gynecology residents in Recife, Brazil, from June 8 to August 30, 2013. The instructional design group included 13 teams, whereas the best practice group included seven teams. A standardized task checklist was used for scenario analysis and the proportion of correctly executed tasks compared (post-test minus pretest).

Results

The instructional design group scored higher than the best practice group for total number of tasks completed (median difference 0.46 vs 0.17; $P < 0.001$; effect size [r]=0.72). Similar results were observed for communication (median difference 0.56 vs 0.22; $P = 0.004$; $r = 0.58$), laboratory evaluation (median difference 0.83 vs 0.00; $P < 0.001$; $r = 0.76$), and mechanical management (median difference 0.25 vs -0.15 ; $P = 0.048$; $r = 0.39$). Speed of learning was also increased. The median differences were 0.20 for the instructional design group compared with 0.05 for the best practice group at 60 seconds ($P = 0.015$; $r = 0.49$), and 0.49 versus 0.26 ($P = 0.001$; $r = 0.65$) at 360 seconds.

Conclusion

The use of simulation training for postpartum hemorrhage that was based on instructional design guidelines yielded better learning outcomes than did training based on best practice.

Introduction

Postpartum hemorrhage (PPH) is the leading cause of maternal mortality worldwide, particularly among deaths that are considered avoidable.¹ Frequently reported assistance failures associated with preventable PPH include delay in diagnosis, poor communication, insufficient teamwork, and lack of adequate education and training.^{2,3} Simulation training has emerged as a widespread strategy to overcome such failures. Systematic reviews of simulation-based healthcare education debate the best way to deliver instructional features to ensure optimal effectiveness of training.^{4,5} Nonetheless, a growing body of evidence recommends that training formats should be based on reliable educational principles, particularly instructional design guidelines.⁶

The Four-Component Instructional Design model⁷ and Merrill's First Principles of Instruction⁸ summarize the main guidelines. According to the guidelines, effective instruction should contain authentic, relevant, and daily real-world problems, emphasize practice at different levels of complexity, and provide feedback and diminishing instructional support through training.⁷

When devising a PPH simulation training format based on instructional design guidelines, the learning objectives should be clearly identified and summarized in strategies that aim to overcome recurrent assistance failures. Such learning objectives include situational awareness enhancement,⁹ improvement of both communication and team-work skills and attitudes,¹⁰ and reinforcement of specific knowledge and skills regarding the correct management of patients with PPH.¹⁻³

The aim of the present study was to test the hypothesis that a PPH simulation training program based on instructional design guidelines would lead to better learning outcomes than the use of a simulation training program based on best practice.

Methods

A pretest–post-test non-equivalent groups study was conducted at two of the local teaching hospitals in Recife, Brazil—Instituto de Medicina Integral Prof. Fernando Figueira and Hospital das Clínicas, Universidade Federal de Pernambuco—from June 8 to August 30, 2013. Obstetrics and gynecology residents working at any of the five local teaching hospitals in Recife were considered eligible and invited to participate. These hospitals provide high-risk maternity services and report an equivalent number of deliveries per resident. The present study was approved by the Ethics Committee of the Instituto de Medicina Integral Prof. Fernando Figueira. All participants provided signed informed consent at the training session.

Of all eligible residents, half worked at the Instituto de Medicina Integral Prof. Fernando Figueira. Therefore, residents from this center formed the entire instructional design group ($n=36$). They were divided into 13 teams: four teams of two individuals and nine teams of three individuals. One participant had to leave the instructional design group for personal reasons before undergoing the post-test. Participants assigned to the best practice group ($n=18$) came from the other four maternity hospitals and underwent training at Hospital das Clínicas, Universidade Federal de Pernambuco. These participants were divided into seven teams: three teams of two individuals and four teams of three individuals. A similar distribution among teams by age, sex, and year of residency was noted for both training formats. Training occurred in half-day sessions, with one or two teams at each.

According to Merrill,⁸ efficient instruction should engage learners in solving real-world problems, activate existing knowledge as a foundation for new knowledge, demonstrate the new knowledge to the learner, provide opportunities for applying the new knowledge, and integrate the new knowledge into the trainee's world. Additionally, the Four-Component Instructional Design model⁷ claims that four interrelated components are essential for a training program: learning tasks, supportive information, just-in-time information (i.e. information provided to the learner at the moment that it is needed), and part-task practice. To clearly establish the learning objectives

of the present study, a PPH guideline was created through discussion with a focus group (formed of invited obstetricians and anesthesiologists who were supervisors of residents) and was based on the most up-to-date clinical evidence.¹⁻³

Table 3.1 List of examples of expected tasks.

Subscales	Examples of tasks
Communication	The participant introduced himself or herself to the patient, explained the actions to the patient, and asked the patient about comorbidities, medications, and allergies
Teamwork	Team-support behavior and adequate sharing of information
Vital Signs	Checked heart rate, blood pressure, rechecked vital signs
Venous Access	Large caliber intravenous line, order blood samples
Exams	Complete blood count, renal and liver function tests, blood type and cross-match
Drug Management	Loading and maintenance doses of oxytocin
Mechanical Management	Massaged uterus, emptied bladder, and checked for lacerations
Surgical Management	Suture or placental tissue removal if the scenario required action

Figure 3.1 outlines the two training formats and the clinical case scenarios. The instructional design group format comprised eight steps, including three training scenarios with increasing levels of complexity. An obstetrician was present in the delivery room to provide “over the shoulder” just-in-time guidance and corrective feedback to participants, whenever necessary. The best practice group received the PPH guideline before undergoing a three-step training format, one of which was a training scenario. This format replicated the best PPH simulation training¹¹ and was created using articles identified in the literature. Seven experts (five from an educational background and two who had healthcare training) used a rating scale based on Merrill’s First Principles of Instruction⁸ to analyze the articles from an instructional design perspective.




Figure 3.1 Description of the steps for the instructional design group and the best practices group training formats. The pretest and post-test scenarios were identical for both training formats.

The pretest and post-test scenarios were the same for both groups. They were offered immediately before and after the training scenarios and presented an intermediate level of complexity. The third instructional design format training scenario (most complex scenario) was identical to the sole training scenario used in the best practice format and corresponded with a previous description.¹¹

All scenarios were scripted as PPH after vaginal delivery and included a standardized patient, a standardized nurse, and a part-task pelvis simulator (EVA Simulador Pós-Parto; ProDelphus, Olinda, Brazil). Consequently, the nature of potentially expected tasks comprised communication, teamwork, and clinical management. The low-cost part-task simulator allowed variation in the site (vaginal laceration, uterus, or both) and intensity of the bleed, and was managed by the standardized patient. The standardized nurses provided the participants with clinical data (vital signs, results of laboratory evaluation, and uterine responses). Both the standardized patients and nurses (who were all volunteer healthcare personnel) received the script for each scenario in advance of training to allow for rehearsal.

At each training session, participants were welcomed, introduced to the training equipment, and asked to suspend their disbelief (i.e. act as though the simulation was a real-life clinical case) and not to share any information about the program until the end of all training sessions. Participants were then asked to form the teams outlined above and provided with the pretest scenario. All scenarios were either run for a maximum of 900 seconds or else interrupted at the point that the team diagnosed and corrected the main cause of the bleeding.

All scenarios were video recorded for either debriefing purposes (the training scenarios) or to enable future analysis for scoring and comparisons (the pretest and post-test scenarios). Two obstetricians were present at each session: BdM attended all the training sessions and 10 additional obstetricians alternated as the second content expert. Each pair of obstetricians was responsible for rating the pretest and post-test scenarios video recordings for each of the teams that they supervised. The experts received the checklists of expected tasks at the beginning of each scenario, whereas the participants received them immediately after attending the first and third training scenario for debriefing purposes.

A standardized checklist of expected tasks (derived from the PPH guideline outlined previously) for each scripted scenario was used in the analysis. A task was considered to be executed if it had been performed by at least one participant (e.g. ordering laboratory evaluation) or, depending on the nature of the task, by all members of the team (i.e. teamwork tasks). Consequently, the unit of analysis was the team rather than individual participants.

At pretest, 38 tasks were expected to be executed; at post-test, 39 were expected. These tasks were distributed among eight learning subscales: communication (9 pretest/9 post-test), teamwork (6/6), vital signs (3/3), venous access (3/3), laboratory evaluation (6/6), mechanical management (4/5), drug management (6/6), and surgical management (1/1). Most pretest and post-test tasks were coincident, particularly for the first five subscales. The expected tasks varied according to the scripted case for the remaining three subscales. The number of tasks correctly executed by each team was registered in total, per subscale, and per time elapsed (in seconds). The entrance of the first team member into the delivery room defined the start of the process (0 seconds). The proportion of correct tasks was calculated for each of these landmarks. The medians of these proportions were used for comparison (instructional design group vs best practice group).

A generalizability analysis was used to estimate inter-rater reliability; a value of 0.93 for one rater (BdM) was considered to represent sufficiently high reliability.¹²

The timepoints used to compare groups of teams were 60, 360, and 900 seconds. The 60-second checkpoint reflected the “golden minute” concept in neonatal resuscitation guidelines; early initiation of basic resuscitation interventions are thought to be essential to prevent progression to circulatory collapse.¹³ The 360-second landmark corresponded to an additional 300 seconds, during which correct identification of the cause of bleeding and corresponding preliminary actions were expected.^{14,15} The 900-second checkpoint was the cut-off for the amount of time required to assess how the team managed the PPH case.

The data were analyzed using SPSS version 22 (IBM, Armonk, NY, USA). The post-test minus pretest difference in the median of the proportions of executed tasks was compared between the two groups for the various landmarks. The Mann–Whitney U test was used for comparison, with a P value

of less than 0.05 considered statistically significant. The effect size (r) was also calculated,¹⁶ and its magnitude was classified as small (0.10), medium (0.30), and large (≥ 0.50).¹⁷

Results

Overall, 54 (55.1%) of the 98 eligible residents attended training during their off-duty hours. Data regarding the proportion of all tasks correctly executed by the two groups is presented in Table 3.2. Both groups presented similar median proportions at pretest. However, when comparing the proportion of all tasks executed from pretest to post-test, a greater increase was found for the instructional design group (median difference 0.46, interquartile range [IQR] 0.147) than for the best practice group (median difference 0.17, IQR 0.262). The U value for the comparison was 5.0 ($P < 0.001$) and the effect size was 0.72.

Table 3.2 Proportion of tasks completed per team for Instructional Design and Best Practice and statistical comparisons

Type of Task	N tasks pre/post	Best Practices based (n = 7 teams)						Instructional Design based (n = 13 teams)						Statistical comparisons between training formats' groups			
		Pre-test		Post-test		Difference		Pre-test		Post-test		Difference		Difference of the medians		Effect size r^d	
		Md	IQR	Md	IQR	Md	IQR	Md	IQR	Md	IQR	Md	IQR	Post-Pre scores	U^e		p^d
Communication	9/9	0.33		0.44		0.22		0.22		0.78		0.56		0.33	13.0	0.004	0.58
Teamwork	6/6	0.67		0.83		0.17		0.67		0.83		0.17		0.00	38.0	0.294	0.15
Vital Signs	3/3	0.67		1.00		0.33		0.67		1.00		0.33		0.00	44.0	0.469	0.03
Venous access	3/3	0.67		0.67		0.00		0.67		1.00		0.33		0.33	25.5	0.057	0.38
Exams	6/6	0.00		0.33		0.00		0.00		0.83		0.83		0.83	3.0	0.000	0.76
Mechanical M	4/5 ^{a)}	0.75		0.60		-0.15		0.75		0.80		0.25		0.40	24.0	0.048	0.39
Drug Management	6/6 ^{a)}	0.33		0.67		0.33		0.17		0.83		0.50		0.17	25.5	0.057	0.36
All total time	37/38 ^{b)}	0.41	0.162	0.58	0.237	0.17	0.262	0.35	0.122	0.79	0.105	0.46	0.147	0.30	5.0	0.000	0.72
All within 60 seconds	33/34 ^{b)}	0.15	0.061	0.21	0.029	0.05	0.092	0.18	0.076	0.35	0.176	0.20	0.131	0.15	18.0	0.015	0.49
All within 360 seconds	33/34 ^{b)}	0.30	0.061	0.53	0.235	0.26	0.237	0.30	0.076	0.76	0.059	0.49	0.135	0.23	9.0	0.001	0.65

Note. Md = Median; IQR = Interquartile range M = management

This table provides an overview of proportions of tasks correctly completed – Medians¹³ and Interquartile ranges (IQRs) – in the ID group and BP group, and statistical comparisons for both pretest and posttest on the seven different subscales as well as on all subscales, all subscales in the first minute, and all subscales in the first six minutes.

^{a)} Tasks for pre-and post-test were not all the same.

^{b)} Four Tasks had no time indication and therefore were not included in the time related analyses

^{c)} Mann-Whitney U-statistic

^{d)} p-value of between-group difference for the Mann-Whitney test (one-sided)

^{e)} effect size $r = z/\sqrt{N}$; z is the z-score corresponding to the Mann-Whitney U-statistic; N is the total number of units in the two groups

The instructional design group exhibited a statistically significant greater increase from pretest to post-test than did the best practice group for three of the eight subscales: communication, laboratory evaluation, and mechanical management (Table 3.2). No comparison was made for the surgical management subscale because it comprised only one task, both at pretest and post-test. Additionally, at post-test, the expected task involved manual removal of the retained placental tissue from the part-task simulator. Some of the teams—five from the instructional design group and three from the best practice group—did not complete this task because they did not want to damage the simulated pelvis. Owing to the heterogeneity of the supervisors' approach to overcoming this problem during training, it was decided to exclude the surgical management subscale from the analysis.

Figure 3.2 presents the post-test minus pretest difference in the proportion of tasks completed by both the instructional design and best practice groups of teams after 60 and 360 seconds. A greater increase from pretest to post-test was observed for the instructional design group versus the best practice group at both time points. As shown in Table 3.2, the instructional design group showed a greater increase of correctly performed tasks after 60 seconds (median difference 0.20, IQR 0.131) than did the best practice group (median difference 0.05, IQR 0.092). The U value for the comparison was 18.0 ($P=0.015$) and the effect size was 0.49, indicating that the instructional design group was not only performing more tasks correctly but also earlier in the process than the best practice group. A similar result was observed after 360 seconds. The median differences for the instructional design and best practice groups were 0.49 (IQR 0.135) and 0.26 (IQR 0.237). The U value for the comparison was 9.0 ($P=0.001$) and the effect size was 0.65.




Figure 3.2 Proportion of tasks completed by the teams in the best practices group (dashed line, N=7 teams) and the instructional design group (solid line, N= 13 teams). The upper panel shows pretest (thin lines) and post-test (bold) proportions, and the lower panel presents the post-pre difference for the two groups.

Discussion

The present study demonstrated the effectiveness of the instructional design format for PPH simulation training. Participants who received such training exhibited a substantially greater increase in the number of executed tasks, and within a shorter time frame, than did participants assigned to the best practice training format. Thus, the findings of the present study confirmed the stated hypothesis.

The instructional design format offered several features not included in the best practice format. These features included broad content discussion at the prior knowledge step, multiple learning strategies, video demonstration of the expected tasks; clinical variation, increasing level of complexity, just-in-time information, corrective feedback, decreasing support, and two debriefings, with self-assessment in the second debriefing session. All these aspects are regarded to be effective for learning through simulation^{7,18,19} with particular recommendation by the Four-Component Instructional Design model.²⁰ Furthermore, potential confounding variables were controlled by delivering both training formats in an equivalent hospital environment.²¹

Although the ideal design for the present study would have been to randomly allocate participants to the two training formats, the locally elevated rates of PPH-related maternal morbidity and mortality required prioritization of a large number of participants through flexibility of the training schedule. Of note, however, both groups displayed similar median proportions at pretest.

The present findings could contribute to minimizing the recognized gap in comparative effectiveness research on simulation training formats. As previously noted,^{22,23} an extensive body of evidence provides plain description of training outcomes,²⁴ or comparison of different instructional sequences,²⁵ but not of different instructional design features.

The level of detail in the present findings allowed some interpretation. For example, the marked increase for the laboratory evaluation subscale could be extrapolated as ability acquired by the trainees to anticipate, recognize, and intercept ongoing management flaws and potential patient compromise.

This acquired ability might lead to increased situational awareness,⁹ a factor that is frequently jeopardized in the substandard care of patients with PPH.^{2,3}

The use of a portable part-task simulator, ordinary labor assistance material, portable filming, and a video exhibition device (electronic notebook) are all elements that favor future training replication and portability. With regard to PPH, these things could be of particular relevance, given that the most challenging aspects of this condition are its unpredictability and omnipresence.¹ Therefore, an effective portable PPH training package might increase the proportion of trained personnel in an undemanding manner. Moreover, the essential structure to the instructional design training format could be easily adapted to other obstetric emergencies, clinical content, and types of trainees.


Despite the nonsignificant differences observed for some of the subscales assessed in the present study, there was an increase in the overall proportion of tasks executed from pretest to post-test. For the teamwork subscale, measurement difficulty is a well-recognized issue owing to its multiple dimensions.^{5,26} The low number of expected tasks for the both the vital signs and venous access subscales could have limited the comparison; however, all the expected tasks were correctly executed in the instructional design group. The unanticipated finding of a negative value for the mechanical management subscale among the best practice group might be attributed to the unfamiliarity of these trainees with the simulator, given that they had attended fewer scenarios than the instructional design group. The time-on-task difference between training formats could be highlighted as a potential limitation because it reflects a core difference between current best practice and instructional design training formats.

In conclusion, the findings of the present study indicated the importance of applying established learning principles, such as instructional design guidelines, when designing simulation training formats. Optimum training time and the impact of particular features on knowledge transfer and clinical outcomes remain to be explored.

References

1. World Health Organization. WHO recommendations for the prevention and treatment of postpartum haemorrhage. Italy, 2012.
2. Bohlmann MK, Rath W. Medical prevention and treatment of postpartum hemorrhage: a comparison of different guidelines. *Arch Gynecol Obstet* 2014;289:555-67.
3. Knight M, Kenyon S, Brocklehurst P, et al. Saving lives, improving mothers' care – lessons learned to inform future maternity care from the UK and Ireland confidential enquiries into maternal deaths and morbidity 2009–12. Oxford, UK 2014.
4. Cook DA, Hamstra SJ, Brydges R, et al. Comparative effectiveness of instructional design features in simulation-based education: systematic review and meta-analysis. *Med Teach* 2013;35:e867-98.
5. Mérien AER, van de Ven J, Mol BW, Houterman S, Oei SG. Multidisciplinary team training in simulation setting for acute obstetric emergencies: a systematic review. *Obstet Gynecol* 2010;115:1021-31.
6. Ameh CA, van den Broek N. Making It Happen: Training health-care providers in emergency obstetric and newborn care. *Best Pract Res Clin Obstet Gynaecol* 2015;29:1077-91.
7. van Merriënboer JGG, Kirschner PA. *Ten Steps to Complex Learning*. 3rd ed. New York, NY: Routledge; 2018.
8. Merrill MD. *First Principles of Instruction: Identifying and designing effective, efficient and engaging instruction* San Francisco, CA, USA: Pfeiffer; 2013.
9. Edozien LC. Situational awareness and its application in the delivery suite. *Obstet Gynecol* 2015;125:65-9.
10. Cornthwaite K, Edwards S, Siassakos D. Reducing risk in maternity by optimising teamwork and leadership: An evidence-based approach to save mothers and babies. *Best Pract Res Clin Obstet Gynaecol* 2013;27:571-81.
11. Clark EA, Fisher J, Arafah J, Druzin M. Team training/simulation. *Clin Obstet Gynecol* 2010;53:265-77.
12. Streiner DL, Norman GR. *Health measurement scales: A practical guide to their development and use*. 4th ed. Oxford ; New York: Oxford University Press; 2008.
13. Ersdal HL, Mduma E, Svensen E, Perlman JM. Early initiation of basic resuscitation interventions including face mask ventilation may reduce birth asphyxia related mortality in low-income countries: a prospective descriptive observational study. *Resuscitation* 2012;83:869-73.

14. Crowe SD, Faulkner B. Lean management system application in creation of a postpartum hemorrhage prevention bundle on postpartum units. *Obstet Gynecol* 2014;123:455-5.
15. Deering SH, Chinn M, Hodor J, Benedetti T, Mandel LS, Goff B. Use of a postpartum hemorrhage simulator for instruction and evaluation of residents. *J Grad Med Educ* 2009;1:260-3.
16. Field A. *Discovering Statistics Using SPSS*. 4th ed. London: Sage; 2013.
17. Cohen J. *Statistical power analysis for the behavioral sciences*. 2nd ed. Hillsdale, N.J.: L. Erlbaum Associates; 1988.
18. Issenberg SB, McGaghie WC, Petrusa ER, Lee Gordon D, Scalese RJ. Features and uses of high-fidelity medical simulations that lead to effective learning: a BEME systematic review. *Med Teach* 2005;27:10-28.
19. McGaghie WC, Issenberg SB, Petrusa ER, Scalese RJ. Effect of practice on standardised learning outcomes in simulation-based medical education. *Med Educ* 2006;40:792-7.
20. Maggio LA, Cate OT, Irby DM, O'Brien BC. Designing Evidence-Based Medicine Training to Optimize the Transfer of Skills From the Classroom to Clinical Practice: Applying the Four Component Instructional Design Model. *Acad Med* 2015;90:1457-61.
21. Cook DA. Avoiding confounded comparisons in education research. *Med Educ* 2009;43:102-4.
22. McGaghie WC, Issenberg SB, Cohen ER, Barsuk JH, Wayne DB. Does simulation-based medical education with deliberate practice yield better results than traditional clinical education? A meta-analytic comparative review of the evidence. *Acad Med* 2011;86:706-11.
23. Cook DA, Hatala R, Brydges R, et al. Technology-Enhanced Simulation for Health Professions Education A Systematic Review and Meta-analysis. *JAMA* 2011;306:978-88.
24. Fialkow MF, Adams CR, Carranza L, Golden SJ, Benedetti TJ, Fernandez R. An in situ standardized patient-based simulation to train postpartum hemorrhage and team skills on a labor and delivery unit. *Simul Healthc* 2014;9:65-71.
25. Stefaniak JE, Turkelson CL. Does the sequence of instruction matter during simulation? *Simul Healthc* 2014;9:15-20.
26. Salas E, Grossman R, Hughes AM, Coultas CW. Measuring team cohesion: observations from the science. *Hum Factors* 2015;57:365-74.





Chapter 4

Self-perceived long-term transfer of learning after postpartum hemorrhage simulation training³

³ This chapter is based on the published article: De Melo BCP, Falbo AR, Sorensen JL, van Merriënboer JJG, & van der Vleuten CPM. (2018). Self-perceived long-term transfer of learning after postpartum hemorrhage simulation training. *International Journal of Gynaecology and Obstetrics: The Official Organ of the International Federation of Gynaecology and Obstetrics*, 141, 261-267. DOI: 10.1002/ijgo.12442

Abstract

Introduction

Effective simulation training should promote long-term transfer (application of acquired knowledge and skills on the job).

Objective

To explore long-term transfer after postpartum hemorrhage simulation training based on either instructional design principles or conventional best practice.

Methods

In this qualitative study, semi-structured interviews with obstetrics and gynecology healthcare practitioners were conducted between August 7 and September 26, 2015, in Recife, Brazil. The participants were randomly selected from each of two postpartum hemorrhage simulations attended two years earlier (one ID and one conventional best practice). Thematic analysis was used to explore (1) residents' perceptions of long-term transfer of learning, (2) ID elements influencing the perceived long-term transfer, and (3) differences in the participants' perceptions according to the type of simulation attended.

Results

There were 12 interview participants. After either simulation format, residents perceived long-term transfer effects. Training design factors influencing transfer were, in their opinion, related to trainees' characteristics, simulation design, and workplace environment. Trainees who participated in the ID-based simulation perceived better communication skills and better overall situational awareness: "I didn't do that before."

Conclusion

All residents perceived long-term transfer after simulation training for postpartum hemorrhage. Those who attended the ID format additionally perceived improvements in communication skills and situational awareness, which are fundamental factors in the management of postpartum hemorrhage.

Introduction

Postpartum hemorrhage (PPH) is the leading cause of maternal mortality worldwide, with most deaths considered preventable and attributable to human factors such as poor communication and lack of situational awareness. To improve the management of PPH, simulation training is strongly recommended.¹⁻³ Effective training, however, relies on workplace application of the knowledge and skills acquired during training—in other words, on transfer of learning.⁴

Systematic reviews on the effectiveness of healthcare simulation training have recommend the implementation of instructional design principles, particularly when aiming to achieve transfer;^{5,6} yet, these principles are sparsely adopted. Data on transfer from healthcare simulation training are scarce because of the complexity of measuring on-the-job performance, and studies have mainly explored the perceptions of trainees on short-term transfer (hours or weeks after training).⁷ Effective training, however, should also promote long-term transfer.⁸

The present study was conducted to explore self-perceived long-term transfer 2 years after having attended PPH simulation training based on either ID principles or current best practice (BP). The ID format applied principles derived from cognitive psychology that are explicitly aimed at optimizing long-term transfer;^{9,10} the ID format used in the present study included multiple elements (Table 4.1).

Table 4.1 List of instructional design guideline elements used in the ID simulation format

What	Concept	Application in this study
<i>Authenticity</i> ⁹	<i>Real life, whole-tasks</i>	<i>Scenarios script details very similar to reality. SKA tasks expected.</i>
<i>Psychological fidelity</i> ¹¹	<i>Degree to which real-task skill(s) are replicated in the simulated task</i>	<i>In situ environment, simulated patient and simulated nurse contributed to trainees' suspension of disbelief (i.e. acting as if the scenario was real)</i>
<i>Engineering fidelity</i> ¹¹	<i>Degree to which the training device or environment replicates the physical characteristics of the real task</i>	<i>Scenarios occurring in situ, with simulated patient and interactive part-task simulator</i>
<i>Paivio's dual coding</i> ¹²	<i>Images visualization associated with instruction help people learn</i>	<i>A circle with continuous arrows summarized training content (Figure 4.1)</i>
<i>Feedback</i> ¹³	<i>Trainees receive information on their performance</i>	<i>Trainees received feedback (debriefing steps) on more than one occasion</i>
<i>Variability</i> ¹⁴	<i>Opportunity to practice different tasks in more than one scenario</i>	<i>Trainees attended three different scenarios</i>
<i>Increasing complexity</i> ¹⁴	<i>Learning tasks are presented in an increasing sequence of complexity</i>	<i>Scenarios presented in a sequence of increasing complexity</i>

SKA – Skills, knowledge and attitude




Figure 4.1 Postpartum hemorrhage management protocol.

The current study was designed to address the following research questions. (1) Do the residents perceive transfer 2 years after having attended PPH simulation training (ID or BP format)? (2) Which factors of the simulation training do they perceive as having positively affected long-term transfer? (3) Does the perceived long-term transfer differ according to the format (BP or ID) of the simulation training?

Methods

In the present qualitative study, semi-structured interviews were conducted between August 7 and September 26, 2015, with health-care practitioners who had attended one of two different PPH simulations (based on ID or BP) 2 years previously.¹⁵ At the time of the simulations, all attendees were residents at one of the five teaching hospitals in Recife, Brazil, in one of the 3 years of the obstetrics and gynecology program. Using a simple randomization method (masking of names), six attendees from each simulation format were selected for invitation to interview from a total of 54 attendees. The study was approved by the ethics committee of Instituto de Medicina Integral Professor Fernando Figueira, Recife, Brazil. All participants gave written informed consent.

The two PPH simulations differed with regard to the ID elements (Table 4.1) and the number of steps. The ID-based simulation contained eight steps: (1) prior knowledge activation, (2) video demonstration, (3) dual-coding PPH protocol discussion, (4) training scenario #1, (5) debriefing, (6) training scenario #2 with immediate feedback, (7) training scenario #3, and (8) debriefing with self-assessment. The training scenarios had an increasing level of complexity. The BP simulation contained three steps: (1) prior knowledge activation, (2) training scenario (identical to ID simulation training scenario #3), and (3) debriefing. In both scenarios, the debriefings included participants watching video of themselves from the simulations. This format replicated the “best PPH simulation” previously¹⁶ selected by seven training experts.

The PPH management protocol was developed by a focus group of obstetricians and anesthesiologists based on best available evidence.^{1,17} The protocol was presented as shown in Figure 4.1, in agreement with Paivio’s dual-coding theory,¹² which states that content presented via a combination of text and pictures is easier to remember than content presented with either method alone.

In both simulation formats, the scenarios included a standardized patient, a standardized nurse, and a part-task pelvis simulator (Postpartum bleeding station; ProDelphus, Olinda, Brazil). The trained domains comprised

skills, knowledge, and attitudes tasks such as communication, teamwork, and clinical management.

The interview guide used open-ended questions to explore residents' perceived long-term transfer (Table 4.2) and was based on Baldwin and Ford's transfer model.¹⁸ According to this model, three factors influence transfer: (1) trainee characteristics (the motivation to attend training, the perceived relevance of the training content, and self-confidence); (2) the training design (application of ID principles); and (3) the workplace environment (organizational feedback, teamwork, and opportunity to use what has been learned).^{18,19}

Table 4.2 Interview guide list of questions for the semi-structured questionnaire (the themes are inspired by Baldwin and Ford, 1988)

Questions	
Introductory Question 1	<i>Do you consider the simulation training to have changed how you manage a PPH case? If so, in what way?</i>
Introductory Question 2	<i>What were the "take home messages" from the simulations?</i>
Trainee characteristics	<i>What motivated you most to attend the simulations? How relevant do you consider the content of the simulations to be?</i>
Training design	<i>How authentic (from a clinical perspective) did you consider the scenarios to be? How much do you recall applying in a real PPH clinical case from what you learned in simulations? How much of what was learned do you recall applying in any real PPH case with an etiology different from those of the simulations? How different from the simulations scenario cases was it? How much do you think visualizing the PPH protocol facilitated your learning? How much do you recall from the debriefing step? How much do you think this feedback contributed to your learning? Do you think the increasingly complex sequence of scenarios contributed to your learning? How would you describe your confidence in attending a real PPH case after the simulations? How did you perceive your confidence during your unsupervised practice (at the rural maternity units)?</i>
Work environment	<i>After the simulations, did you have any form of feedback (comments, support, patient clinical outcome information) from the maternity unit supervisors after managing a real PPH case? Was it in your residency hospital or a hospital you work at as a staff member? After the simulations, how would you describe the teamwork on the PPH cases you attended?</i>
Comments & Suggestions	<i>Do you have any comments and/or suggestions regarding the simulations? What would they be?</i>

A fifth-year medical student was trained to conduct all one-to-one interviews in a private room in the hospital. The participants received two visual aids to facilitate recollection: the picture summarizing the PPH protocol (Fig. 1) and a picture summarizing the PPH simulation steps from each training format.¹⁵ The interviews were audio-recorded and typically lasted 20 minutes. They were transcribed verbatim and the participants' identities were coded according to the format of the attended simulation (Resident_ID# or Resident_BP#).

Thematic analysis²⁰ was used to analyze the interviews. Two researchers (BCPM and ARF), who were blinded to the participants' identity but not to the format of the attended simulation, independently read the transcripts of each interview and highlighted, labeled, and grouped relevant quotations in a coding table.

In a first discussion round, the readers compared the highlighted quotations and coding tables and agreed on two preconceived main categories: (1) perceived transfer and (2) simulation factors perceived as potentially affecting transfer. The simulation factors category was composed of three subcategories that were based on the model by Baldwin and Ford:¹⁸ (2a) trainee characteristics, (2b) simulation design, and (2c) work environment. One particular group of similar quotations did not fit into the preconceived main categories and was categorized as "systematic approach to PPH management".¹⁰

Subsequently, each reader independently created a table for each simulation format (BP or ID) and displayed the findings according to the main categories and subcategories. They then discussed differences in perceived long-term transfer per format. A final table containing the summary columns for each simulation format was translated into English for interpretation and discussion by the other authors. Saturation of the findings was achieved once the collected data allowed sufficient understanding of the dimensions and properties of our key concepts.²¹ No further sampling of residents was deemed necessary.

The English-language table contained quotations of the residents, organized by category, and notes from the readers. The notes described

reflections regarding the residents' perception of overall transfer; the potential effects of different simulation factors (in particular ID elements) on transfer; and observed differences in perceived transfer depending on the simulation format.

Results

All 12 potential interviewees accepted the invitations. The mean age of the interviewees was 29 years (range 28–31 years), the mean time since graduation from medical school was 4 years (range 2.5–5 years), and 11 interviewees were female.

The first research question was addressed by ascertaining whether residents from both simulation formats perceived long-term transfer effects. Residents consistently reported a perceived change in PPH management and the use of a more systematic approach to problem-solving after the simulations (Table 4.3).

Table 4.3 Extracted segments of Participants' interview responses

Research Questions	Quotations
<p>First research question: <i>How do participants perceive long-term transfer while performing on the job 2 years after attending the simulations (ID or BP)?</i></p>	<p>Perceived transfer <i>Resident_ID6: "What changed the most was the sequence of actions. What to start with: massage the uterus first, then go to medication, right? ... Acknowledging teamwork more, being aware of the need to call people to help, stabilize the patient, get a venous access, give her oxygen, monitor her vital signs... These were the things that really stuck in our minds after the training ... Now I have a more clearly defined sequence of actions ..."</i></p> <p><i>Resident_BP4: "I think it was mostly, like... more speed in my reasoning! Bleeding: then I do this, then this, then that. Didn't work! Then this, afterwards that ..."</i></p>
<p>Second research question: <i>Which factors of simulation do they perceive as positively affecting this long-term transfer awareness?</i></p>	
Trainee characteristics	
<p>Motivation and relevance of content</p>	<p><i>Resident_ID3: "If you work at a maternity, you have to practice. ... You need to have the young personnel trained to be aware of it"</i></p> <p><i>Resident_BP6: "Very frequent. Managing it properly will save lives."</i></p>
<p>Self-confidence</p>	<p><i>Resident_ID5: "A lot, 100%. You realize you are more confident, so you end up assuming more the leadership because of the training."</i></p>
Simulation design	
<p>Authenticity</p>	<p><i>Resident_BP1: "For sure we deal with very similar cases in our day-to-day work"</i></p> <p><i>Resident_ID1: "I recall I got very tense at that point, everybody got nervous. In this sense, it was very real."</i></p>
<p>Dual coding</p>	<p><i>Resident_ID1: "It is still in my mind today. I think it is great. I remember it every time. I recall the circle."</i></p> <p><i>Resident_BP1: "It helps a lot. ... Because we can see it and recall the sequence very clearly."</i></p>
<p>Feedback</p>	<p><i>Resident_BP1: "I recall we were in a room watching what we had done. It was very useful. ... You can analyze what you did wrong,</i></p>

then you see what would have been right and you remember it.”

Resident_BP1: “I remember well that the last case was much more complex, and we managed it much faster than the initial (simpler) cases.”

Increasing complexity (ID simulation only)

Resident_ID1: “It helped a lot. We were doing more and more increasingly more complex tasks.” **Resident_ID3:** “I think it [the increasing complexity of the scenarios] allowed us to gradually build up more knowledge”

Work environment

Feedback **Resident_BP1:** “From a supervisor, once, at my residency. He congratulated me saying I knew how to take action, [how to] handle the case.”

Teamwork and workplace learning skills **Resident_ID5:** “There was this one case, I was with a colleague who had also attended the training. So, things went very smoothly, we were thinking alike and getting feedback from each other. It was very interesting.”

Resident_ID4: “The anesthesiologists are the ones who complicate our lives the most. I was very upset one day when they kind of neglected a case ... only when the patient began to become unstable did they realize it was for real.”

Third research question: *Are there differences in perceived long-term transfer according to the type of simulation attended (BP or ID)?*

Differences in perceptions according to PPH simulation format

Resident_ID4: “First is that thing of introducing yourself, explaining to the patient what is going on and your actions ... I didn’t do that before. And collect exams, to remember to collect the exams ...”

Resident_ID2: “... already arriving at the scene talking to the patient, already managing the case, ... remembering to collect exams ...”

The findings relating to the second research question (influence of simulation factors on transfer) were presented as follows: (a) trainee characteristics (relevance, motivation, and self-confidence), (b) training design (authenticity, dual coding, feedback by debriefing, and variability and increasing complexity), and (c) work environment (organizational feedback, teamwork, and opportunity to use what was learned).

The reported motivation for attending the simulations was a desire to learn and improve professional skills. The residents acknowledged that the content of the simulations (management of PPH) was relevant to them. Moreover, they perceived heightened self-confidence in the management of patients with PPH in their real work environment after the simulations, in particular when working in an unsupervised situation (outside their residency programs). The simulations gave them the confidence to assume team leadership when necessary (Table 4.3).

With regard to the training design, the scenarios were considered authentic in so far as they genuinely replicated clinical cases encountered in the residents' workplace practice, with similar reasoning challenges. Furthermore, the simulations occurred in situ (in a real environment) in the presence of a simulated nurse and a simulated patient. All these items contributed to the psychological fidelity (mental similarity to reality) of the simulations, reflected by a perceived sense of anxiety during the simulations (Table 4.3). The engineering fidelity (physical similarity) of the part-task simulator was also mentioned as a factor contributing to the authenticity (overall similarity to real cases) of the simulations.

The dual-coding strategy was also acknowledged as facilitating transfer. Residents strongly recollected the visual representation of the PPH protocol and specifically referred to the central circle with continuous arrows, which reflected the need to execute continuous and simultaneous tasks.

The residents reported that watching themselves on video during the debriefing (feedback) in both simulation formats allowed them to identify and reflect on their own errors and make plans for how to improve their skills. They also acknowledged that the safe feedback environment facilitated learning.

Residents who participated in the ID format were exposed to more than one training scenario and debriefing, which enabled them to try out variations

and practice under conditions of increasing complexity. Residents attending this format reported that exposure to this training sequence facilitated learning (Table 4.3).

As for working-environment factors, organizational feedback was described as poor. With one exception, all residents said they had not received any workplace feedback (feedback from supervisors or institutional feedback).

The residents perceived an improvement in teamwork skills because they were both more aware of them after participating in the simulations and more motivated to take on a leadership role when faced with a real patient with PPH. Residents who had managed a woman with PPH along with a colleague who had also attended the simulation stressed the smoothness of the actions and the shared responsibility in re-evaluating the diagnosis and revising the management plan. However, the residents also cited teamwork conflicts related to resistance from colleagues from other healthcare disciplines who had not attended the simulation training (Table 4.3).

With regard to the third research question, residents who had attended the ID-based simulation attributed the perceived transfer to improvements in both communication and teamwork skills, and also to situational awareness in the workplace. Moreover, only residents who had attended the ID simulation format reported communication with the bleeding woman and the collection of blood samples for examination early in the process, which reflects an awareness that the patient is potentially instable (increased situational awareness) (Table 4.3). The findings relating to the second research question (influence of simulation factors on transfer) were presented as follows: (a) trainee characteristics (relevance, motivation, and self-confidence), (b) training design (authenticity, dual coding, feedback by debriefing, and variability and increasing complexity), and (c) work environment (organizational feedback, teamwork, and opportunity to use what was learned).

The reported motivation for attending the simulations was a desire to learn and improve professional skills. The residents acknowledged that the content of the simulations (management of PPH) was relevant to them. Moreover, they perceived heightened self-confidence in the management of patients with PPH in their real work environment after the simulations, in

particular when working in an unsupervised situation (outside their residency programs). The simulations gave them the confidence to assume team leadership when necessary (Table 4.3).

With regard to the training design, the scenarios were considered authentic in so far as they genuinely replicated clinical cases encountered in the residents' workplace practice, with similar reasoning challenges. Furthermore, the simulations occurred in situ (in a real environment) in the presence of a simulated nurse and a simulated patient. All these items contributed to the psychological fidelity (mental similarity to reality) of the simulations, reflected by a perceived sense of anxiety during the simulations (Table 4.3). The engineering fidelity (physical similarity) of the part-task simulator was also mentioned as a factor contributing to the authenticity (overall similarity to real cases) of the simulations.

The dual-coding strategy was also acknowledged as facilitating transfer. Residents strongly recollected the visual representation of the PPH protocol and specifically referred to the central circle with continuous arrows, which reflected the need to execute continuous and simultaneous tasks.

The residents reported that watching themselves on video during the debriefing (feedback) in both simulation formats allowed them to identify and reflect on their own errors and make plans for how to improve their skills. They also acknowledged that the safe feedback environment facilitated learning.

Residents who participated in the ID format were exposed to more than one training scenario and debriefing, which enabled them to try out variations and practice under conditions of increasing complexity. Residents attending this format reported that exposure to this training sequence facilitated learning (Table 4.3).

As for working-environment factors, organizational feedback was described as poor. With one exception, all residents said they had not received any workplace feedback (feedback from supervisors or institutional feedback).

The residents perceived an improvement in teamwork skills because they were both more aware of them after participating in the simulations and more motivated to take on a leadership role when faced with a real patient with PPH. Residents who had managed a woman with PPH along with a colleague

who had also attended the simulation stressed the smoothness of the actions and the shared responsibility in re-evaluating the diagnosis and revising the management plan. However, the residents also cited teamwork conflicts related to resistance from colleagues from other healthcare disciplines who had not attended the simulation training (Table 4.3).

With regard to the third research question, residents who had attended the ID-based simulation attributed the perceived transfer to improvements in both communication and teamwork skills, and also to situational awareness in the workplace. Moreover, only residents who had attended the ID simulation format reported communication with the bleeding woman and the collection of blood samples for examination early in the process, which reflects an awareness that the patient is potentially instable (increased situational awareness) (Table 4.3).

Discussion

In the present study, residents perceived positive long-term transfer 2 years after having attended simulation training for PPH, regardless of the simulation format (ID and BP). The reported motivation to attend the simulations was a desire to improve professional skills. The residents acknowledged that PPH was a relevant training topic and described an increase in self-confidence. Attendees of either simulation format reported the following ID elements as contributors to transfer: authenticity,⁹ use of Paivio's dual-coding strategy,¹² and feedback during debriefing.^{14,22} Residents who participated in the ID-based simulation additionally mentioned variability and increasing complexity as ID elements contributing to transfer. Residents from both groups reported an increase in teamwork skills. They also reported poor organizational feedback and team struggles with colleagues who had not attended the simulation. Residents who had attended the ID format perceived better transfer of communication and teamwork skills and higher situational awareness than those having attended the BP format.

The present study makes several contributions to the transfer literature. Whereas most studies^{4,7} have explored transfer in the short term (immediately after training or a few weeks later), the present study demonstrated positive transfer 2 years after the PPH simulation training (long-term transfer). Moreover, the perceived improvements in communication skills and situational awareness reinforce the recommendation of applying ID principles when devising simulations, in particular simulations for high-stake situations such as PPH.^{1,5,6,10} In addition, because the study explored the perceptions of participants who, in their majority, had already concluded their residency, the study provides a wide range of perceptions deriving from actual workplace performance. This is likely to have had a positive influence on the perception of long-term training transfer.¹⁴

With regard to the influence of trainee characteristics, the motivation to improve one's professional skills has long been recognized as an important predictor for the effectiveness of transfer.^{4,19,23} The present study confirms this notion. The acknowledgement of PPH as a relevant topic for simulation training can be explained by the facts that PPH is a preventable cause of maternal death and that there has been a worrisome increase in the rate of PPH, which has led to recommendations for healthcare personnel to participate in simulation training.^{1,2,17} The long-time gap between training and the present evaluation of transfer reinforces the strength of the finding that self-confidence was increased.²³ Whereas most previous studies have assessed self-confidence in the short term,^{4,7,19} residents in the present study were consulted after several on-the-job exposures to PPH, which will have promoted better on-the-job learning.¹⁴

As for the training design, the following ID elements were reported as contributors to training transfer. First, authenticity,^{9,10} because the scenarios were close to reality and possessed psychological and engineering fidelity, which also contributed to the "suspension of disbelief" among the trainees.¹¹ Although there is an ongoing debate as to the exact definition of authenticity,^{9-11,24} there is a general consensus that it is important for trainees to perceive scenarios as "challenging" and to feel motivated to exert effort;¹⁹ this was indeed the case with the present scenarios. Second, the finding that

visualization of the PPH protocol helped the residents to recall what had been learned reinforces Paivio's dual-coding theory.¹² This might be of particular relevance to simulations of high-stake situations such as PPH, where several tasks have to be performed simultaneously and in collaboration.²⁵ Third, feedback during debriefing is broadly recognized as being important for learning;^{10,22} such feedback was provided more than once in the ID group. The residents stressed the importance of having the opportunity to learn from their own mistakes, a finding that is supported by evidence that error management improves transfer.^{4,25} Finally, the fact that attendees of the ID format acknowledged the variability and increasing complexity of the scenarios as contributors to transfer is in agreement with the literature on complex learning.^{4,14}

When discussing the impact of the work environment (opportunity to use skills in practice, organizational feedback, teamwork skills) on transfer, the high prevalence of PPH1,¹⁷ is an important factor to consider. Given this high prevalence, the trainees had multiple on-the-job opportunities to apply the acquired skills, but it also emerged that poor organizational feedback was ubiquitous⁴, highlighting the need to implement new policies for organizational feedback. With regard to teamwork, the perceived empowerment to take on a leadership role corroborates the need to train teamwork skills, preferably in a multidisciplinary fashion as illustrated by the reported conflicts with colleagues from other disciplines.²⁵

The perception of enhanced communication skills and situational awareness by ID format residents reinforces the recommendation to apply ID principles when designing simulation training, particularly for high-stake situations such as PPH. The achievement of long-term transfer is particularly relevant to simulations for clinical conditions in which the optimal coordination of simultaneous tasks from multiple domains has an important impact on patient outcomes.^{1,2,14,25}

The fact that the present analysis was based on self-perceived outcomes may be seen as a limitation. Indeed, strict analysis of transfer requires the overcoming of a few methodological challenges. For instance, to analyze PPH management in real life, uninterrupted video recording of all deliveries would

be necessary (with all the accompanying costs and operational implications) because the occurrence of PPH is unpredictable. Another potential limitation is related to the fact that the ID and BP simulation formats contained a different number of scenarios. This reflects a core difference between the two formats: with the larger number of scenarios, the ID format provided multiple practice opportunities, variability, and increasing complexity, which promotes cognitive elaboration because learners are encouraged to compare and contrast scenarios with each other. In fact, these aspects reinforce the trustworthiness of the present findings, which were obtained 2 years after the actual simulations and after potential contamination from unrelated work experiences.^{10,14}

In conclusion, the application of ID principles to PPH simulation training led to improvements in the perceived transfer of communication, teamwork, and situational awareness skills. Future studies should explore the contribution of specific ID elements to long-term transfer and their potential impact on clinical outcomes.

References

1. World Health Organization. WHO recommendations for the prevention and treatment of postpartum haemorrhage. Italy, 2012.
2. Wong CA, Scott S, Jones RL, Walzer J, Geller S. The state of Illinois obstetric hemorrhage project: pre-project and post-training examination scores. *J Matern Fetal Neonatal Med* 2016;29:845-9.
3. Edozien LC. Situational awareness and its application in the delivery suite. *Obstet Gynecol* 2015;125:65-9.
4. Salas E, Tannenbaum SI, Kraiger K, Smith-Jentsch KA. The science of training and development in organizations: What matters in practice. *Psychol Sci Public Interest* 2012;13:74-101.
5. Cook DA, Hamstra SJ, Brydges R, et al. Comparative effectiveness of instructional design features in simulation-based education: systematic review and meta-analysis. *Med Teach* 2013;35:e867-98.
6. McGaghie WC, Issenberg SB, Petrusa ER, Scalese RJ. A critical review of simulation-based medical education research: 2003-2009. *Med Educ* 2010;44:50-63.

7. Mema B, Harris I. The barriers and facilitators to transfer of ultrasound-guided central venous line skills from simulation to practice: exploring perceptions of learners and supervisors. *Teach Learn Med* 2016;28:115-24.
8. Burke LA, Hutchins HM. Training transfer: An integrative literature review. *Hum Resour Dev Rev* 2016;6:263-96.
9. Merrill MD. First Principles of Instruction. *ETR&D* 2002;50:43-59.
10. van Merriënboer JG, Kirschner PA. *Ten Steps to Complex Learning*. 3rd ed. New York, NY: Routledge; 2018.
11. Maran NF, Glavin RF. Low- to high-fidelity simulation – a continuum of medical education? *Med Educ* 2003;37:22-8.
12. Clark JM, Paivio A. Dual Coding Theory and education. *Educ Psychol Rev* 1991;3:149-210.
13. Clark RE, Voogel A. Transfer of training principles for instructional design. *ECTJ* 1985;33:113-23.
14. van Merriënboer JG, Kester L, Paas F. Teaching complex rather than simple tasks: Balancing intrinsic and germane load to enhance transfer of learning. *Appl Cogn Psychol* 2006;20:343-52.
15. de Melo BC, Falbo AR, Muijtjens AM, van der Vleuten CP, van Merriënboer JJ. The use of instructional design guidelines to increase effectiveness of postpartum hemorrhage simulation training. *Int J Gynaecol Obstet* 2017;137:99-105.
16. Clark EA, Fisher J, Arafeh J, Druzin M. Team training/simulation. *Clin Obstet Gynecol* 2010;53:265-77.
17. Mavrides E, Allard S, Chandrachan E, et al. Prevention and management of postpartum haemorrhage: Green-top Guideline No. 52. *BJOG* 2017;124:e106-e49.
18. Baldwin TT, Ford JK. Transfer of training: A review and directions for future research. *Pers Psychol* 1988;41:63-105.
19. Blume BD, Ford JK, Baldwin TT, Huang JL. Transfer of training: A meta-analytic review. *J Manage* 2009;36:1065-105.
20. Braun V, Clarke V. Using thematic analysis in psychology. *Qual Res Psychol* 2006;3:77-101.
21. Morse JM. The significance of saturation. *Qual Health Res* 1995;5:147-9.
22. Fanning RM, Gaba DM. The role of debriefing in simulation-based learning. *Simul Healthc* 2007;2:115-25.
23. Liaw SY, Scherpbier A, Rethans JJ, Klainin-Yobas P. Assessment for simulation learning

outcomes: A comparison of knowledge and self-reported confidence with observed clinical performance. *Nurse Educ Today* 2012;32:e35-9.

24. Grierson LE. Information processing, specificity of practice, and the transfer of learning: considerations for reconsidering fidelity. *Adv Health Sci Educ Theory Pract* 2014;19:281-9.
25. Stocker M, Burmester M, Allen M. Optimisation of simulated team training through the application of learning theories: a debate for a conceptual framework. *BMC Medical Education* 2014;14:2-9.



This chapter is embargoed at request

Chapter 5

Effects of an in situ instructional design based postpartum hemorrhage simulation training on patient outcomes: an uncontrolled before-and-after study⁴

⁴This chapter is based on a manuscript submitted for publication: De Melo BCP, van der Vleuten CPM, Muijtjens AMM, Falbo AR, Katz L, & van Merriënboer JIG. (submitted). Effects of an in situ instructional design based postpartum hemorrhage simulation training on patient outcomes: an uncontrolled before-and-after study



Chapter 6

General Discussion

General Discussion

This thesis aimed to explore the use of evidence-based instructional design (ID) guidelines for simulation training applied to a postpartum hemorrhage (PPH) content. Over the past decade, the worldwide spread of simulation training has been propelled by a genuine will to improve patient outcomes.¹⁻³ With the aim of optimizing simulation effectiveness, the use of ID guidelines has been recommended for this training strategy, in particular when focusing on the complex learning demands of high-risk situations.^{4,5} PPH is an example of a high-risk situation, with wide ecological validity and sound epidemiological relevance as the leading cause of maternal deaths worldwide with most of these considered as preventable.⁶⁻⁸

We aimed to explore the use of evidence-based instructional design in four different aspects related to training effectiveness: learning, transfer of learning, patient outcomes, and the use of ID guidelines in PPH simulation trainings as currently described in the literature. Our verified findings provide support to the current recommendation of adopting ID guidelines, as they demonstrate a positive influence on each of the analyzed aspects of training effectiveness. In addition, the verified neglect of ID guidelines use by the majority of simulation trainings described in the literature reinforces the relevance of our findings.

We began this thesis' studies series by analyzing the use of ID guidelines by PPH simulation trainings described in published articles. We found a pervasive neglect of its use by the majority of analyzed simulation trainings, as presented in Chapter 2. Then, as described in Chapter 3, we conducted a pretest-post-test learning outcomes comparison of residents who attended two different PPH simulation trainings: one based on ID guidelines and the other on current best practices. On the post-test, we found the residents who attended the ID based PPH simulation training to execute a greater number of tasks in a shorter period of time than their peers who attended the best practices format. In the study presented in Chapter 4, we explored the self-perceived long-term training transfer after PPH simulation training and found a

greater situational awareness and better communication skills perceptions by the residents who attended the ID based simulation. Ultimately, we analyzed the impact of the ID based PPH simulation training on patient outcomes and, as described in Chapter 5, we found an improvement in PPH management, with fewer patients receiving a greater mean dose of therapeutic oxytocin, which likely reflects an increased situational awareness after the training.

The chapters presented in this thesis may be perceived as having an hierarchical connection, with some parallel to Kirkpatrick's training evaluation model.^{9,10} According to this model, training effectiveness should be assessed on four different levels: (1) learner reaction, (2) learning outcomes, (3) behavior (transfer), and (4) results.

In our studies, we did not establish effects on Kirkpatrick's first level (learner reaction) but decided to explore the use of ID guidelines in previously published PPH simulation training studies, as presented in Chapter 2. The reason for this replacement was based on critiques on the Kirkpatrick model, which indicate little correlation between reaction and learning while also underscoring the model's simple taxonomy of training outcomes.¹¹⁻¹⁴ For the analysis of the remaining perspectives, we avoided simple replication of previously applied study designs and aimed at taking a step further in our analysis strategies. Therefore, we explored multiple aspects of simulation effectiveness through a "wide-angle lens" analysis,¹⁵⁻¹⁷ at each of our research questions listed below:

1) How are instructional design guidelines described and used in the current postpartum hemorrhage healthcare simulation literature?

2) What are the different learning outcomes when two instructionally different postpartum hemorrhage simulation training programmes are compared, one based on instructional design guidelines and the other on current best practices?

3) Is there long-term transfer after these two different training formats? Is there a difference in learning outcomes according to the attended training format? What instructional design features may have facilitated and/or hindered long-term transfer?

4) What is the impact on patient outcome at a teaching hospital after Obstetrics and Gynecology residents attending an instructional design based postpartum hemorrhage simulation training programme format?

As an example of a widened-angle analysis, our learning outcomes comparison study, presented in Chapter 3, had well specified complex learning tasks for our ID based PPH simulation training. Its whole-tasks learning approach differed from most often described studies in the learning outcomes simulation literature of single procedural tasks by promoting tasks from multiple dimensions (e.g. communication, procedural skills, teamwork, etc.).¹⁸⁻²²

Another example of a widened training assessment was our approach to exploring learners' perceived long-term transfer at the workplace after the simulation, presented in Chapter 4. Exploring such transfer two years after the training represents an analysis on a much longer period of time than usually reported.^{23,24} In addition, we adopted the Baldwin and Ford training transfer model²⁵ for our analysis to explore not only the influence of the training design on transfer but also the influence of trainees' characteristics (i.e. motivation to learn) and workplace environment (i.e. transfer of training conditions such as workplace feedback). Lastly, for the patient outcomes analysis, presented in Chapter 5, the study was designed to explore patient outcomes analysis in greater level of details than the usually reported impact on single procedural outcomes.^{18,23,26} Our patient outcomes study presented results on frequencies of specific PPH-related complications, duration of hospitalization, and specific management measures such as drugs usages and dosages. This is the type of analysis recommended by systematic reviews on simulation training translational outcomes, which critique the frequently reported analysis of single procedural outcomes.²⁷

Based on our presented findings, we expect to have successfully demonstrated the overarching benefits of ID guidelines for simulation training design. Our findings demonstrated positive outcomes throughout the complete training effectiveness spectrum – not only did

it lead to significant greater positive learning outcomes, but also to better perceived long-term transfer of learning and improved patient care.

Healthcare simulation training's ultimate goal is patient outcome improvement and, therefore, once embracing the challenge of promoting such training strategy it should be made effective by adopting ID guidelines. Moreover, as a first step, faculty should be made aware of the relevance of adopting such guidelines for designing and implementing simulation training.

Theoretical implications and future research

A few theoretical implications may be derived from our findings of a verified paucity of adherence to evidence-based instructional guidelines in current simulation training design. Moreover, the importance of these guidelines is reinforced by our additional positive outcomes of the use of such guidelines on simulation training effectiveness. These findings urge us to reflect upon the reasons that lead to such recurrent neglect from the healthcare simulation community with regards to training design.

An example of a theoretical implication is the need to further explore the reasons for such neglect from the stakeholders in charge of promoting simulation trainings worldwide. A natural conclusion derived from our presented findings is the sense of widespread "lack of awareness" from these stakeholders with regards to the necessity of adopting ID guidelines. The following step consists, therefore, in exploring possible strategies to promote faculty awareness and overcome potential resistances with regards to such necessity when designing simulation trainings.^{28,29}

Effectively promoting faculty awareness with regards to optimal simulation training design is fundamental, particularly when considering simulation's large spectrum of practical purposes. For instance, simulation trainings may span from curriculum-integrated learning to high stakes certificates.^{28,29} In addition, they may be applied to promote either learning of complex tasks (using whole-tasks in high fidelity or in-situ simulation) or procedural tasks (using part-tasks for part-task trainers), as well as training

single specialties and/or multidisciplinary tasks.^{21,22,30,31} Therefore, in our current era of evidence-based medicine, patient-centered care and the need for a lifelong learning culture among healthcare personnel, a learning strategy such as simulation must be promoted effectively, preferably based on both clinical and instructional best available evidence.³²⁻³⁴

Further exploring the effectiveness of specific instructional design features and/or combinations of these design features may also contribute to the current body of knowledge for optimal simulation training design. For instance, our study presented in Chapter 3 (the learning outcomes study) demonstrated the positive learning outcomes of an ID based PPH simulation format consisting of three training scenarios. However, future research may, for example, explore optimal training length with regards to the number of multiple scenarios sufficient to sustain complex learning through repetitive and simple-to-complex practice, or even, how to best demonstrate the to-be-learned tasks.^{31,35-37}

Further studies should also explore the impact of ID based training designed for healthcare workers with different levels of expertise, that is, not only for residents, but also for the whole continuum, from undergraduates to senior healthcare personnel. It is known that the effectiveness of training designs is heavily dependent on the prior knowledge or level of expertise of the trainees. Instructional design should, therefore, be adjusted accordingly taking into account the “expertise reversal effect” – avoiding redundant instruction to the experienced learner.^{38,39} Furthermore, it is essential to promote training for multiple disciplines, preferably promoting an integration of disciplines in teams.^{40,41}

Additional aspects of simulation training effectiveness beyond its design and content also remain to be explored. For instance, not only the faculty should be accounted for when promoting awareness with regards to the need to implement effective simulation training, but also the different stakeholders involved in promoting simulation, from both extremes of the administrative hierarchy: from the executive board to those responsible for ground work. A full-hospital simulation is still a challenge (simulation training exercises that include an entire care pathway and multiple hospital departments, thus truly

mimicking real clinical practice and exploring how to overcome the operational constraints implicated with routine hospital trainings across the entire patient pathway may lead to a significant contribution to simulation transfer.⁴² Simulation training transfer is significantly influenced by workplace-feedback and opportunities to use different aspects of the “post-training elements”, as described by Baldwin and Ford.²⁵ Promoting full-hospital simulation may significantly contribute to better understanding relevant training transfer constraints, which are reinforced by the workplace environment.^{43,44}

One other aspect of simulation training that is still underexplored is simulation cost-effectiveness.⁴⁵ In our series of studies, the combination of a low-technological, low-cost, part-task simulator with a simulated patient, maximized training tasks possibilities within a limited budget. Further exploration of an optimal balance between the financial investment required by simulation designs and settings and their impact on patient outcomes should be investigated further. Currently, the concept of high-value cost-conscious care (HVCC) has been introduced into the healthcare cost literature.⁴⁶ HVCC claims tests and treatments should have “high value”, meaning high benefit and low harm, in order to be cost-effective and widely promoted. Exploring for such a balance should also be a goal to the simulation training research field, as well on exploring how simulation may contribute to the training of healthcare personnel for HVCC.⁴⁷

Studies Strengths and Limitations

Among the strengths of our thesis, we highlight both the choice for the whole-task complex learning demands of our selected content, PPH, and the multiple practice opportunity provided by the three training scenarios organized in a simple-to-complex sequence. We believe they had a particularly relevant contribution to our verified positive findings on all different training effectiveness levels: learning outcomes, training transfer and patient outcomes. Our presented hybrid-simulation modality, as we combined a low technological, low-cost, part-task simulator with a simulated patient, allowed us to achieve multiple training outcomes – from communication to

procedural tasks and also contributed to training authenticity.³⁷ In addition, the easy reproducibility of the presented ID based training formats should also be pointed as a strength, in particular when considering the relevance of widely promoting healthcare simulation training for high-risk clinical conditions, such as PPH.^{48,49}

Our acknowledged limitations for our reported studies may be associated to those inherent to the constraints of a doctoral dissertation, mostly related to time and funding. For instance, our studies aimed at a single content, PPH. In spite of its wide external validity to other high-risk situations, an ID based simulation training design with positive training effectiveness outcomes should be replicated for other contents, in particular, those with demands for complex learning, both from the field of OBGYN and/or related to other healthcare specialties.^{31,50}

A single temporal sample of residents was included for the analysis of patient outcomes in a single center. Future studies should aim at a maximized impact analysis, including more residents in multiple centers, preferably through a stepped wedge cluster approach.⁵¹ Such study type includes an initial period in which no clusters are exposed to the intervention with subsequent clusters' progressively randomized and included to either control or intervention groups. It is used for the evaluation of interventions delivered at the level of the cluster and allows for robust scientific evaluations to be sought from such multiple cluster strategy, allowing, therefore a wider observation of facts.

While our presented study explored self-perceived long-term transfer, a different exploration strategy of the simulation training transfer aspect should also be aimed for. For instance, other methodologies to explore preferably real-time-on-the-job transfer should be considered, such as developing strategies to adequately quantify and analyze such tasks execution would need to be developed.⁴³ Meanwhile our findings of low institutional feedback reinforce our previous recommendation to further explore how to embrace faculty awareness with regards to simulation training. In addition, measuring and reporting such on-the-job-tasks in the healthcare environment implies a further exploration of how to overcome operational and ethical impediments.⁴⁴

Additional potential limitations were minimized and should in fact be considered as a study strength. For instance, the frequently observed seasonal influence in patient outcomes studies⁵² was minimized once we collected our patient outcome data at the same time of the year, one year before and immediately after the training. Such an operational orchestration should be underscored as an additional effort contributing to evidence-based healthcare simulation body of knowledge.²⁷

Implications for practice

A few tips on implications for practice may be derived from our findings and are presented in Box 1.

Box 1. Essential Implications for Practice Tips:

Use evidence-based ID guidelines for simulation training	
	<i>Multiple scenarios</i>
1	<i>Simple-to-complex sequence of scenarios</i> <i>Provide feedback and self-assessment opportunities</i>
2	<i>Prioritize whole-task learning content</i>
3	<i>Design simulation trainings for a multidisciplinary target-group</i>
4	<i>Promote faculty awareness with regards to the need of designing effective ID-based simulation</i>
5	<i>Promote institutional feedback routines to improve transfer</i>

Based on our presented findings we recommend designing healthcare simulation trainings carefully adopting evidence based ID guidelines. Multiple practices opportunities should always be provided while promoting clinical variation, simple-to-complex sequencing, and feedback with self-assessment opportunities.

Effective simulation training should always take into account the routine and non-routine aspects of the selected learning content and whole-task practice would provide such learning opportunities. Such precaution may be

especially relevant for complex high-risk situations, in particular those that are epidemiologically relevant such as PPH. As a first step, learning tasks should be well sought, therefore, preferably based on a carefully conducted needs assessment.

Designing simulation trainings for a multidisciplinary audience will contribute to a better transfer of learning. Reaching this goal implies, however, faculty awareness with regards to the benefits of effectively designed training strategies. They must be made aware of the contribution of simulation as an essential stage on the multitude of intertwined elements involved in promoting positive impact on patient outcomes. Not only the faculty, but the whole spectrum of the administrative hierarchy should be made aware of these benefits in order to also embrace and contribute to providing institutional feedback, which is an essential step in promoting better training transfer.

Promoting effective simulation training aimed at maximizing patient safety and improving patient outcomes has become particularly relevant in this current era of residency programs duty's restriction hours. These sets of recommendations should significantly contribute to reaching such patient outcomes improvement.

Take home message

The use of instructional design guidelines for PPH simulation training leads to simulation training effectiveness in terms of learning, transfer, and results. Therefore, in order to reach the ultimate goal of improving patient outcomes, simulation training must be designed to be effective and should thus be based on evidence-based instructional design guidelines.

References

1. Draycott TJ. Not all training for obstetric emergencies is equal, or effective. *BJOG* 2016;124:651.
2. Gaba DM. The future vision of simulation in health care. *Qual Saf Health Care*

- 2004;13:i2-i10.
3. Norman G. Simulation comes of age. *Adv Health Sci Educ Theory Pract* 2014;19:143-6.
 4. Cook DA, Hamstra SJ, Brydges R, et al. Comparative effectiveness of instructional design features in simulation-based education: systematic review and meta-analysis. *Med Teach* 2013;35:e867-98.
 5. McGaghie WC, Issenberg SB, Petrusa ER, Scalese RJ. A critical review of simulation-based medical education research: 2003-2009. *Med Educ* 2010;44:50-63.
 6. Bohlmann MK, Rath W. Medical prevention and treatment of postpartum hemorrhage: a comparison of different guidelines. *Arch Gynecol Obstet* 2014;289:555-67.
 7. GBD 2015 Maternal Mortality Collaborators. Global, regional, and national levels of maternal mortality, 1990–2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet* 2016;388:1775-812.
 8. Likis FE, Sathe NA, Morgans AK, et al. Management of postpartum hemorrhage. Comparative effectiveness review No. 151. . In: 15-EHC013-EF. PbtVE-bPCuCN---IAPN, ed. Rockville, MD: Agency for Healthcare Research and Quality; 2015.
 9. Kirkpatrick DL. Techniques for evaluating training programs. *Train Dev J* 1979:78-92.
 10. Kirkpatrick DL. Revisiting Kirkpatrick's Four-Level Model. *Train Dev J* 1996.
 11. Holton III EF. Holton's Evaluation Model: New evidence and construct elaborations. *Adv Dev Hum Resour* 2005;7:37-54.
 12. Holton III EF. The flawed Four Level Evaluation Model. *Hum Resour Dev Q* 1996;7:5-21.
 13. Yardley S, Dornan T. Kirkpatrick's levels and education 'evidence'. *Med Educ* 2012;46:97-106.
 14. Bates R. A critical analysis of evaluation practice: the Kirkpatrick model and the principle of beneficence. *Eval Program Plann* 2004;27:341-7.
 15. Cook DA. How much evidence does it take? A cumulative meta-analysis of outcomes of simulation-based education. *Med Educ* 2014;48:750-60.
 16. Cook DA. If you teach them, they will learn: why medical education needs comparative effectiveness research. *Adv Health Sci Educ Theory Pract* 2012;17:305-10.
 17. Bordage G. Conceptual frameworks to illuminate and magnify. *Med Educ* 2009;43:312-9.
 18. Todsén T, Henriksen MV, Kromann CB, Konge L, Eldrup J, Ringsted C. Short- and long-term transfer of urethral catheterization skills from simulation training to performance on patients. *BMC Medical Education* 2013.
 19. Cartier V, Inan C, Zingg W, Delhumeau C, Walder B, Savoldelli GL. Simulation-based medical education training improves short and long-term competency in, and

- knowledge of central venous catheter insertion: A before and after intervention study. *Eur J Anaesthesiol* 2016;33:568-74.
20. Dilaveri CA, Szostek JH, Wang AT, Cook DA. Simulation training for breast and pelvic physical examination: a systematic review and meta-analysis. *BJOG* 2013;120:1171-82.
 21. Diederich E, Mahnken JD, Rigler SK, Williamson TL, Tarver S, Sharpe MR. The effect of model fidelity on learning outcomes of a simulation-based education program for central venous catheter insertion. *Simul Healthc* 2015;10:360-7.
 22. Hossack T, Chris BB, Beer J, Thompson G. A cost-effective, easily reproducible, suprapubic catheter insertion simulation training model. *Urology* 2013;82:955-8.
 23. Mema B, Harris I. The barriers and facilitators to transfer of ultrasound-guided central venous line skills from simulation to practice: Exploring perceptions of learners and supervisors. *Teach Learn Med* 2016;28:115-24.
 24. Vadnais MA, Dodge LE, Awtrey CS, Ricciotti HA, Golen TH, Hacker MR. Assessment of long-term knowledge retention following single-day simulation training for uncommon but critical obstetrical events. *J Matern Fetal Neonatal Med* 2012;25:1640-5.
 25. Baldwin TT, Ford JK. Transfer of training: A review and directions for future research. *Pers Psychol* 1988;41:63-105.
 26. Isaranuwachai W, Brydges R, Carnahan H, Backstein D, Dubrowski A. Comparing the cost-effectiveness of simulation modalities: a case study of peripheral intravenous catheterization training. *Adv Health Sci Educ Theory Pract* 2014;19:219-32.
 27. Zendejas B, Brydges R, Wang AT, Cook DA. Patient outcomes in simulation-based medical education: A systematic review. *J Gen Intern Med* 2013;28:1078-89.
 28. Peterson DT, Watts PI, Epps CA, White ML. Simulation faculty development: a tiered approach. *Simul Healthc* 2017;12:254-9.
 29. Cheng A, Grant V, Dieckman P, Arora S, Robinson T, Eppich W. Faculty development for simulation programs: five issues for the future of debriefing training. *Simul Healthc* 2015;10:217-22.
 30. Meriën AER, van de Ven J, Mol BW, Houterman S, Oei SG. Multidisciplinary team training in a simulation setting for acute obstetric emergencies: a systematic review. *Obstet Gynecol* 2010;115:1021-31.
 31. Vandewaetere M, Manhaeve D, Aertgeerts B, Clarebout G, Van Merrienboer JJ, Roex A. 4C/ID in medical education: How to design an educational program based on whole-task learning: AMEE Guide No. 93. *Med Teach* 2015;37:4-20.
 32. Maggio LA, Cate OT, Irby DM, O'Brien BC. Designing evidence-based medicine training to optimize the transfer of skills from the classroom to clinical practice: applying the

- Four-Component Instructional Design Model. *Acad Med* 2015;90:1457-61.
33. Hecht L, Buhse S, Meyer G. Effectiveness of training in evidence-based medicine skills for healthcare professionals: a systematic review. *BMC Med Educ* 2016;16:103.
 34. de Bruin ABH, van Merriënboer JGG. Bridging Cognitive Load and Self-Regulated Learning research: A complementary approach to contemporary issues in educational research. *Learn Instr* 2017;51:1-9.
 35. van Merriënboer JGG, Kirschner PA. *Ten Steps to Complex Learning*. 3rd ed. New York, NY: Routledge; 2018.
 36. Kirschner PA, Sweller J, Clark RE. Why minimal guidance during instruction does not work: An analysis of the failure of constructivist, discovery, problem-based, experiential, and inquiry-based teaching. *Educ Psychol* 2006;41:75-86.
 37. Chiniara G, Cole G, Brisbin K, et al. Simulation in healthcare: a taxonomy and a conceptual framework for instructional design and media selection. *Med Teach* 2013;35:e1380-95.
 38. Paas F, Renkl A, Sweller J. Cognitive Load Theory and instructional design: Recent developments. *Educ Psychol* 2003;38:1-4.
 39. Fraser KL, Ayres P, Sweller J. Cognitive Load Theory for the design of medical simulations. *Simul Healthc* 2015;10:295-307.
 40. Eichbaum Q. Collaboration and teamwork in the health professions: rethinking the role of conflict. *Acad Med* 2017.
 41. Stocker M, Burmester M, Allen M. Optimisation of simulated team training through the application of learning theories: a debate for a conceptual framework. *BMC Med Educ* 2014;14:69.
 42. Arora S, Cox C, Davies S, et al. Towards the next frontier for simulation-based training: full-hospital simulation across the entire patient pathway. *Ann Surg* 2014;260:252-8.
 43. Salas E, Tannenbaum SI, Kraiger K, Smith-Jentsch KA. The science of training and development in organizations: What matters in practice. *Psychol Sci Public Interest* 2012;13:74-101.
 44. Grossman RS, E. The transfer of training: what really matters. *Int J Train Dev* 2011;15:103-20.
 45. Zendejas B, Wang AT, Brydges R, Hamstra SJ, Cook DA. Cost: The missing outcome in simulation-based medical education research - a systematic review. *Surgery* 2013;153:160-76.
 46. Owens DK, Qaseem A, Chou R, Shekelle P. High-Value, Cost-Conscious Health Care: concepts for clinicians to evaluate the benefits, harms, and costs of medical

- interventions. *Ann Intern Med* 2011;154:174-81.
47. Stammen LA, Stalmeijer RE, Paternotte E, Pool AO, Drissen EW, Scheele F, Stassen LPS. Training physicians to provide High-Value, Cost-Conscious Care: a systematic review. *JAMA* 2015;314:2384-2400.
 48. Ameh CA, van den Broek N. Making It Happen: Training health-care providers in emergency obstetric and newborn care. *Best Pract Res Clin Obstet Gynaecol* 2015;29:1077-91.
 49. Nelissen E, Ersdal H, Ostergaard D, et al. Helping Mothers Survive Bleeding After Birth: an evaluation of simulation-based training in a low-resource setting. *Acta Obstet Gynecol Scand* 2014;93:287-95.
 50. Szulewski A, Brindley P, van Merriënboer JG. Decision Making in Acute Care Medicine. In: Brindley PGC, P., ed. *Optimizing Crisis Resource Management to Improve Patient Safety and Team Performance A handbook for all acute care health professionals*. 1 ed: Royal College of Physicians and Surgeons of Canada; 2017:13-20.
 51. Hemming K, Haines TP, Chilton PJ, Girling AJ, Lilford RJ. The stepped wedge cluster randomised trial: Rationale, design, analysis, and reporting. *BMJ* 2015;350:h391.
 52. de Melo B, Amorim M, Katz L, Coutinho I, Figueiroa JN. Hypertension, pregnancy and weather: Is seasonality involved? *Rev Assoc Med Bras* 2014;60:105-10.

Summary

Chapter 1 consists of the thesis introduction, where we present the rationale that motivated our studies series. Simulation ubiquitous spread for the past decade was propelled by the ultimate goal of improving patient outcomes, while at the same time providing training opportunities for both novice and experienced personnel in a supervised “error safe” environment. However, more solid evidence with regards to its optimal training effectiveness is still being sought. Systematic reviews on simulation training effectiveness recommend the adoption of instructional design (ID) guidelines for designing such trainings. ID guidelines were developed with a basis on sound cognitive psychological theories of how people learn. They have a unique role in promoting complex learning, which is the integration and acquisition of knowledge, skills and attitudes along with the ability to properly coordinate the qualitatively different constituent skills. Two models summarize the steps described by the main ID guidelines: Merrill’s First Principles of Instruction and the Four-Component Instructional Design model. According to the guidelines expressed in those models, effective instruction should contain authentic, relevant, and daily real-world problems, emphasize practice at different levels of complexity, and provide feedback and diminishing instructional support through training. Postpartum hemorrhage (PPH) was the selected content for our studies series due to both its wide ecological validity for other high-risk situations and its epidemiological relevance. PPH remains the leading cause of maternal mortality in the world, with most resulting deaths considered as preventable for being due to failures in teamwork, communication and management issues. When considering training effectiveness, the widely adopted Kirkpatrick training evaluation model describes four different levels: learners’ satisfaction, learning outcomes, transfer from training to the workplace, and results. In our studies series, we aimed to explore the current use of ID guidelines by simulation training programs described in the literature and its impact on learning, transfer and patient outcomes (results).

In **Chapter 2**, we presented our analysis of the use of ID guidelines in simulation trainings for a high-risk situation, as described in the literature. We explored the adherence to such guidelines in PPH simulation training programs as described in articles that were retrieved from Pubmed, Eric, and Google Scholar between January 2007 and March 2017. Invited raters (N = 40) analyzed simulation training programs described in 32 articles, assessing their adherence to ID guidelines, based on Merrill's First Principles of Instruction, by using a 5-point Likert rating scale. This scale contained items related to key ID features categorized into five subscales: authenticity, activation of prior knowledge, demonstration, application, and integration/transfer. The articles were divided into subsets, which were distributed to the raters for analysis with a varying number of 7 to 10 reviewers per article subset. We then calculated the Likert-scale mean score, per subscale, and interrater reliability (IRR). We found a low level of adherence (< 3.00), with the IRR being excellent for the authenticity subscale, good-to-excellent for the integration/transfer subscale, fair-to-good for the activation of prior knowledge and application subscales, and poor for the demonstration subscale. In addition, we discussed the low level of adherence to evidence-based ID-guidelines in simulation training programs for high risk situations, such as PPH. We underscore the need to further promote faculty awareness with regards to the use of ID guidelines for simulation training in particular, when aiming to improve its effectiveness, especially, transfer of learning.

In **Chapter 3** we present a comparison between learning outcomes of two instructionally different PPH simulation training programs, one based on ID guidelines and the other on current best practices (BP). The presented pretest–post-test non-equivalent groups study was conducted among obstetrics and gynecology residents and included 13 teams; seven teams attended the ID based simulation and six the BP. Using a standardized task checklist, we compared the proportion of correctly executed tasks (post-test minus pretest). The presented “whole tasks” simulation trainings were planned to ideally manage a PPH condition and represent the core point of the subsequent studies. Our PPH ID-based simulation training adopted several elements from the four-component instructional design (4C/ID) model, with

overarching principles from other ID guidelines. As its first step, the problem was well designed – optimally managing PPH – and the reflection upon content relevance was promoted through a prior knowledge activation step with the OBGYN residents. As the following steps, the residents who attended the ID based PPH simulation training were exposed to well defined learning tasks, demonstration, variability of cases, increasing complexity, just-in-time information, supportive information, self-assessment, and debriefing sessions that promoted reflection on their own errors. Such a rich exposure to a variety of ID features aimed at promoting complex learning, a goal that was successfully reached. At post-test, the residents who attended the ID based PPH simulation training performed better (and were faster) in essential learning tasks and in specific subscales, results that reflect better development of cognitive schemata. In conclusion, in Chapter 3, we demonstrated that the use of simulation training for PPH based on evidence-based ID guidelines yielded better learning outcomes than did training based on best practice.

A long-term transfer analysis after such PPH simulation is presented in **Chapter 4**, where we show our qualitative study. Twelve OBGYN residents who attended one of the two PPH simulation trainings formats were interviewed using semi-structured interviews. In this Chapter, we also present the Baldwin and Ford (1988) training transfer model to explore such transfer. According to this model, three factors influence transfer: (1) trainee characteristics (the motivation to attend training, the perceived relevance of the training content, and self-confidence); (2) the training design (application of ID principles); and (3) the workplace environment (organizational feedback, team-work, and opportunity to use what has been learned). Our aim was to answer whether trainees perceived a long-term transfer effect two years after attending the simulation trainings. Using thematic analysis steps, we explored which training aspects trainees perceived as positively affecting this long-term transfer, which instructional design features facilitated and/or hindered long-term transfer, and whether there were differences in the participants' perceptions according to the type of simulation attended. We analyzed how the residents attending both training formats perceived positive long-term transfer two years after having attended simulation training for PPH. They acknowledge

PPH as a relevant training topic and reported being motivated to attend the simulations as a desire to improve professional skills and an increase in self-confidence after the training. Attendees of either simulation format reported the following ID elements as key contributors to transfer: authenticity, use of Paivio's dual-coding strategy (a combination of text and pictures that makes information easier to remember), and feedback during debriefing. Residents who participated in the ID-based simulation additionally mentioned variability and increasing complexity as ID elements contributing to transfer. Residents from both groups reported an increase in teamwork skills, but also appointed poor organizational feedback and team struggles with colleagues who had not attended the simulation. Residents who had attended the ID format perceived better transfer of communication and teamwork skills and higher situational awareness than those having attended the BP format. In conclusion, Chapter 4 presents how long-term transfer was perceived by the residents; however, a systematic approach to problem solving was perceived only by those who attended the ID based PPH simulation training. In addition, trainees who participated in the ID-based simulation perceived better communication skills and better overall situational awareness: "I didn't do that before." Such findings may reflect better long-term situational awareness – an ability to anticipate, recognize, and intercept unfolding error chains, a skill that is of extreme relevance in the adequate management of several high-risk clinical conditions, such as PPH.

In Chapter 5 we presented an uncontrolled before-and-after study that analyzed the impact on patient outcomes at a teaching hospital, after the OBGYN residents attended the ID-based PPH simulation training programme. Data from all women delivering at the maternity of this teaching hospital one year before the simulation were compared to those of all women delivering after the simulation was performed. To minimize seasonality biases, the same three months of the year (June to August) were used for the comparison. We described the contribution of well-defined patient outcome measures whose analysis went beyond single procedural outcomes as usually described in the literature. We presented a detailed analysis of specific and relevant PPH outcomes measures. Our main outcome measures were therapeutic

uterotonics usage rates within 24h of birth (TUW24h) and blood transfusion rates (BTR), while our secondary outcome measures included uterotonic use rates and dosages for each drug, postpartum Hb <6g/dL, B-Lynch suture, uterine artery embolization, hysterectomy, renal insufficiency, obstetric intensive care unit (ICU) admission, mechanical ventilation, length of hospital stay, near-miss maternal mortality and maternal death. The results presented in Chapter 5 correspond to 1388 women delivering in the analyzed months before the training and 1357 in the following year. The PPH rates were 100 cases (7.2% of deliveries) and 80 cases (5.9% of deliveries), respectively. No significant differences were found in the comparison between primary post- and pre-simulation outcomes, although an increase was identified in the therapeutic oxytocin mean dosage IU within 24h of birth. For all other outcome measures, there were no statistical differences. Such increased oxytocin mean dosage in fewer PPH patients may in fact reflect improved ongoing vigilance of the bleeding woman after simulation, since such dosages are administered on a progressive and continuous basis. The study presented in Chapter 5 included only OBGYN residents and was conducted in a single center. Future studies should not only include other specialties, aiming at the improvement of teamwork skills, but also be designed in a multicenter stepped-wedged approach (random and sequential crossover of clusters from control to intervention, until all clusters provide the simulation). Therefore, the verified increase in the therapeutic oxytocin mean dosage in fewer and selected PPH cases may indicate better situational awareness when managing women with PPH after an ID-based simulation training.

Chapter 6 is dedicated to the discussion of our thesis, where we present a summary of the findings of our studies series and reflect on their theoretical implications. We discuss our findings on the use of ID guidelines for simulation by previously published articles, which ranges from a concerning low adherence to such guidelines to a wide range of positive uses of ID guidelines for PPH simulation training. We also highlight our thesis strengths, such as the sound analysis of the Kirkpatrick training evaluation model levels, a detailed task analysis for desired learning outcomes, a long-term transfer analysis, and detailed patient outcomes measures. We also reflect upon a potential limitation,

in particular, the single center patient outcomes analysis. We recommend future studies including multiple centers in a stepped wedged approach and including multiple specialties for adequately training teamwork. We conclude our findings by reinforcing the recommendation from systematic reviews that ID guidelines must be used when designing simulation training. This PhD thesis also underscores the importance of improving faculty awareness with regards to this topic as an essential strategy for the improvement of patient outcomes through simulation training.

Samenvatting

Hoofdstuk 1 vormt de inleiding tot dit proefschrift en legt uit waarom de hier gepresenteerde onderzoeken zijn uitgevoerd. De afgelopen tien jaar hebben simulaties een hoge vlucht genomen in het medisch onderwijs. De oorzaak hiervan is het streven naar betere patientuitkomsten en de mogelijkheden die simulaties bieden om zowel beginnende als ervaren beroepsbeoefenaars te trainen in een gesuperviseerde, 'foutenvrije' omgeving. Hoe effectief simulaties zijn voor dergelijke trainingsdoeleinden is echter nog onvoldoende onderzocht. Systematische reviews van de effectiviteit van simulatietraining bevelen aan onderwijskundige ontwerprichtlijnen (Instructional Design of ID richtlijnen) te gebruiken bij het ontwerpen van deze trainingen. Zulke richtlijnen zijn ontwikkeld op basis van solide theorieën uit de cognitieve psychologie over de manier waarop mensen leren. Ze spelen een unieke rol bij het stimuleren van complex leren: de verwerving en integratie van kennis, vaardigheden en attitudes in combinatie met het vermogen de kwalitatief verschillende deelvaardigheden goed op elkaar af te stemmen. De stappen die onderdeel uitmaken van de voornaamste ID richtlijnen zijn samengevat in twee modellen: Merrill's First Principles of Instruction en het Four-Component Instructional Design (4C/ID) model. Volgens de in deze modellen beschreven richtlijnen heeft effectieve instructie de volgende elementen: er wordt gewerkt met authentieke, relevante en alledaagse problemen, er wordt geoefend op diverse complexiteitsniveaus, er wordt feedback gegeven, en de ondersteuning door opleiders wordt tijdens de training gaandeweg afgebouwd. Voor de hier beschreven onderzoeken richtten we ons op haemorrhagia post partum (HPP), aangezien deze aandoening een hoge ecologische validiteit heeft voor andere hoog risico-aandoeningen en vanwege de epidemiologische relevantie ervan. HPP is nog steeds de voornaamste oorzaak van moedersterfte wereldwijd. In de meeste gevallen had de fatale afloop vermeden kunnen worden, aangezien deze vaak veroorzaakt wordt door slechte samenwerking, communicatie en beleid. Voor

wat betreft de effectiviteit van training, geeft het algemeen geaccepteerde trainingsevaluatiemodel van Kirkpatrick vier niveaus aan: tevredenheid van leerders, leeruitkomsten, transfer van de trainingssituatie naar de werkvloer en resultaten. Voor dit proefschrift onderzochten we het gebruik van ID-richtlijnen bij simulatietrainingsprogramma's zoals beschreven in de literatuur en het effect ervan op leren, transfer en patientuitkomsten.

In **hoofdstuk 2** presenteren we onze analyse van het gebruik van ID-richtlijnen in simulatietrainingen voor hoog risico-situaties zoals beschreven in de literatuur. In artikelen gepubliceerd tussen januari 2007 en maart 2017 op Pubmed, Eric en Google Scholar gingen we na hoe zulke richtlijnen in HPP simulatietrainingsprogramma's opgevolgd worden. De door ons ingeschakelde beoordelaars (N = 40) analyseerden de simulatietrainingsprogramma's beschreven in 32 artikelen. Op basis van Merrill's First Principles of Instruction beoordeelden ze de mate waarin de ID-richtlijnen toegepast werden. Het oordeel werd uitgedrukt op een 5-punts Likertschaal. Deze schaal omvat items die betrekking hebben op fundamentele ID-kenmerken verdeeld over vijf subschalen: authenticiteit, activatie van reeds aanwezige kennis, demonstratie, toepassing en integratie/overdracht. De artikelen werden verdeeld in groepen die werden toegewezen aan de beoordelaars (zeven tot tien beoordelaars per groep). Daarna werden de gemiddelde score voor de gehele schaal, de gemiddelde scores per subschaal en de interbeoordelaarsbetrouwbaarheid (IRR) berekend. De mate waarin de programma's de richtlijnen opvolgen bleek laag te zijn (< 3,00), met een zeer goede IRR voor de subschaal authenticiteit, een goede tot zeer goede IRR voor de subschaal integratie/overdracht, een redelijk tot goede IRR voor de subschaal activatie van reeds aanwezige kennis en een slechte IRR voor de subschaal demonstratie. Verder bespreken we de geringe mate waarin de evidence-based ID-richtlijnen worden opgevolgd in simulatietrainingsprogramma's voor hoog risico-situaties zoals HPP. We benadrukken de noodzaak om opleiders meer bewust te maken van het gebruik van ID-richtlijnen, vooral bij simulatietrainingen, en met name om de effectiviteit van de overdracht van het geleerde te verbeteren.

In **hoofdstuk 3** presenteren we een vergelijking tussen leeruitkomsten van twee HPP-simulatietrainingsprogramma's, waarbij de ene gebaseerd was

op ID-richtlijnen en de andere op de huidige best practices (BP). Dit pretest-posttest niet-equivalente groepenonderzoek werd verricht onder arts-assistenten verloskunde en gynaecologie en omvatte 13 teams. Zeven teams volgden de ID-simulatie en zes de BP-simulatie. Met een gestandaardiseerde takenchecklist vergeleken we het percentage correct uitgevoerde taken (posttest min pretest). De gepresenteerde “gehele taken” simulatietrainingen waren gericht op een optimale behandeling van HPP en vertegenwoordigen de belangrijkste taakaspecten zoals geïdentificeerd in eerdere onderzoeken. Onze op ID-richtlijnen gebaseerde HPP-training omvatte diverse elementen van het vier-componenten onderwijsontwerpmodel (4C/ID) en de voornaamste principes van andere ID-richtlijnen. Als eerste stap werd het probleem – optimale HPP-behandeling – goed gedefinieerd. In de volgende stap, waarin reeds aanwezige kennis geactiveerd werd, werden de arts-assistenten gestimuleerd te reflecteren op relevante aspecten. In de daaropvolgende stappen werden de arts-assistenten die deelnamen aan de ID-training geconfronteerd met goed gedefinieerde leertaken, hands-on demonstraties, een scala aan casussen van oplopende complexiteit, informatie die net op tijd beschikbaar komt, ondersteunende informatie, zelfbeoordeling en nabesprekingen om hen te stimuleren te reflecteren op hun fouten. Een dergelijke rijke confrontatie met veel ID-kenmerken heeft als doel complex leren te bevorderen. Dit doel werd bereikt. De arts-assistenten die de ID-training hadden gevolgd behaalden hogere scores op de posttest (en waren sneller) op realistische leertaken en op specifieke subschalen, hetgeen erop wijst dat hun cognitieve schemata beter ontwikkeld waren. Ten slotte laat hoofdstuk 3 zien dat het gebruik van een HPP-simulatietraining gebaseerd op evidence-based ID-richtlijnen betere leeruitkomsten oplevert dan een training gebaseerd op best practices.

In **hoofdstuk 4** wordt een kwalitatieve analyse gepresenteerd van langetermijntransfer na een dergelijke HPP-training. Twaalf arts-assistenten verloskunde en gynaecologie die een van de twee HPP-trainingen hadden gevolgd werden ondervraagd met behulp van semi-gestructureerde interviews. In dit hoofdstuk presenteren we ook het Training Transfer Model van Baldwin en Ford (1988) om de transfer van training te onderzoeken. Volgens dit model wordt transfer beïnvloed door drie factoren: (1) deelnemerskenmerken: motivatie om de training bij te wonen, veronderstelde

relevantie van de inhoud van de training en zelfvertrouwen; (2) ontwerp van de training: toepassing van ID-principes; en (3) werkomgeving: feedback vanuit de organisatie, teamwerk en mogelijkheden om het geleerde te gebruiken. Doel was om te zien of de deelnemers vonden dat langetermijntransfer had plaatsgevonden in de twee jaar na afloop van de HPP-training. Met behulp van thematische analysestappen onderzochten we welke aspecten van de training de deelnemers zagen als positieve bijdragen aan deze langetermijntransfer, welke onderwijskundige ontwerpkenmerken de transfer makkelijker of moeilijker maakten, en of de deelnemers aan de verschillende typen simulatie andere ervaringen hadden. We analyseerden wat de arts-assistenten die een van beide trainingen hadden gevolgd vonden van de langetermijntransfer twee jaar nadat ze deze hadden ondergaan. De deelnemers stelden dat HPP een relevant onderwerp is voor een training en dat ze gemotiveerd waren om de simulaties bij te wonen om zo hun professionele vaardigheden te verbeteren, en dat hun zelfvertrouwen was toegenomen na de training. De deelnemers vonden dat de volgende ID-kenmerken het meest bijdroegen aan transfer: authenticiteit, het gebruik van Paivio's dual-coding strategy (een combinatie van tekst en beelden die het makkelijker maakt informatie te onthouden) en feedback tijdens de nabespreking. De arts-assistenten die deelnamen aan de ID-training noemden ook variabiliteit en toenemende complexiteit als ID-elementen die bijdragen aan transfer. Arts-assistenten in beide groepen vonden dat hun samenwerkingsvaardigheden waren verbeterd, maar gaven ook aan dat de feedback vanuit de organisatie soms slecht was en dat er problemen waren met collega's in het team die de training niet hadden gevolgd. De communicatieve en samenwerkingsvaardigheden van degenen die de ID-training hadden gevolgd waren meer verbeterd en hun 'situational awareness' was hoger dan van degenen die de BP-training hadden gevolgd. Ten slotte laat hoofdstuk 4 zien hoe de arts-assistenten aankeken tegen de transfer op de lange termijn. Alleen degenen die de ID-training hadden gevolgd vonden dat ze een systematische aanpak om problemen op te lossen hadden geleerd. Bovendien vonden deze deelnemers dat ze betere communicatieve vaardigheden hadden en de klinische situatie beter konden overzien: "Dit deed ik vroeger niet." Dergelijke resultaten kunnen erop wijzen dat ze een betere en meeromvattende 'situational awareness' hebben: het vermogen om te anticiperen, situaties te herkennen en eerder in te grijpen

wanneer cumulatie van fouten dreigt. Het laatste is een vaardigheid die zeer belangrijk is om hoog-risico aandoeningen, waaronder HPP, te behandelen.

In **hoofdstuk 5** presenteren we een pretest-postteststudie zonder controlegroep waarin we het effect van training op patientuitkomsten nagingen in een opleidingsziekenhuis waar de arts-assistenten verloskunde en gynaecologie hadden deelgenomen aan de ID-simulatietraining voor HPP. Gegevens van alle vrouwen die een jaar vóór de training op de kraamafdeling van dit ziekenhuis een baby ter wereld brachten werden vergeleken met gegevens van alle vrouwen die een jaar na de training bevielen. Om seizoensinvloeden te minimaliseren werden dezelfde drie maanden gebruikt (juni t/m augustus). We beschrijven het effect van de training op goed-gedefinieerde patientuitkomsten waarvan de analyse verder ging dan afzonderlijke procedureresultaten zoals gewoonlijk beschreven in de literatuur en presenteren een gedetailleerde analyse van specifieke en relevante HPP uitkomstmaten. De belangrijkste uitkomstmaten waren het therapeutisch gebruik van uterotonica in de eerste 24 uur na de geboorte (TUW24h) en de mate waarin bloedtransfusies gegeven werden (BTR). Onze secundaire uitkomstmaten waren onder andere de mate waarin uterotonica werden gegeven en de doses van elk middel, post-partum Hb lager dan 6 g/dL, toepassing van B-Lynch hechtingen, embolisatie van de a. uterina, uterusextirpatie, nierfalen, opname op de verloskundige Intensive Care-afdeling, mechanische ventilatie, opnameduur, bijna fatale afloop voor de moeder en dood van de moeder. De resultaten die in dit hoofdstuk gepresenteerd worden hebben betrekking op 1388 vrouwen die in een periode van drie maanden vóór de training een baby ter wereld brachten en 1357 vrouwen die in dezelfde periode van drie maanden in het jaar daarna een baby ter wereld brachten. Het aantal HPP-gevallen in beide groepen was respectievelijk 100 (7,2% van de bevallingen) en 80 (5,9%). Er werden geen significante verschillen in primaire uitkomstmaten gevonden in beide groepen. Wel werd een toename geconstateerd in de gemiddelde therapeutische oxytocinedosis toegediend binnen 24 uur na de bevalling. Voor alle andere uitkomstmaten werden geen statistische verschillen gevonden. Een dergelijke toegenomen gemiddelde oxytocinedosis bij minder HPP-patiënten kan wijzen op een meer intensieve bewaking van het bloedverlies na de simulatie, aangezien oxytocine progressief en continu wordt toegediend. Het onderzoek

dat in dit hoofdstuk wordt gepresenteerd, omvatte alleen arts-assistenten verloskunde en gynaecologie en werd in één ziekenhuis verricht. Nader onderzoek zou niet alleen andere specialismen moeten omvatten en zich moeten richten op de verbetering van samenwerkingsvaardigheden maar ook meerdere ziekenhuizen moeten omvatten en een 'stepped-wedge' ontwerp moeten hebben (gerandomiseerde en sequentiële uitwisseling van clusters van de controle- naar de interventiegroep, totdat alle clusters de simulatie aanbieden). Aldus kan de waargenomen toename van de gemiddelde dosis oxytocine bij minder en geselecteerde HPP-patiënten wijzen op een betere 'situational awareness' onder behandelaars van vrouwen met HPP na het volgen van een ID-simulatietraining.

In **hoofdstuk 6** worden de hier gepresenteerde studies samengevat en gaan we in op de theoretische implicaties van de resultaten. We bespreken onze bevindingen met het gebruik van ID-richtlijnen voor het ontwerpen van simulatietrainingen zoals beschreven in de literatuur, die uiteenlopen van een zorgwekkende veronachtzaming van dergelijke richtlijnen tot een scala aan manieren om ID-richtlijnen op positieve wijze te gebruiken voor HPP-simulatietrainingen. We geven ook de sterke punten van ons onderzoek weer, zoals de solide analyse van de niveaus van het Kirkpatrick trainingsevaluatiemodel, de gedetailleerde taakanalyse om te komen tot gewenste leeruitkomsten, de analyse van de langetermijntransfer en de gedetailleerde maatregelen ter verbetering van patientuitkomsten. We bespreken ook de mogelijke beperkingen, in het bijzonder het feit dat de patientuitkomsten van slechts een ziekenhuis zijn geanalyseerd. Nader onderzoek zal in meerdere ziekenhuizen moeten plaatsvinden, een 'stepped-wedge' onderzoeksontwerp moeten hebben en meerdere specialismen moeten omvatten om te komen tot een goede training van samenwerkingsvaardigheden. We eindigen met een herhaling van de aanbeveling die naar voren kwam uit de systematische reviews: dat bij het ontwerp van simulatie-trainingen gebruik moet worden gemaakt van ID-richtlijnen. Dit proefschrift benadrukt ook hoe belangrijk het is om opleiders meer bewust te maken van dit thema, als een essentiële strategie voor het verbeteren van patientuitkomsten door middel van simulatie-trainingen.

Valorization

This Chapter meets the requirements of Article 22 of the Regulation Governing the Attainment of Doctoral Degrees on the inclusion of a valorization paragraph in all PhD dissertations at Maastricht University as of 1 September 2014. Knowledge valorization is the “process of creating value from knowledge by making knowledge suitable for translation into competitive products, services, processes and new commercial activities”.

Relevance. The studies presented in this thesis demonstrate positive training outcomes of an instructional design based obstetrics simulation training. The practical implications of these findings may be found on several fronts – from responding to demands on continuous healthcare education to positive impacts on patient outcomes.

The need for evidence-based simulation training which makes good use of instructional design principles emanated from the high budget demands of this training strategy. After initial euphoria, simulation training needed to be optimized and its effectiveness needed to be soundly proved¹. This thesis may then contribute to this aim by presenting an effective simulation training format that can be potentially applied to multiple different training contents.

The presented studies explored the outcomes of instructional design based simulation training and the selected training content was postpartum hemorrhage (PPH), the leading cause of maternal deaths worldwide². Maternal mortality rates remain unacceptably high, despite the verified global improvement over the past decades, especially because the majority of cases are considered preventable. Management failures frequently associated with high mortality rates are described as miscommunications, teamwork malfunction, and lack of situational awareness^{3,4}. Therefore, the positive training outcomes of the explored simulation training may span across several domains beyond PPH clinical management itself.

This thesis aimed to contribute to demands from the international scientific community within a global perspective. However, presenting the

local context at which the studies were conducted may further contribute to reflections upon their relevance. The studies were conducted in the city of Recife, located in the Northeast region of Brazil, a middle-income country. The Northeast region holds 30% of the Brazilian population and accounts for 13% of Brazil's GDP. Its wide socio-economical inequalities lead to healthcare assistance challenges varying from those encountered by high-income countries to those from low-income countries. In spite of improvements on maternal mortality rates, women still die at high rates worldwide, but foremost in low-and-middle-income countries (LMIC).

Brazilian healthcare coverage is free and universal since the creation of Brazil's Sistema Único de Saúde (SUS) by the 1988 Brazilian Constitution. Along with healthcare assistance among SUS attributions lays the planning and provision of healthcare professionals continuous education⁵. Therefore, healthcare policy makers and healthcare community are continuously in search for effective healthcare personnel training and the findings from this thesis may be largely welcomed in benefit of maternal healthcare assistance.

In addition to the challenges encountered by maternal healthcare professionals worldwide, those in LMIC face a daily stressful routine of work overload and personnel shortage. Such reality leads to undesirable neglect to quality assistance and patient safety compromise. However, improving patient safety and quality of care may be achieved through effective personnel training and does not necessarily demand prohibitively expensive investments. By providing effective simulation training, a positive impact may be perceived in several key perspectives: from avoiding patient harm in its individual consequences and societal economical burden to introducing a safe learning environment and a culture in which errors are viewed as an opportunity for learning⁶.

Effective strategies contributing to the reduction of maternal mortality benefit society as a whole. In societies economically still under development and with fragile institutions, the absence of a mother brings even further significant compromise to the offspring in various different aspects (vaccination rates completion, neurocognitive development and academic outcomes)². As long as the goal of safe pregnancy and childbirth for all women and all

girls in all countries is still not reached, research and innovation are essential ingredients for success⁷. Therefore, providing effective evidence based postpartum hemorrhage simulation training may significantly contribute to improve our current prospect.

Target groups. Our findings on effective evidence-based simulation training using instructional design principles may benefit healthcare professionals who are in constant need of updating their knowledge, skills and overall competencies. Healthcare professionals are constantly facing changes in an evolving society, from technological advances to interpersonal interactions. Their continuous demands for complex learning and proper integration of a wide range of training contents may find an effective response strategy in simulation training. In addition, such training strategy may also contribute to their need to self-regulate the deep foundations of the psychological sciences of clinical reasoning^{8,9}.

Overall, healthcare policy makers and stakeholders may also benefit from our presented findings. For those in charge of providing strategies for healthcare professionals to succeed in their continuous search for up-to-date knowledge and skills becoming aware of effective simulation training designs that are easily applicable and reproducible is a helping hand.

Activities/Products/Schedule & Implementation. As previously described, the studies reported in this thesis were conducted in Recife, a city located in the impoverished Northeast region of Brazil. Therefore, it represents a large potential to contribute to the local healthcare system. Our aim is to offer instructional design based simulation trainings in a wide range of contents to healthcare professionals across clinical disciplines.

As a first step, the presented PPH training will be replicated at the Maternity where the studies were conducted, Instituto de Medicina Integral Prof. Fernando Figueira (IMIP), to incoming residents and healthcare teams. Then, the training will be amplified to other epidemiological relevant contents such as eclampsia and sepsis. In sequence, it should be further offered to other healthcare professionals and residents from other disciplines, to further promote teamwork interactions. Later, rare-event contents will be also incorporated and offered as simulation training for the multidisciplinary healthcare teams.

In a longer-term perspective, the plan is to offer effective simulation training to healthcare professionals outside the institution. A simulation training and innovation research center is currently being planned and developed to accommodate training routine to a wider audience through an agreement with healthcare policy makers. The schedule for inauguration is predicted for late 2019.

Innovation. The simulation training format explored in the reported studies provided opportunity for repetitive practice through a variation of multiple clinical scenarios presented in a simple-to-complex sequence. Multiple debriefing opportunities were also provided. Such design led to significant positive learning outcomes, long-term transfer and a positive impact on patient outcomes. These findings and design represent a long-awaited innovation at the current frontier of knowledge in healthcare simulation training research and their benefits were soundly demonstrated^{10,11}.

References

1. Draycott T. Not all training for obstetric emergencies is equal, or effective. *BJOG* 2017;124:651.
2. Graham W, Woodd S, Byass P, et al. Diversity and divergence: the dynamic burden of poor maternal health. *Lancet* 2016;388:2164-2175.
3. Knight M, Kenyon S, Brocklehurst P, et al. Saving lives, improving mothers' care – lessons learned to inform future maternity care from the UK and Ireland confidential enquiries into maternal deaths and morbidity 2009–12. Oxford, UK2014.
4. Miller S, Abalos E, Chamillard M, et al. Beyond too little, too late and too much, too soon: a pathway towards evidence-based, respectful maternity care worldwide. *Lancet* 2016;388:2176-2192.
5. Batista K, Gonçalves O. Formação dos Profissionais de Saúde para o SUS: significado e cuidado. *Saúde Soc* 2011;20:884-889.
6. Flott K, Fontana G, Dhingra-Kumar N, Yu A, Durkin M, Darzi A. *Lancet* 2017;389.
7. Depoortere E, Matthews Z, Nove A, et al. The need for innovation and implementation research for maternal and newborn health. *Lancet* 2016;388:1374.
8. Coiera E. The fate of medicine in the time of AI. *Lancet* 2018.

9. de Bruin ABH, van Merriënboer JJG. Bridging Cognitive Load and Self-Regulated Learning Research: A complementary approach to contemporary issues in educational research. *Learn Instr* 2017;51:1-9.
10. McGaghie WC, Issenberg SB, Petrusa ER, Scalese RJ. A critical review of simulation-based medical education research: 2003-2009. *Med Educ* 2010;44:50-63.
11. Cook DA, Hamstra SJ, Brydges R, et al. Comparative effectiveness of instructional design features in simulation-based education: Systematic review and meta-analysis. *Med Teach* 2012;35:e867-e898.

Acknowledgment

Writing a thank you note is always challenging. You start planning on keeping it short, but then you recall all the relevant support you received during the PhD journey and become convinced that they should be acknowledged. You end up on a retrospective journey of reflecting which paths led you this way. Unavoidably, the list becomes too long, but all names were recalled with love and gratitude. At the end, some may be omitted, but the gratefulness will always be present. These are the people I would like to honestly say thank you to.

This thesis is a result of a collective project made possible through the support of the Instituto de Medicina Integral Prof. Fernando Figueira (IMIP) and Faculdade Pernambucana de Saúde (FPS). It is aligned with the ideas and principles promoted by **Prof. Fernando Figueira** since IMIP's inception "Always there for the people, working for a better world". Living in the impoverished Northeast region of Brazil, this sentence makes more than sense. My special thanks to **Gilliat Falbo** for inviting me and giving me a vote of confidence for the challenge. I thank you for being a lighthouse since my early professional years. I would also like to thank **Antonio Carlos Figueira** and **Carlos Santos da Figueira** for making this PhD possible and **Fernando Menezes** for articulating the partnership between IMIP/FPS and the School of Health Professions (SHE) at Maastricht University. IMIP is my professional home, where I attended my Obstetrics and Gynecology residency and my Masters in Maternal and Child Healthcare.

To **Ana Falbo** for all the support, in all times, when most needed. I once described her as a Henderson–Hasselbalch equation. You must see her adapt herself, yourself and the environment, and you will understand. Such characteristic could not be more appropriated and effective. My honest thank you.

To **Cees Van der Vleuten** for being present in my very early steps in SHE supportive research environment, since we first met, in Recife. You realistically presented the high standard of support and guidance provided to those who would be admitted into SHE PhD program. I can only be thankful for receiving such support and much more from SHE. The safe learning and stimulating research SHE environment will be a constant and happy recollection. I am also very thankful for all the wise guidance and great expertise in promptly recognizing my current and next puzzle/struggle and immediately showing me the way forward.

To **Jeroen van Merriënboer** for such careful, detailed, consistent and patient supervision. Each and every time, when needed. “Take the reader by the hand”, not only told, but provided by example, became my mantra.

To both of you, **Cees and Jeroen**, together, as a team. I would always be certain of replies to my emails, in the best and briefest way. At each Skype meeting, a perfect combination of supervision profiles, always looking ahead, pushing me forward and facing the bright side of the story. And last but not least, I am aware of the privilege to chat over science and research struggles at each month with such experienced researchers. All of this while sharing the fundamental stepping stones on the content of my thesis. Having this as a routine for the past years was certainly a privilege. I sure learned much from you. Thank you very much.

To **all team from SHE**, always supportive and friendly.

To **all colleagues and friends** from all different departments at **IMIP** for all the support. To those from the **Centro de Atenção à Mulher-CAM**, all healthcare team colleagues OBGYN, Neonatologists, Anesthesiologists, Surgeons, Obstetric Intensive Care, nursing staff, nutrition, cleaning and transport. I would particularly like to acknowledge the support from our CAM Coordinator **Adriana Scavuzzi** and also **Flavia Anchielle** in the most critical times. To my dear High-Risk Obstetrics Ward team from 4CAM (**Ana**

Porto and **Isabela**, you are missed everyday) in fundamental guidance in many fronts and **Gláucia Guerra** for her daily support, both personally and professionally. Also, to our dear 4CAM fellows in high risk obstetrics through these PhD years: **Carol Del Mar**, **Juliana Silveira**, **Gabriela Melo** and **Liana Lira**, **Judith Pontual** and **Gabriela Correia** and our lovely nursing and support staff – **Sandra Machado**, you rule! To those who were available to support me in my needed substitutions, my special thanks to **Djenane** and **all Delivery Room team** – thank you for every shared horray! To those whom I shared the daily struggle: **Colegiado do CAM; UTI Obstétrica; Triagem e Sala de Parto; Diretoria de Pesquisa; Diretoria de Pós-graduação; Centro Acadêmico; Núcleo de TeleSaúde-NTES; Educação à Distância-EAD; Presidência** and **Superintendência** for all institutional support.

To all who shared the struggle of PhD life work balance: **Nara Cavalcanti**, **Natasha Calheiros**, **Hegla Prado**, **Manuela Hazin**, **Tereza Rebeca**, **Karina Melo** and **Paula Diniz**.

To all colleagues from **FPS**, from the early talk with **Pacheco** to the current sharing with **Taciana Duque**, **Edvaldo Souza**, **Patricia Bezerra**, **Cristiano Souza**, **Suênia França**, **Flavia Orange**, **Carla Araújo** and **Luciano Guerra**. In special, thank you to those who shared the SHE/Maastricht experience and gave me support along the way: **Antonio Cavalcanti**, **Hegla Prado**, **Mirela Rebêlo**, **Tereza Rebeca**, **Emanuelle Valente**, **Taciana Albuquerque**, **Talita Menezes**, **Rafael Oliveira** and **Alexandre Sales**.

All colleagues from the other our **High-Risk Maternities in Pernambuco** – **Centro Integrado Amaury de Medeiros-CISAM**, **Hospital das Clínicas-HC**, **Hospital Barão de Lucena-HBL**, **Hospital Agamenon Magalhães-HAM** – who struggle with the daily challenge of our local obstetric assistance. In particular, to the team from **HC**, specially **Dr Antonio Carlos Barbosa** and **Elias Melo** for supporting me while conducting the PhD and participating in the study.

To all my colleagues whom I have met through my professional journey

and supported me by their actions and/or words or by sharing their passions for knowledge and applying the best available knowledge to women in need. In particular, those who took their time to discuss my PhD project in its early stages, **Melania Amorim, Leila Katz, Isabela Coutinho, Guilherme Cecatti, João Paulo Souza.**

I would specially like to thank **Unidade de Cardiologia e Medicina Fetal-UCMF** for all the learning, support and example of “When there is a will, there is a way” as a student and in my early professional years. **Sandra Mattos and team**, your example of learning, transfer of learning and positive impact on children with cardiopathy should be always congratulated. I am proud of being part of the team.

All of those who joined me at the **Diretório Acadêmico Umberto Câmara – DAMUC** throughout MedSchool, for sharing the vision of making a better world possible starting by challenging the teaching standards at the time. Being part of DAMUC and participating in the **Direção Executiva Nacional dos Estudantes de Medicina-DENEM** meetings was my first exposure to the topic of Medical Education, active learning and Problem Based Learning.

To the **Serviço de Cirurgia Geral e Transplante Hepático at the Universidade de Pernambuco-UPE**, where I attended my General Surgery Residency. The unit praise for quality of care is a role model for those who work at the Sistema Único de Saúde-SUS.

To all who collaborated as experts in instruction and/or simulation and contributed to the pilot stage of the rating scale for the study presented in Chapter 2: David Merrill, Jake Einfeld, Bert Hoogveld, Jung Lim, Irene Tijam, Jeroen Van Merrienboer and Gilliat Falbo. And I can only be more than thankful **to all healthcare experts who contributed as raters** by analyzing the simulation training programs: Angela Hewett, AnnSofia Thomsen, Carina Georg, Carla Polido, Caroline Ong, Cecilia Nilsson, Cris Attoe, Ebbe Thinggaard, Edvaldo Souza, Elaine Rossi, Elizabeth Santos, Heather

Straub, Ieda Aleluia, Jeffrey Cheng, Jette Sorensen, Jos van Roosmalen, Kristopherson Lustosa, Lawrence Grierson, Leigh Kinsman, Ligia Ribeiro, Louis Boyer, Marcia Costa, Marco Carvalho Filho, Marleen Gronier, Melanie Chichester, Patricia Bezerra, Patricia Cury, Roxana Knobel, Ruth Endacott, Sanne Peters, Sharon Maslovitz, Signe Egenberg, Simon Cooper, Steven Andersen, Tereza Rebecca Lima, Thiago Appoloni, Tia Andrigueti, Tiago Grangeia, Tobias Todsén, Yoshikazu Asada.

For the learning outcome study, presented in Chapter 3, I would like to thank **Leila Katz** and **Isabela Coutinho** for all the expertise, collaboration and support in the detailed planning of the scripts of the scenarios. To **ProDelphus** and **Marcos Lyra** and team for all the support in designing the postpartum hemorrhage part-task simulator. **To all who attended the trainings:** residents, supervising obstetricians, video raters, and simulated nurses: Leila Katz, Luciana Schuler, Cinthia Komuro, Carol Del Mar, Djenane Siqueira, Cinthia Brito, Sheila Leoni, Dirce Santos, Edilberto Rocha, Danielle Gattás, Valeska, Pricila, Layane Alves, Jizany Tavares, Juliana Carvalho, Socorro. My special thanks to the volunteer simulated patient who contributed out of friendship and in the belief of contributing to obstetrical healthcare assistance: **Vanessa Bahé, Paula Bonfim, Nélia de Paula**, thank you so much.

For their contribution in the long-term transfer of learning study, presented in Chapter 4, I would like to thank **Jette Led Sorensen** and **Ana Claudia Conrado**.

To all women whose data were analyzed in Chapter 5.

To the learning culture of my parents, Celso and Lucia, who inherited it from their parents. As a family of teachers, knowledge was always valued, as well as the importance of sharing it. Thank you, **Celso, Lucia** and **Mila** for all the constant support, incentive and brainstorming.

To my partner and best friend **Juliano** for the daily support and constant

stimuli. Always looking at the half full part of the glass was essential. And for sharing the daily upbringing of our reason for all, **Bernardo**.

To all my friends who supported me, each in their own special way, with a word of support when needed. **Luciana Schuler** and **Cinthia Komuro**. **Leila, Isabela, Melania, Ana Porto, Gláucia, Telma, Cynthia Coelho, Gisele, Gabi, Manuela, Silvinha, Lenora, Patricia Melo, Luiza Barros, Bruno Coelho, Fernando Leão, Silvio Melo, Luciano Domingues, José Ferrer, Cynthia Brito, Lucia Röhr, Dani Couto**.

Being online full time allows you to get online support from many dear Whatsapp groups who must receive my thank you note: **Meninas Douradas, Tagarelas24h, Forever Young; Id, Ego & Superego; Simplesmente 3B; Médicos 2001.1**.

And I hope this final thank you note may perhaps serve as tip for future PhD candidates. To the following essential apps and facebook profiles: **Accuradio** - *the Classical Concentration* channel makes you work at the busiest café; **PhD comics** - puts your spirits high at any time; and **Writing about Writing** - gives valuable tips and puts you back in track with their most needed memes: "You should be writing".

Curriculum Vitae

Brena Melo was born in Santa Barbara, California, USA, in 1977 to a Brazilian couple of PhD students from the University of California. The family returned to Recife, Brazil, before her third birthday. When she was nine years old, the family spent one year in Boston when her parents were admitted to the Massachusetts Institute of Technology (MIT), her father as a Fulbright fellow and mother as special graduate student. The children attended Lawrence School, an award-winning public elementary school in USA. This set of experiences exposed Brena to an environment of excellent educational standards in which knowledge, arts, sports, science and research were enormously prioritized. Still abroad, her little sister broke her arm at school and was admitted to Boston Children's Hospital. Once again, the exposure to an institution of excellence made a life-long impression and awakened her passion for the dynamics of a hospital.

Back to Brazil, after concluding her school years, Brena was admitted to the Universidade Federal de Pernambuco (UFPE) where she graduated from Medical School, in 2001. In the following year she was admitted at the Instituto de Medicina Integral Prof. Fernando Figueira (IMIP) for Obstetrics & Gynecology residency. After concluding this first residency, she decided to widen her clinical scope by attending General Surgery residency at the Serviço de Cirurgia Geral e Transplante Hepático at the Universidade de Pernambuco (UPE) a reference unit in quality of care. After this second residency, she returned to IMIP for her master's degree in Maternal and Child Health in which she studied postpartum outcome in women with preeclampsia according to their uterine artery Doppler. At the same time, she started to work at the Obstetrics Intensive Care Unit and as residency supervisor at the High-Risk Obstetric Ward at the same institution. She also worked at the delivery room of other High-Risk Obstetrics Maternities in the state of Pernambuco: Centro Integrado de Saúde Amaury de Medeiros (CISAM), Hospital Agamenon Magalhães (HAM) and Hospital da Clínicas (HC).

At all these workplaces she was constantly exposed to the daily struggle of the obstetrics assistance typical to low-middle income countries, with constant patient overload and personnel shortage. However, the passion for teaching, sharing and discussing knowledge with peers, residents and students was a constant stimulus to continue her journey as a scholar.

In 2009, Brena spent one month as an observer at St George's Hospital Fetal Medicine Unit exploring partnerships possibilities with IMIP. Her awareness of a standard of excellence in education drove her search and was aligned with IMIP's institutional culture. In her way back home, she was consulted upon her interest in studying medical education at Maastricht University (UM) as result of a partnership between IMIP, its medical school, Faculdade Pernambucana de Saúde (FPS), and the School of Health Professions Education (SHE), at UM, The Netherlands. The opportunity to attend a school of academic excellence was promptly celebrated and the challenge to unveil a new academic field happily accepted. At SHE, she became passionate with instructional design guidelines and its potential practical application in simulation training. She was trained as a scholar and looks forward to locally applying her acquired skills in the training of healthcare professionals.

Brena lives in Recife, Brazil, with her husband, Juliano, and their six years old son, Bernardo.

She Dissertations Series

The SHE Dissertation Series publishes dissertations of PhD candidates from the School of Health Professions Education (SHE) who defended their PhD theses at Maastricht University. The most recent ones are listed below. For more information go to: <https://she.mumc.maastrichtuniversity.nl>

Eppich W. (10-10-2018) Learning through Talk: The Role of Discourse in Medical Education

Wenrich, M. (12-9-2018) Guided Bedside Teaching for Early Learners: Benefits and Impact for Students and Clinical Teachers

Marei, H. (07-09-2018) Application of Virtual Patients in Undergraduate Dental Education

Waterval, D. (26-04-2018) Copy but not paste, an exploration of crossborder medical curriculum partnerships

Smirnova, A. (04-04-2018) Unpacking quality in residency training and health care delivery

Hikspoor, J. (05-12-2017) Development of the heart and vessels in the caudal part of the human body

Boymans, T. (06-10-2017) Hip arthroplasty in the very elderly: anatomical and clinical considerations

Zaidi, Z. (04-10-2017) Cultural hegemony in medical education: exploring the visibility of culture in health professions

Harrison, C. (20-09-2017) Feedback in the context of high-stakes assessment: can summative be formative?

Mekonen, H. (30-06-2017) Development of the axial musculo-skeletal system in humans

Taylor, T. (29-03-2017) Exploring Fatigue as a Social Construct: Implications for Work Hour Reform in Postgraduate Medical Education

McLellan, L. (29-03-2017) Prescribing the right medicine: Perspectives on education and practice

Ignacio, J. (09-02-2017) Stress Management in Crisis Event Simulations for Enhancing Performance

Bolink, S. (19-01-2017) Functional outcome assessment following total hip and knee arthroplasty; Implementing wearable motion sensors

Beckers, J. (09-12-2016) With a little help from my e-portfolio. Supporting students' self-directed learning in senior vocational education

Giroldi, E. (07-12-2016) Towards skilled doctor-patient communication. Putting goal-directed and context-specific communication into (educational) practice

Huwendiek, S. (25-11-2016) Virtual patients for learning of clinical reasoning

Bohle-Carbonell, K. (28-09-2016) May I ask you...? The influence of individual, dyadic & network factors on the emergence of information exchange in teams

Ginsburg, S. (01-09-2016) Hidden in plain sight, the untapped potential of written assessment comments

Koops, W. (08-06-2016) Computer-supported collaborative learning in clinical clerkships

Schlegel, C. (08-06-2016) Simulated and standardized patients in health profession education: the impact of quality improvement

Sorensen, J. (01-06-2016) Obstetric simulation: designing simulation-based medical education and the role of physical fidelity

Kok, E. (01-04-2016) Developing visual expertise: from shades of grey to diagnostic reasoning in radiology

